

eCHT REU 2024

Spectra, Spectral Sequences, and (Co)Fibers smashed with tmf

Irina Bobkova, Jack Carlisle, Emmett Fitz,
Mattie Ji, Peter Kilway, Hillary Kim,
Kolton O'Neal, Jacob Shuckman, Scotty Tilton

June 13, 2024

Note: These are Live \TeX ed notes from the eCHT REU written by Scotty Tilton based on lectures by Irina Bobkova and Jack Carlisle. Scotty may have added a little commentary to add to the exposition, and it's more than likely that Scotty added some of his own errors to the lecture. Please email me at stilton@ucsd.edu if (and when) you find errors. Thanks!

Contents


Day 4	3
Lecture 1	3

Day 4

Lecture 1

Yesterday we computed a homotopy fixed point spectral sequence. Suppose that $C_2 \leadsto \mathcal{E}$ where \mathcal{E} is a spectrum.

$$E_2^{s,t} = H^s(C_2; \pi_t \mathcal{E}) \Rightarrow \pi_{t-s} \mathcal{E}^{hC_2}.$$



The final E_∞ page is $\pi_* E^{hC_2}$. This gives $\pi_* E = \mathbb{Z}[u^{\pm 1}]$ where $|u| = 2$. (If this were a polynomial ring where x is a variable, then u would have the same weight the x^2 “part.”)

Question 1

How do we compute $\pi_*(E^{hC_2} \wedge V(0))$.

There is a spectral sequence

$$\underbrace{E_2^{s,t} = H^s(C_2, \pi_t(E \wedge V(0)))}_{\text{Question 1}} \Rightarrow \underbrace{\pi_{t-s} E^{hC_2} \wedge V(0)}_{\text{Question 2}}$$

For Question 1, we have

$$\mathbb{S} \xrightarrow{2} \mathbb{S} \rightarrow V(0) \quad (\text{fiber sequence})$$

$$E \wedge \mathbb{S} \xrightarrow{2} E \wedge \mathbb{S} \rightarrow E \wedge V(0) \quad (\text{fiber sequence})$$

This induces a long exact sequence in homotopy groups.

$$\begin{array}{ccccccc} & & & & \cdots & \longrightarrow & \pi_{k+1}(E \wedge V(0)) \\ & & & \swarrow & & & \uparrow \\ \pi_k(E \wedge \mathbb{S}) & \xrightarrow{2} & \pi_k(E \wedge \mathbb{S}) & \longrightarrow & \pi_k(E \wedge \mathbb{S}) & & \\ & & \swarrow & & \uparrow & & \\ \pi_{k-1}(E \wedge \mathbb{S}) & \xrightarrow{2} & \cdots & & & & \end{array}$$

After following the LES (using our spectral sequence E_∞ page on the the previous page), we get

$$\pi_t(E \wedge V(0)) = \begin{cases} \mathbb{Z}/2 & t \text{ even} \\ 0 & t \text{ odd} \end{cases}.$$

Therefore, our $E_2^{s,t}$ group we were looking for is coming from

$$H^s(C_2, \pi_t(E \wedge V(0))).$$

$$\begin{array}{ccccccc} & & & & \cdots & \longrightarrow & \pi_3(E \wedge V(0)) \\ & & & \swarrow & & & \uparrow \\ \pi_2(E \wedge \mathbb{S}) & \xrightarrow{u-22u} & \pi_2(E \wedge \mathbb{S}) & \longrightarrow & \pi_2(E \wedge \mathbb{S}) & & \\ & & \swarrow & & \uparrow & & \\ \pi_1(E \wedge \mathbb{S}) & \xrightarrow{2} & \pi_1(E \wedge \mathbb{S}) & \longrightarrow & \pi_1(E \wedge \mathbb{S}) & & \\ & & \swarrow & & \uparrow & & \\ \pi_0(E \wedge \mathbb{S}) & \xrightarrow{2} & \pi_0(E \wedge \mathbb{S}) & & & & \end{array}$$

Definition 1: pullbacks

Let \mathcal{C} be a category which contains the diagram

$$\begin{array}{ccc} & Y & \\ & \downarrow g & \\ X & \xrightarrow{f} & Z \end{array}$$

A **pullback** of this diagram W is 3 pieces of information

- An object $W \in \text{ob}(\mathcal{C})$
- A map $W \xrightarrow{p_1} X$
- A map $W \xrightarrow{p_2} Y$

such that

- The diagram $\begin{array}{ccc} W & \xrightarrow{p_2} & Y \\ p_1 \downarrow & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$ commutes^a

- If someone hands you a commutative diagram $\begin{array}{ccc} A & \xrightarrow{h_2} & Y \\ h_1 \downarrow & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$, then there is a *UNIQUE* map \tilde{h} such that

$$\begin{array}{ccccc} A & & & & \\ & \searrow \tilde{h} & & & \\ & & W & \xrightarrow{p_2} & Y \\ & \swarrow h_1 & \downarrow p_1 & \lrcorner & \downarrow g \\ & & X & \xrightarrow{f} & Z \end{array}$$

^b

Let \mathcal{C} be a category with a subcategory \mathcal{D} . We say \mathcal{D} is **closed under pullbacks** by morphisms in \mathcal{C} if for all

arrows $X \xrightarrow{f} Z$ in \mathcal{D} and for all $Y \xrightarrow{g} Z$ in \mathcal{C} such that we can form the pullback $\begin{array}{ccc} W & \xrightarrow{p_1} & Y \\ \downarrow p_2 & \lrcorner & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$, then the arrow

$W \xrightarrow{p_1} Z$ is in \mathcal{D} .

^a“ \lrcorner ” is a ‘long’ hand for commutes, and people usually suppress it from notation.

^bAs a shorthand, people usually write the pullback like this:

$$\begin{array}{ccc} W & \longrightarrow & Y \\ \downarrow & \lrcorner & \downarrow \\ X & \longrightarrow & Z \end{array}$$

Definition 2: pushouts

Let \mathcal{C} be a category which contains the diagram

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ \downarrow f & & \\ X & & \end{array}.$$

A **pushout** of this diagram W is 3 pieces of information

- An object $W \in \text{ob}(\mathcal{C})$
- A map $X \xrightarrow{i_1} W$
- A map $Y \xrightarrow{i_2} W$

such that

- The diagram $\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ \downarrow f & & \downarrow i_2 \\ X & \xrightarrow{i_1} & W \end{array}$ commutes

- If someone hands you a commutative diagram $\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ \downarrow f & & \downarrow \ell_2 \\ X & \xrightarrow{\ell_1} & A \end{array}$, then there is a *UNIQUE* map $\tilde{\ell}$ such that

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ \downarrow f & & \downarrow i_2 \\ X & \xrightarrow{i_1} & W \end{array} \quad \begin{array}{c} \searrow \ell_2 \\ \downarrow \tilde{\ell} \\ \searrow \ell_1 \end{array} \quad \begin{array}{c} \\ \\ A \end{array}.$$

^a

^aAs a shorthand, people usually write the pushout like this:

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ \downarrow f & \lrcorner & \downarrow i_2 \\ X & \xrightarrow{i_1} & W \end{array}$$

Definition 3: (Homotopy) Fixed points

Let $G \curvearrowright X$ where X is a topological space. The **fixed points** of X are

Bibliography

Index

closed under pullbacks, 5

fixed points, 6

pullback, 5

pushout, 6