

## Theorems related to Vector Calculus

### Question 1:

#### part 1a:

Evaluate the scalar line integral:

$$\int_C (6x^2 + 4y) ds \quad \text{where} \quad \mathbf{r}_C(t) = \begin{bmatrix} 3t + 2 \\ -t + 3 \end{bmatrix} \quad \text{and } t \in [-1, 1]$$

#### part 1b:

Evaluate the scalar line integral:

$$\int_C (x^2 + 2y) ds \quad \text{where} \quad \mathbf{r}_C(t) = \begin{bmatrix} 1 - 2t \\ 2 - t \end{bmatrix} \quad \text{and } t \in [0, 3]$$

#### part 1c:

Evaluate the scalar line integral:

$$\int_C 2x \cdot \sin(y) \cdot ds \quad \text{where} \quad \mathbf{r}_C(t) = \begin{bmatrix} \cos(t) \\ t \end{bmatrix} \quad \text{and } t \in [0, \pi/2]$$

### Question 2:

#### part 2a:

Evaluate the vector line integral:

$$\int_C \begin{bmatrix} -6x \\ 4y \end{bmatrix} \cdot d\mathbf{r} \quad \text{where} \quad \mathbf{r}_C(t) = \begin{bmatrix} t^2 \\ 1/t \end{bmatrix} \quad \text{and } t \in [1, 2]$$

#### part 2b:

Evaluate the vector line integral:

$$\int_C \begin{bmatrix} y \\ x \end{bmatrix} \cdot d\mathbf{r} \quad \text{where} \quad \mathbf{r}_C(t) = \begin{bmatrix} t^3 \\ t^4 \end{bmatrix} \quad \text{and } t \in [0, 2]$$

**part 2c:**

Evaluate the vector line integral:

$$\int_C \begin{bmatrix} 2x \\ y^2 \\ -z \end{bmatrix} \cdot d\mathbf{r} \quad \text{where} \quad \mathbf{r}_C(t) = \begin{bmatrix} -t^2 \\ t+4 \\ 3t+2 \end{bmatrix} \quad \text{and } t \in [-1, 1]$$

**Question 3:**

**part 3a:**

Is the vector field  $\mathbf{F}(x, y, z) = \begin{bmatrix} y^2 \sin(z) \\ 2xy \sin(z) \\ xy^2 \cos(z) \end{bmatrix}$  conservative? If yes, use the **gradient theorem** to evaluate the vector line integral:

$$\int_C \begin{bmatrix} y^2 \sin(z) \\ 2xy \sin(z) \\ xy^2 \cos(z) \end{bmatrix} \cdot d\mathbf{r} \quad \text{where} \quad \mathbf{r}_{\text{initial}} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad \text{and} \quad \mathbf{r}_{\text{final}} = \begin{bmatrix} 1 \\ 1 \\ \pi/2 \end{bmatrix}$$

**part 3b:**

Is the vector field  $\mathbf{F}(x, y, z) = \begin{bmatrix} xy \\ xy \\ z \end{bmatrix}$  conservative? If yes, use the **gradient theorem** to evaluate the vector line integral:

$$\int_C \begin{bmatrix} xy \\ xy \\ z \end{bmatrix} \cdot d\mathbf{r} \quad \text{where} \quad \mathbf{r}_{\text{initial}} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad \text{and} \quad \mathbf{r}_{\text{final}} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

**Question 4:**

**part 4a:**

The region  $\sigma$  on the right is

$$\sigma = \{(x, y) | 0 \leq x \leq 6 \text{ and } 0 \leq y \leq 3 - (1/2)x\}$$


Use Green's theorem to compute the loop integral

$$\int_{\partial\sigma} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r}$$

where  $\partial\sigma$  is the counterclockwise oriented boundary of  $\sigma$ .

Next, compute each of  $\int_{C_1} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r}$ ;  $\int_{C_2} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r}$ ; and  $\int_{C_3} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r}$  and show that

$$\int_{\partial\sigma} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r} = \int_{C_1} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r} + \int_{C_2} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r} + \int_{C_3} \begin{bmatrix} 3x - 5y \\ x - 2y \end{bmatrix} \cdot d\mathbf{r}$$



**part 4b:**


The region  $\sigma$  on the right is

$$\sigma = \{(x, y) | 0 \leq x \leq 1 \text{ and } x^3 \leq y \leq x\}$$

Use Green's theorem to compute the loop integral

$$\int_{\partial\sigma} \begin{bmatrix} 2xy \\ x+y \end{bmatrix} \cdot d\mathbf{r}$$

where  $\partial\sigma$  is the counterclockwise oriented boundary of  $\sigma$ .



Next, compute each of  $\int_{C_1} \begin{bmatrix} 2xy \\ x+y \end{bmatrix} \cdot d\mathbf{r}$ ; and  $\int_{C_2} \begin{bmatrix} 2xy \\ x+y \end{bmatrix} \cdot d\mathbf{r}$  and show that

$$\int_{\partial\sigma} \begin{bmatrix} 2xy \\ x+y \end{bmatrix} \cdot d\mathbf{r} = \int_{C_1} \begin{bmatrix} 2xy \\ x+y \end{bmatrix} \cdot d\mathbf{r} + \int_{C_2} \begin{bmatrix} 2xy \\ x+y \end{bmatrix} \cdot d\mathbf{r} + \int_{C_3} \begin{bmatrix} 2xy \\ x+y \end{bmatrix} \cdot d\mathbf{r}$$

**part 4c:**


The region  $\sigma$  on the right is

$$\sigma = \{(x, y) | -4 \leq x \leq 0 \text{ and } 0 \leq y \leq 2 + (1/2)x\}$$

Use Green's theorem to compute the loop integral

$$\int_{\partial\sigma} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r}$$

where  $\partial\sigma$  is the counterclockwise oriented boundary of  $\sigma$ .



Next, compute each of  $\int_{C_1} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r}$ ;  $\int_{C_2} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r}$ ; and  $\int_{C_3} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r}$  and show that

$$\int_{\partial\sigma} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r} = \int_{C_1} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r} + \int_{C_2} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r} + \int_{C_3} \begin{bmatrix} x+2y \\ 5x+y \end{bmatrix} \cdot d\mathbf{r}$$

### Question 5:


The region  $\sigma$  on the right is

$$\sigma = \{(x, y) | 0 \leq x \leq 1 \text{ and } 0 \leq y \leq 1 - x^2\}$$

Use Gauss's divergence theorem to compute the flux integral

$$\int_{\partial\sigma} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix}$$

where  $\partial\sigma$  is the counterclockwise oriented boundary of  $\sigma$ .



Next, compute each of  $\int_{C_1} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix}$ ;  $\int_{C_2} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix}$ ; and  $\int_{C_3} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix}$  and show that

$$\int_{\partial\sigma} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix} = \int_{C_1} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix} + \int_{C_2} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix} + \int_{C_3} \begin{bmatrix} x^2 \\ y \end{bmatrix} \cdot \begin{bmatrix} dy \\ -dx \end{bmatrix}$$