

Omega-3 Free Fatty Acids for the Maintenance of Remission in Crohn Disease

The EPIC Randomized Controlled Trials

Brian G. Feagan, MD

William J. Sandborn, MD

Ulrich Mittmann, MD

Simon Bar-Meir, MD

Geert D'Haens, MD, PhD

Marc Bradette, MD

Albert Cohen, MD

Chrystian Dallaire, MD

Terry P. Ponich, MD

John W. D. McDonald, MD, PhD

Xavier Hébuterne, MD, PhD

Pierre Paré, MD

Pavel Klvana, MD

Yaron Niv, MD

Sandro Ardizzone, MD

Olga Alexeeva, MD

Alaa Rostom, MD

Gediminas Kiudelis, MD

Johannes Spleiss, MSc

Denise Gilgen, PhD

Margaret K. Vandervoort, MSc

Cindy J. Wong, MSc

Guang Yong Zou, PhD

Allan Donner, PhD

Paul Rutgeerts, MD, PhD

EFFECTIVE THERAPY TO MAINTAIN remission in Crohn disease is an unmet medical need. Although formulations of 5-aminosalicylic acid agents remain widely used for this purpose, strong evidence exists that they are not effective.¹

See also Patient Page.

Context Maintenance therapy for Crohn disease features the use of immunosuppressive drugs, which are associated with an increased risk of infection. Identification of safe and effective maintenance strategies is a priority.

Objective To determine whether the oral administration of omega-3 free fatty acids is more effective than placebo for prevention of relapse of Crohn disease.

Design, Setting, and Patients Two randomized, double-blind, placebo-controlled studies (Epanova Program in Crohn's Study 1 [EPIC-1] and EPIC-2) conducted between January 2003 and February 2007 at 98 centers in Canada, Europe, Israel, and the United States. Data from 363 and 375 patients with quiescent Crohn disease were evaluated in EPIC-1 and EPIC-2, respectively.

Interventions Patients with a Crohn's Disease Activity Index (CDAI) score of less than 150 were randomly assigned to receive either 4 g/d of omega-3 free fatty acids or placebo for up to 58 weeks. No other treatments for Crohn disease were permitted.

Main Outcome Measure Clinical relapse, as defined by a CDAI score of 150 points or greater and an increase of more than 70 points from the baseline value, or initiation of treatment for active Crohn disease.

Results For EPIC-1, 188 patients were assigned to receive omega-3 free fatty acids and 186 patients to receive placebo. Corresponding numbers for EPIC-2 were 189 and 190 patients, respectively. The rate of relapse at 1 year in EPIC-1 was 31.6% in patients who received omega-3 free fatty acids and 35.7% in those who received placebo (hazard ratio, 0.82; 95% confidence interval, 0.51-1.19; $P=.30$). Corresponding values for EPIC-2 were 47.8% and 48.8% (hazard ratio, 0.90; 95% confidence interval, 0.67-1.21; $P=.48$). Serious adverse events were uncommon and mostly related to Crohn disease.

Conclusion In these trials, treatment with omega-3 free fatty acids was not effective for the prevention of relapse in Crohn disease.

Trial Registration clinicaltrials.gov Identifiers: EPIC-1: NCT00613197, EPIC-2: NCT00074542

JAMA. 2008;299(14):1690-1697

www.jama.com

Immunosuppressive agents such as purine antimetabolites,^{2,3} methotrexate,⁴ and tumor necrosis factor α antagonists⁵⁻⁸ are moderately effective for maintaining remission; however, their use is associated with an increased risk of infection.⁹⁻¹¹ Therefore, development of a safe, inexpensive, and effective orally administered agent is a research priority.

Omega-3 free fatty acids are anti-inflammatory substances found in marine fish that have several health benefits. These compounds have been used to treat inflammatory disorders such as rheumatoid

Author Affiliations are listed at the end of this article.

Corresponding Author: Brian G. Feagan, MD, Robarts Clinical Trials, Robarts Research Institute, University of Western Ontario, 100 Perth Dr, London, ON, Canada N6A 5K8 (bfeagan@robarts.ca).

arthritis^{12,13} and IgA nephropathy.¹⁴ Additionally, administration of omega-3 free fatty acids reduces the concentration of serum triglycerides in patients with dyslipidemia¹⁵⁻¹⁷ and decreases the risk of death from cardiac disease.¹⁸⁻²⁰ However, previous clinical trials that evaluated omega-3 free fatty acids for maintenance of remission in Crohn disease have yielded inconsistent results. Belluzzi et al²¹ demonstrated a 33% absolute reduction in the 1-year risk of relapse in patients treated with 2.7 g/d of omega-3 free fatty acids compared with placebo. In contrast, no benefit was observed in a second trial in which patients received 5 g/d of highly concentrated omega-3 free fatty acids.²² Based on these observations, we conducted 2 large-scale trials of high-dose omega-3 free fatty acids as maintenance therapy in patients with quiescent Crohn disease. Two independent trials were performed, in complementary populations of patients, to meet regulatory expectations.

METHODS

These multicenter, randomized, double-blind, placebo-controlled studies were conducted in Canada, Europe, Israel, and the United States between January 2003 and February 2007. Fifty-one sites participated in Epanova Program in Crohn's Study 1 (EPIC-1), and 47 sites participated in EPIC-2. The institutional review board at each site approved the protocols. All patients provided written informed consent.

Eligible patients were adults with a confirmed diagnosis of Crohn disease and a Crohn's Disease Activity Index (CDAI)²³ score of less than 150 points. In EPIC-1, patients were eligible if they had experienced a disease exacerbation within the past year and were in remission for at least 3 months but not longer than 12 months. In EPIC-2, patients with active disease were treated with a standardized 16-week tapering course of either prednisone²⁴ or budesonide²⁵ at initial doses of 40 mg once daily and 9 mg once daily, respectively. Eight weeks after the initiation of corticosteroid treatment, the CDAI score was assessed; if it was less than

150 points, the patient was eligible for randomization. At this time, patients were receiving either 20 mg/d of prednisone or 6 mg/d of budesonide.

Patients were not eligible if they were allergic or intolerant of omega-3 free fatty acids or fish products; if they required treatment with 5-aminosalicylates, immunosuppressives, tumor necrosis factor α antagonists, or investigational drugs; had an ostomy or short bowel syndrome; or if they had a bowel resection in the preceding 3 months or had clinically important bowel obstruction (defined by the presence of proximal dilatation demonstrated radiographically) in the 3 months prior to randomization. Patients with malignancy (with the exception of resected basal or squamous cell skin cancer or carcinoma of the cervix in situ), those with substance abuse, or those with severe medical diseases other than Crohn disease were not included.

Study Design

Eligible patients were randomly assigned, in a 1:1 ratio using a random number generator, to receive four 1-g gelatin capsules of omega-3 free fatty acids (Epanova; Tillotts Pharma AG, Ziefen, Switzerland) or 4 capsules of identical-appearing placebo each day for 52 weeks in EPIC-1 or for 58 weeks in EPIC-2. Epanova consists of 50% to 60% eicosapentanoic acid and 15% to 25% docosahexanoic acid as a free fatty acid encapsulated in a delayed-release soft gelatin capsule. The placebo capsule consisted of 1 g of medium-chain triglyceride oil. To enhance tolerability, patients began with an initial dose of 1 g/d of the study drug for the first 7 days; this was increased to 2 g/d daily for 7 days and then further increased to the final dose of 4 g/d, administered as 2 capsules twice daily.

Both studies used a central randomization procedure that was stratified by the induction therapy received prior to randomization or the duration of remission (For EPIC-1: 5-aminosalicylates; corticosteroids and in remission for 3 to 6 months; corticosteroids and in remission for 6 to 12 months; or im-

munosuppressives. For EPIC-2: prednisone or budesonide for induction therapy). Neither the investigators nor the patients were aware of the treatment assignment.

No new medications for Crohn disease were allowed following randomization. Participants in EPIC-2 had the dose of corticosteroid tapered. Those receiving budesonide decreased the daily dose to 3 mg by week 4, with discontinuation at week 8. Patients receiving prednisone decreased the daily dose by 2.5 mg/wk, with discontinuation at week 8. An increase in the dose of corticosteroid to control symptoms was considered a failure of treatment.

Follow-up Schedule

Patients were evaluated in the clinic at weeks -1, 0 (randomization), 4, 16, 30, 44, and 52 for EPIC-1 and at weeks -8, 0 (randomization), 2, 8, 16, 30, 44, and 58 for EPIC-2. At each visit patients were given a physical examination, blood was drawn for biochemical testing, and results of the CDAI and 36-Item Short-Form Health Survey (SF-36)²⁶ were assessed.

Outcome Measures

The primary efficacy outcome was the occurrence of a relapse, defined as either an increase of the CDAI score to 150 points or greater and an increase of more than 70 points from the score calculated at the randomization visit, or the initiation of prohibited medical therapy or surgery for symptomatic active Crohn disease.⁴ Secondary outcomes were the change in mean scores on the CDAI and SF-36, the change in the mean serum triglyceride concentration, and the occurrence of adverse events, which were classified using the Medical Dictionary for Regulatory Activities.²⁷


Adherence to the study drug as assigned was evaluated at each clinic visit by interviewing the patient and by capsule counts. Patients were considered adherent if they received at least 75% of the capsules between 2 clinic visits, with no interruption of medication for more than 14 consecutive days.

Statistical Analysis

Baseline characteristics were analyzed using the *t* test or the χ^2 test as appropriate. For the primary analysis the time to a re-

lapse was compared by treatment group using the log-rank test. Kaplan-Meier plots were used to depict the proportion of patients remaining in remission over time.

Figure 1. Disposition of Patients During the Conduct of EPIC-1 and EPIC-2



EPIC indicates Epanova Program in Crohn's Study.

Hazard ratios (HRs) and associated confidence intervals (CIs) were obtained using Cox regression with treatment assignment as the only independent variable. For EPIC-1, a prespecified subgroup analysis was also conducted to evaluate the primary outcome in patients whose remission had been induced by corticosteroids. A Bonferroni correction ($P \leq .025$) was applied for this comparison. The mean changes in CDAI, SF-36 scores, and serum triglyceride concentrations were compared by the *t* test. Point estimates and 95% CIs of the between-group difference in the proportion of patients with adverse events were obtained. The proportion of patients compliant with the study drug was compared using the χ^2 test.

All analyses were performed according to a modified intent-to-treat principle in which data from all patients who received at least 1 dose of the study drug and who provided any postrandomization data were analyzed. A 2-sided *P* value less than .05 was considered statistically significant. SAS version 8.2²⁸ was used to perform these analyses; however, CIs were estimated using StatXact version 8.0.²⁹

Both trials were designed to detect a 15%³⁰ reduction in relapse rate with 80% power at the .05 significance level. We estimated that the relapse rate in the placebo group would be 70% in EPIC-1.²¹ For EPIC-2, we estimated that 80% of the patients would achieve clinical remission at the discontinuation of corticosteroid induction therapy and that the rate of relapse in the patients who were subsequently assigned to placebo would be 80%.³¹ Accordingly, a total of 306 patients and 309 patients were required to detect an absolute difference of 15% in the relapse rate for EPIC-1 and EPIC-2, respectively. We planned to randomize 360 patients in EPIC-1 and 364 patients in EPIC-2 to allow for a non-evaluable rate of 15%.

RESULTS

The disposition of the patients is shown in FIGURE 1. In EPIC-1, 383 patients were randomized. During the conduct of EPIC-1, it was discovered that a single site had not conducted the trial according to International Conference on Har-

monisation E6 guidelines³² and had not followed the protocol on numerous occasions. Following consultation with regulatory agencies, all of the data collected on the 9 patients randomized from this site were excluded from analysis. Of the remaining 374 patients, 2 did not receive study drug, 8 had no postrandomization relapse assessment, and 1 was re-randomized in error. Thus, a total of 363 patients were included in the modified intent-to-treat population; of these, 183 received omega-3 free fatty acids and 180 received placebo. In EPIC-2, 379 patients were randomized, 2 did not receive study drug, and 2 had no postrandomization assessment. Thus, a total of 375 patients were included in the modified intent-to-treat population; of these, 187 received omega-3 free fatty acids and 188 received placebo.

In EPIC-1, 26 patients who did not experience relapse discontinued treatment prematurely in the omega-3 free fatty acids group compared with 29 in the placebo group ($P = .61$). In EPIC-2, 30 patients who did not experience relapse discontinued treatment prematurely in the omega-3 free fatty acids group compared with 18 in the placebo group ($P = .06$). No important differences were observed between the treatment groups in the reasons for withdrawal in either trial (Figure 1).

The baseline characteristics of the patients were similar for both treatment groups in each trial (TABLE 1).³³

Primary Outcome

FIGURE 2 shows the proportion of patients who remained in remission over time. No significant differences were ob-

served between the 2 treatment groups in either trial. In EPIC-1, 54 patients treated with omega-3 free fatty acids and 62 patients treated with placebo experienced a clinical relapse. The proportion of patients assigned to receive omega-3 free fatty acids who experienced a relapse within 360 days was estimated to be 31.6%, compared with 35.7% for those who received placebo (HR, 0.82; 95% CI, 0.57-1.19; $P = .30$). In EPIC-2, 84 patients treated with omega-3 free fatty acids and 94 patients treated with placebo experienced a clinical relapse. The proportion of patients assigned to receive omega-3 free fatty acids who experienced a relapse within 360 days was estimated to be 47.8%, compared with 48.8% of those who received placebo (HR, 0.90; 95% CI, 0.67-1.21; $P = .48$).

Table 1. Baseline Characteristics of the Study Patients in EPIC-1 and EPIC-2^a

Characteristic	EPIC-1			EPIC-2		
	Omega-3 Free Fatty Acids (n = 183)	Placebo (n = 180)	P Value	Omega-3 Free Fatty Acids (n = 187)	Placebo (n = 188)	P Value
Characteristic						
Age, mean (SD), y	40.5 (15.2)	38.2 (13.1)	.12	38.5 (13.8)	40.0 (13.6)	.28
Male sex, No. (%)	88 (48.1)	74 (41.1)	.18	81 (43.3)	79 (42.0)	.80
Weight, mean (SD), kg	69.0 (13.2)	67.9 (13.8)	.42	69.8 (15.5)	68.4 (16.1)	.40
Current smoker, No. (%)	56 (30.6)	62 (34.4)	.43	47 (25.1)	70 (37.2)	.01
Remission induction therapy, No. (%)						
Oral 5-ASA or antibiotics	82 (44.8)	83 (46.1)	.97			
Steroids, in remission 3-6 mo	57 (31.1)	57 (31.7)				
Steroids, in remission >6-12 mo	36 (19.7)	32 (17.8)				
Immunosuppressives	8 (4.4)	8 (4.4)				
Prednisone				106 (56.7)	110 (58.5)	.72
Budesonide				81 (43.3)	78 (41.5)	
Harvey-Bradshaw index, mean (SD) ^b	2.3 (2.0)	2.3 (2.3)	.82	6.2 (4.2)	6.0 (3.6)	.58
Crohn's Disease Activity Index, mean (SD) ^c	79.6 (40.7)	75.4 (43.5)	.34	77.4 (44.8)	78.0 (41.6)	.90
Months since diagnosis						
Mean (SD)	85.7 (106.3)	82.7 (98.9)	.78	84.6 (94.0)	90.6 (108.5)	.57
Median (range)	42.0 (4-540)	42.5 (1-465)		51.0 (1-427)	42.0 (2-482)	
Previous surgery for Crohn disease, No. (%)	61 (33.3)	64 (35.6)	.66	56 (29.9)	67 (35.6)	.24
Crohn disease therapy prior 12 mo, No. (%)						
Oral 5-ASA therapy	142 (77.6)	141 (78.3)	.87	102 (54.5)	90 (47.9)	.20
Topical rectal therapy	9 (4.9)	8 (4.4)	.83	5 (2.7)	11 (5.9)	.13
Systemic corticosteroids	97 (53.0)	94 (52.2)	.88	39 (20.9)	36 (19.1)	.68
Immune-modifying agents	12 (6.6)	15 (8.3)	.52	13 (7.0)	13 (6.9)	.99
Immune modifiers/biologics	3 (1.6)	2 (1.1)	.67	4 (2.1)	3 (1.6)	.70
Antibiotic therapy	37 (20.2)	32 (17.8)	.55	23 (12.3)	25 (13.3)	.77

Abbreviations: ASA, aminosaliclate; EPIC, Epanova Program in Crohn's Study.

^aValues for comparison of continuous outcomes were derived by the *t* test. Pearson χ^2 test was used to compare categorical variables.

^bHarvey-Bradshaw scores range from 0 to approximately 14. Higher scores indicate worse disease activity; scores of ≤ 4 are associated with clinical remission. Baseline score for

EPIC-2 was assessed at the screening visit prior to steroid tapering.

^cScores of ≤ 150 points are associated with quiescent disease, scores > 150 points indicate active disease, and scores > 450 points are seen with extremely severe disease.

In the subgroup of patients in EPIC-1 who had received induction therapy with corticosteroids, no significant difference was observed between the 2 treatment groups. Thirty-eight percent of those who received omega-3 free fatty acids experienced a relapse at 360

days compared with 39.5% in the placebo group ($P = .60$).


Disease Activity and Quality of Life

No important differences were observed in the changes in mean CDAI and SF-36 scores in either study (data not shown).

Adherence to Study Drug

In EPIC-1, 79.2% of patients who received omega-3 free fatty acids had adequate adherence to study drug, compared with 75.6% of those who received placebo ($P = .40$). The corresponding values for EPIC-2

Figure 2. Kaplan-Meier Estimates of the Time to Relapse in the Omega-3 Free Fatty Acids and Placebo Groups for EPIC-1 and EPIC-2



EPIC indicates Epanova Program in Crohn's Study.

Table 2. Incidence of Adverse Events, EPIC-1 and EPIC-2^a

Adverse Event	No. (%)					
	EPIC-1			EPIC-2		
	Omega-3 Free Fatty Acids (n = 187)	Placebo (n = 184)	Difference (95% CI) ^b	Omega-3 Free Fatty Acids (n = 189)	Placebo (n = 188)	Difference (95% CI) ^b
Crohn disease worsening	47 (25.1)	57 (31.0)	-5.8 (-14.9 to 3.3)	78 (41.3)	89 (47.3)	-6.1 (-16.0 to 4.0)
Abdominal pain	45 (24.1)	41 (22.3)	1.8 (-6.8 to 10.4)	65 (34.4)	62 (33.0)	1.4 (-8.1 to 10.9)
Diarrhea	35 (18.7)	21 (11.4)	7.3 (0 to 14.7)	44 (23.3)	37 (19.7)	3.6 (-4.7 to 11.9)
Arthralgia	21 (11.2)	15 (8.2)	3.1 (-3.1 to 9.3)	29 (15.3)	28 (14.9)	0.5 (-6.9 to 7.8)
Nasopharyngitis	18 (9.6)	13 (7.1)	2.6 (-3.2 to 8.5)	29 (15.3)	27 (14.4)	1.0 (-6.3 to 8.3)
Nausea	17 (9.1)	4 (2.2)	6.9 (2.4 to 12.2)	30 (15.9)	19 (10.1)	5.8 (-1.0 to 12.7)
Headache	12 (6.4)	10 (5.4)	1.0 (-4.1 to 6.1)	12 (6.3)	20 (10.6)	-4.3 (-10.2 to 1.4)
Influenza	12 (6.4)	14 (7.6)	-1.2 (-6.7 to 4.2)	6 (3.2)	7 (3.7)	-0.5 (-4.7 to 3.5)
Flatulence	11 (5.9)	13 (7.1)	-1.2 (-6.5 to 4.0)	12 (6.3)	12 (6.4)	6.3 (-5.2 to 5.2)
Gastroenteritis	10 (5.3)	10 (5.4)	-0.1 (-5.0 to 4.8)	6 (3.2)	8 (4.3)	-1.1 (-5.4 to 3.0)
Fatigue	10 (5.3)	6 (3.3)	2.1 (-2.3 to 6.7)	22 (11.6)	22 (11.7)	-0.1 (-6.7 to 6.6)
Dysgeusia	2 (1.1)	1 (0.5)	0.5 (-2.0 to 3.3)	10 (5.3)	2 (1.1)	4.2 (0.8 to 8.5)
Dyspepsia	4 (2.1)	5 (2.7)	-0.6 (-4.3 to 3.0)	14 (7.4)	13 (6.9)	-0.5 (-4.9 to 6.0)
Hematochezia	0	2 (1.1)	-1.1 (-3.9 to 0.9)	9 (4.8)	10 (5.3)	-0.6 (-5.3 to 4.2)
Sinusitis	1 (0.5)	3 (1.6)	-1.1 (-4.2 to 1.5)	3 (1.6)	15 (8.0)	-6.4 (-11.4 to -2.3)
Vomiting	4 (2.1)	2 (1.1)	1.1 (-2.0 to 4.4)	13 (6.9)	11 (5.9)	1.0 (-4.1 to 6.3)

Abbreviations: CI, confidence interval; EPIC, Epanova Program in Crohn's Study.

^a Table shows the proportion of patients experiencing adverse events by more than 5% of the patients in any treatment group in EPIC-1 or EPIC-2. Seven pregnancies occurred during the trials; 2 occurred in patients who received omega-3 free fatty acids and 5 in those who received placebo.

^b 95% CIs obtained using inversion of 2-sided χ^2 tests.

were 75.4% and 81.4%, respectively ($P=.16$).

Serum Triglycerides

A significant decrease in serum triglyceride concentration was observed in patients assigned to receive omega-3 free fatty acids. In EPIC-1, the mean decrease after 30 weeks of treatment was -21.5 mg/dL (to convert to millimoles per liter, multiply by 0.0113), compared with a mean increase of 16.5 mg/dL in patients who received placebo ($P<.001$). Corresponding values for EPIC-2 were -27.1 mg/dL in patients treated with omega-3 free fatty acids, compared with -5.1 mg/dL in patients receiving placebo ($P=.02$).

Adverse Events

No important differences were observed between the treatment groups in the incidence of adverse events (TABLE 2). The overall frequency of adverse events was higher in EPIC-2 because these patients had active disease at the time of entry and were receiving treatment with corticosteroids. The omega-3 free fatty acid formulation was well tolerated, and previously observed adverse events such as diarrhea and bloating²¹ were uncommon. No differences in laboratory results were identified apart from the reduced triglyceride concentration observed in patients who received omega-3 free fatty acids.

COMMENT

These 2 large-scale trials had similar results. No beneficial effect of high-dose omega-3 free fatty acid treatment was observed for the prevention of relapse over 1 year of follow-up. These findings are distinctly different from those of Belluzzi et al,²¹ who reported a 33% absolute reduction in the rate of relapse following treatment with a similar omega-3 free fatty acid formulation in a randomized placebo-controlled trial that evaluated 78 patients.

There are 3 possible explanations for these discordant results. First, the design of the study by Belluzzi et al²¹ differed from the EPIC trials in that only patients at a high risk of relapse, de-

finied by the presence of an elevated serum concentration of acute-phase reactants, were included. It could be speculated that the patients evaluated in the EPIC studies were at such low risk that the trials lacked sufficient statistical power to detect the prespecified 15% benefit of treatment. While this hypothesis might be plausible in the case of EPIC-1, given that the 1-year placebo relapse rate was only 35.7%, it is not tenable for EPIC-2, in which 51.2% of patients relapsed. Furthermore, our results are consistent with those of Lorenz-Meyer et al,²² who also demonstrated no benefit of omega-3 fatty acids in a placebo-controlled relapse prevention study that evaluated 204 patients.

A second possible explanation concerns the different formulations used in the studies. The formulation evaluated in the study by Belluzzi et al was a hard gelatin capsule, in contrast to the soft gelatin capsule used in the current trials. However, pharmacokinetic studies have shown that a similar concentration of omega-3 free fatty acids is incorporated into cell membranes following administration of either preparation (J.S., written communication, January 2004, data on file). Moreover, although no beneficial effect was observed for the prevention of relapse, a statistically significant decrease in the serum concentration of triglycerides was observed in patients who received omega-3 free fatty acids. This finding, which is consistent with our analysis of the capsule counts, indicates the patients had adequate exposure to the study drug and that poor compliance cannot account for the negative results.

Finally, we believe that the most plausible explanation for the discordant findings is that the study by Belluzzi et al, which only evaluated 78 patients, was incorrect in concluding that treatment with omega-3 free fatty acids is beneficial. Several studies have demonstrated that small clinical trials are more susceptible to bias, are often less methodologically rigorous, and produce relatively less precise estimates of the true effect of treatment than large, multicenter studies.³⁴⁻³⁶

Our results are important because the use of alternative medicines in general, and omega-3 free fatty acid formulations in particular, is widespread among patients with inflammatory bowel disease.³⁷⁻⁴¹ This may be due, in part, to dissemination of the positive results obtained in the trial by Belluzzi et al. Given the negative results observed in the EPIC trials and in the trial by Lorenz-Meyer et al,²² we do not endorse this practice, since patients with Crohn disease who are at risk for relapse would be better served by taking medications of known efficacy.

The omega-3 free fatty acid formulation was well tolerated. In contrast to the results obtained in the study by Belluzzi et al,²¹ during the first month of treatment only 2.7% of patients had to discontinue the drug due to diarrhea. The increased tolerability was likely due to the gradual increase in dosing specified by the protocols. Although a theoretical risk of bleeding due to platelet dysfunction exists with omega-3 free fatty acid treatment,^{42,43} no serious bleeding problems were encountered despite the administration of a relatively high drug dose.

In summary, in the EPIC-1 and EPIC-2 trials administration of a high dose of omega-3 free fatty acids did not reduce the rate of relapse in patients with quiescent Crohn disease.

Author Affiliations: Robarts Clinical Trials, Robarts Research Institute (Drs Feagan, Zou, and Donner and Mss Vandervoort and Wong) and Department of Epidemiology and Biostatistics (Drs Feagan, Zou, and Donner), University of Western Ontario, London, Ontario, Canada; Mayo Clinic, Rochester, Minnesota (Dr Sandborn); MP Consulting, Basel, Switzerland (Dr Mittmann); Sheba Medical Center, Tel Hashomer, Israel (Dr Bar-Meir); Imelda General Hospital, Imeldalaan Bonheiden, Belgium (Dr D'Haens); Pavillon Hôtel-Dieu de Québec, Québec City, Québec, Canada (Dr Bradette); Jewish General Hospital, Montreal, Québec, Canada (Dr Cohen); Pavillon Saint-François d'Assise, Québec City (Dr Dallaire); London Health Sciences Centre, London, Ontario, Canada (Drs Ponich and McDonald); Centre Hospitalier Universitaire, University of Nice Sophia Antipolis, Nice, France (Dr Hébuterne); Hôpital Saint-Sacrement, Québec City (Dr Paré); University Hospital, Ostrava, Czech Republic (Dr Klvana); Rabin Medical Center, Petah Tiqva, Israel (Dr Niv); Ospedale Luigi Sacco, Milan, Italy (Dr Ardizzone); Regional Clinical Hospital, Nizhny Novgorod, Russia (Dr Alexeeva); University of Calgary, Calgary, Alberta, Canada (Dr Rostom); Kaunas Medical University, Kaunas, Lithuania (Dr Kiudelis); Tillotts Pharma AG, Ziefen, Switzerland (Mr Speiss and Dr Gilgen); and University Hospital Gasthuisberg, Herestraat, Leuven, Belgium (Dr Rutgeerts).

Author Contributions: Dr Feagan had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Feagan, Sandborn, Mittmann, Niv, Vandervoort, Rutgeerts.

Acquisition of data: Sandborn, Bar-Meir, D'Haens, Bradette, Cohen, Dallaire, Ponich, McDonald, Hébuterne, Klvana, Ardizzone, Alexeeva, Rostom, Kiudelis, Spleiss, Gilgen.

Analysis and interpretation of data: Feagan, Sandborn, Mittmann, Bar-Meir, Paré, Wong, Zou, Donner, Rutgeerts.

Drafting of the manuscript: Feagan, Hébuterne, Klvana, Ardizzone, Kiudelis.

Critical revision of the manuscript for important intellectual content: Feagan, Sandborn, Mittmann, Bar-Meir, D'Haens, Bradette, Cohen, Dallaire, Ponich, McDonald, Paré, Niv, Alexeeva, Rostom, Spleiss, Gilgen, Vandervoort, Wong, Zou, Donner, Rutgeerts.

Statistical analysis: Wong, Zou, Donner.

Obtained funding: Sandborn, Klvana, Alexeeva, Spleiss, Rutgeerts.

Administrative, technical, or material support: Feagan, Sandborn, Bar-Meir, D'Haens, Spleiss, Gilgen, Vandervoort.

Study supervision: Feagan, Mittmann, Cohen, Dallaire, Ponich, McDonald, Hébuterne, Paré, Klvana, Ardizzone, Alexeeva, Kiudelis, Vandervoort, Rutgeerts.

Financial Disclosures: Dr Feagan reported receiving grant and research support from Schering-Plough, Otsuka, Millennium, Tillotts, Abbott, Protein Design Labs, Boehringer-Ingelheim, Novartis, Centocor, Berlex, Synta, Schering Canada, Elan/Biogen, UCB Pharma, BMS, Procter & Gamble, and Napo Pharma; receiving consulting fees from Synta, Millennium, Schering Canada, Celltech, Centocor, Elan/Biogen, Janssen-Ortho, Protein Design Labs, ISIS, Teva Pharmaceuticals, Santarus, Schering-Plough, Bristol-Myers Squibb, Celgene, CombinatorRx Inc, UCB Pharma, Napo Pharma, Abbott, Procter & Gamble, Osiris, Berlex, AstraZeneca, GeneLogic Inc, Cerimon Pharm, Tioga Pharm, Serono, Genentech, Tillotts Pharma AG, and CombinatoRx Inc; serving on the speakers bureau of AstraZeneca; and serving on the scientific advisory board of Protein Design Labs, AstraZeneca, Elan/Biogen, Celltech, Synta, Schering Canada, Celgene, and Tillotts Pharma AG. Dr Sandborn reported receiving grants or funding from Abbott Laboratories, Bristol-Myers Squibb, Centocor, Chemocentryx, Elan Pharmaceuticals Inc, Otsuka American Pharmaceuticals Inc, PDL Biopharma, Procter & Gamble, Prometheus Laboratories, Salix Pharmaceuticals Inc, Schering-Plough, Serono, Shire Pharmaceuticals, Targacept Inc, Techlab Inc, Tillotts Pharma AG, and UCB Pharma (previously Celltech Therapeutics Ltd); receiving consulting fees from GlaxoSmithKline (previously Glaxo Wellcome), Abbott Laboratories (fees paid to the Mayo Clinic), ActoGenix NV, AGI Therapeutics Inc, Ajinomoto Pharmaceuticals, Alba Therapeutics, Alzyme plc, Alza, Amgen Inc, Ardea Biosciences, Aspreva Pharmaceuticals, AstraZeneca, Atrix Laboratories Inc (acquired by QLT in 2004), Avidia Inc, Berlex, Bexel Pharmaceuticals Inc, BioBalance, Boehringer-Ingelheim Inc, Bristol Meyers Squibb, Celgene, Celltech (acquired by UCB Pharma in 2004) (fees paid to the Mayo Clinic), Centocor (fees paid to the Mayo Clinic), Cerimon Pharmaceutical Inc, Chemocentryx, CombinatoRx Inc, Com-Mentis, Corautes Genetics, Cosmo Technologies Ltd, CuraGen, Effective Pharmaceuticals Inc, Eisai Medical Research Inc, Elan Pharmaceuticals Inc, Entero-medics, Enzo Therapeutics Inc, Eurand, FlexPharm Ltd (previously named Enterotech), Genencor International, Genentech, H3 Pharma (name changed to Debiopharm SA in 2005), Hoffman LaRoche Inc (Roche), Hutchison Medipharma Ltd, Inflabloc Pharmaceuticals (previously named Pharmadigm Inc), Intek Phar-

maceutical, ISIS Pharmaceuticals, Jacobus Pharmaceutical Co, Johnson & Johnson Pharmaceutical Research & Development, LigoCyte Pharmaceuticals, McNeil Consumer and Specialty Pharmaceuticals, Medarex Inc, Merck Research Laboratories, Millennium Pharmaceuticals Inc, Nissin Kyojin Pharmaceutical Co Ltd, Novartis, NPS Pharmaceuticals, Ocera Therapeutics Inc (previously named Renovia Inc), Ono Pharma USA, Otsuka American Pharmaceuticals Inc, PDL Biopharma (previously named Protein Design Labs) (fees paid to the Mayo Clinic), Pfizer, Procter & Gamble (fees paid to the Mayo Clinic), Prometheus Laboratories, Renovis Inc, Salix Pharmaceuticals Inc, Sangstat (acquired by Genzyme in 2003), Schering-Plough, Serono Inc, Shire Pharmaceuticals, Synta Pharmaceuticals, Targacept, Teva Pharmaceuticals, Therakos, Tioga Pharmaceuticals, UCB Pharma (fees paid to the Mayo Clinic), Vela Pharmaceuticals Inc, and ViaCell Inc; and receiving honoraria for speaking at continuing medical education events indirectly sponsored by Abbott Laboratories, Acan Pharmaceuticals, AstraZeneca, Centocor, Elan Pharmaceuticals Inc, Falk Pharma, Otsuka American Pharmaceuticals Inc, PDL Biopharma, Procter & Gamble, Prometheus Laboratories, Salix Pharmaceuticals Inc, Schering-Plough, Shire Pharmaceuticals, and UCB Pharma. Dr Mittmann reported receiving product development consultancy fees from Tillotts Pharma AG. Dr Bar-Meir reported receiving consulting fees from Tillotts Pharma AG. Dr Rostom reported receiving consulting fees from Novartis and Bayer and an unrestricted educational grant from ALBA Therapeutics. Mr Spleiss reported having employee equity ownership in Tillotts Pharma AG. Dr Rutgeerts reported receiving consulting fees from Centocor, Schering-Plough, UCB Abbott, Elan-Biogen, PDL, Avidia, Bristol-Myers Squibb, Millennium, and Tillotts; lecture fees from Centocor, Schering-Plough, UCB, Abbott, and Elan-Biogen; and grant support from Centocor, Schering-Plough, Abbott, and UCB. Drs D'Haens, Bradette, Cohen, Dallaire, Ponich, McDonald, Kiudelis, Hébuterne, Paré, Klvana, Niv, Ardizzone, Alexeeva, and Rostom were site investigators and were compensated for performing clinical measurements. No other financial disclosures were reported.

Funding/Support: This study was funded by Tillotts Pharma.

Role of the Sponsor: The trial was designed by the academic investigators in collaboration with representatives of Tillotts, and representatives of Tillotts reviewed and approved the final version of the manuscript.

Data and Safety Monitoring Board: University of Health Sciences Center, Denver, Colorado: J. Singleton; Charité-Universitätsmedizin Berlin: H. Lochs; Wolfson Institute of Preventive Medicine, London, UK: J. Cuzick. **Data Coordinating Center:** Roberts Clinical Trials, Roberts Research Institute, The University of Western Ontario, London, Ontario: F. Babar, B. Bergman, M. Bertin, T. Clayton, M. Hammond, M. Hopkins, L. Jensen, E. Karugu, L. Liddiard, F. Livingstone, M. Mathers, C. Mueller, S. Nelson, J. Rémillard, I. Ruocco, B. Sarazin, B. Sharpe, L. Smith, H. Sun, K. Toulouse. **Participating Investigators:** Atlanta Gastroenterology Associates, Atlanta, Georgia: D. Wolf, M. Monroe; Carle Clinic Association, Urbana, Illinois: E. Greenberg, V. Carlton; Cleveland Clinic Foundation, Cleveland, Ohio: D. Seidner, S. Zurc; Clinical Research Institute of Michigan, Chesterfield: R. Fogel, A. Leyson-Fiel; Health Sciences Centre, Calgary, Alberta: R. Panaccione, A. Daye; Health Sciences Centre, Edmonton, Alberta: R. Fedorak, C. McCabe-Woodrow; Health Sciences Centre, Winnipeg, Manitoba: C. Bernstein, T. Rawsthorne; Hôpital Maisonneuve-Rosemont, Montreal, Québec: G. Aumais, L. Henri; Hôpital St-Sacrement, Québec City, Québec: P. Paré, S. Rousseau; Hôtel Dieu de Lévis, Lévis, Québec: R. Bourdages, M. Berber; Hotel Dieu Hospital, Kingston, Ontario: W. Depew, D. Brady; Jew-

ish General Hospital, Montreal, Québec: A. Cohen, N. Desjardins; LAIR Centre, Vancouver, British Columbia: F. Anderson, I. Wong; London Health Sciences Centre South Street Campus, London Ontario: T. Ponich, P. Walton-Mennill; London Health Sciences Centre University Campus, London, Ontario: J. McDonald, W. McCaw; Long Island Clinical Research Associates, Great Neck, New York: S. Katz, M. Mustac; Mayo Clinical Rochester, Minnesota: W. Sandborn, R. Jeché; Montreal General Hospital, Montreal, Québec: G. Wild, N. Pellerin; Mount Sinai Hospital, Toronto, Ontario: H. Steinhart, S. Mikolajinis; Pavillon-Hôtel-Dieu de Québec CHUQ, Québec City, Québec: M. Bradette, S. Lepire; Pavillon St-François d'Assise CHUQ, Québec City, Québec: C. Dallaire, F. Bernard; Royal Alexandra Hospital, Edmonton, Alberta: R. Bailey, D. Meyer; Royal Victoria Hospital, Montreal, Québec: A. Bitton, M. Bernier; Saint John Regional Hospital, Saint John, New Brunswick: A. Cockeram, H. Long; St Joseph's Health Centre, London, Ontario: R. Sy, L. Moyer; St Michael's Hospital, Toronto, Ontario: J. Baker, G. Stewart; St Paul's Hospital GI Research, Vancouver, British Columbia: R. Enns, S. Patterson; Sunnybrook & Women's College Health Science Centre, Toronto, Ontario: F. Saibil, M. Morgan; The Ottawa Hospital-Civic Campus, Ottawa, Ontario: A. Rostom, P. Waddell; University of Chicago Medical Center, Chicago, Illinois: S. Hanauer, H. Brenner; University of Kentucky Medical Branch, Lexington: T. Winter, J. Reynolds; University of North Carolina at Chapel Hill: K. Isaacs; D. Walkup; University of Texas Medical Branch, Galveston: J. Sellin, J. Hirsch; Victoria General Hospital QEII Health Sciences Centre, Halifax, Nova Scotia: D. McIntosh, C. Kelly; Albrecht-Ludwigs-University, Freiburg, Germany: J. Rasenack, I. Borsukiewicz-Krupa; Internistische Praxis, Medizin, Berlin, Germany: H.-J. Kramm, I. Borsukiewicz-Krupa; Barzilai Medical Center, Ashkelon, Israel: M. Faschik, I. Rosen-Leibowitz; CHU de Grenoble, Grenoble, France: B. Bonaz, S. Zachayus; District Hospital, Usti nad Orlici, Czech Republic: L. Gabalec, L. Pribylova; District Hospital, Ceske Budejovice, Czech Republic: O. Shonova, L. Pribylova; Faculty Hospital, Prague, Czech Republic: J. Benes, L. Pribylova; Hadassah University Hospital, Jerusalem, Israel: E. Goldin, I. Rosen-Leibowitz; Heilig Hartziekenhuis, Roeselare, Belgium: F. Baert, W. Walravens; Hillel Yaffe Medical Center, Hadera, Israel: Z. Fireman, I. Rosen-Leibowitz; Hôpital Civil, Strasbourg, France: D. Vetter, S. Zachayus; Hôpital Saint Antoine, Paris, France: J. P. Gendre, S. Zachayus; Hôpital Nord, Marseille, France: J.-C. Grimaud, S. Zachayus; Imelda Ziekenhuis, Bonheiden, Belgium: G. D'Haens, K. Lauwers; Internistische Gemeinschaftspraxis, Rödermark-Oberoden, Germany: S. Orlemann, I. Borsukiewicz-Krupa; Kaunas Medical University Clinics, Kaunas, Lithuania: G. Kiudelis, L. Didziapetriene; Klinikum der Universität Regensburg, Regensburg, Germany: H. Herfarth, I. Borsukiewicz-Krupa; Klinikum Mannheim, Mannheim, Germany: U. Böcker, I. Borsukiewicz-Krupa; Masaryk Hospital, Usti nad Labem, Czech Republic: J. Stehlik, L. Pribylova; Medizinische Hochschule Hannover, Hannover, Germany: M. Manns, I. Borsukiewicz-Krupa; Medizinische Universitäts, Erlangen, Germany: D. Raitel, I. Borsukiewicz-Krupa; Ospedale G. B. Morgagni, Forlì, Italy: D. Valpiani, P. Zamboni; Ospedale Luigi Sacco, Milan, Italy: S. Ardizzone, P. Zamboni; Ospedale Nuovo, Imola, Italy: C. Brignola, P. Zamboni; Ospedale Santissima Annunziata, Canto, Italy: G. Zoli, P. Zamboni; Policlinico S. Orsola Malpighi, Bologna, Italy: A. Belluzzi, T. Perrone; Rabin Medical Center Beilinson Campus, Petah Tiqwa, Israel: Y. Niv, O. Ben-Shitrit; Ranbam Medical Center, Haifa, Israel: A. Eliakim, I. Rosen-Leibowitz; Soroka University Medical Center, Beer Sheva, Israel: A. Fich, I. Rosen-Leibowitz; Tel Aviv Sourasky Medical Center, Tel Aviv, Israel: I. Dotan, I. Rosen-Leibowitz; The

Chaim Sheba Medical Center, Tel Hashomer, Israel: A. Lang, I. Rosen-Leibowitz; The Edith Wolfson Medical Center, Holon, Israel: Y. Avni, I. Rosen-Leibowitz; ULB Hôpital Erasme, Brussels, Belgium: A. Van Gossum, W. Walravens; University Hospital, Ostrava, Czech Republic: P. Klvana, L. Pribylova; University Hospital, Pilsen, Czech Republic: J. Kozeluhova, L. Pribylova; University of Nice Sophia Antipolis, Nice, France: X. Hébuterne; UZ Gashuisberg, Leuven, Belgium: P. Rutgeerts, K. Lauwers; Vilnius University Hospital, Vilnius, Lithuania: A. Irnius, L. Didziapetrienė; Western Galilee Hospital, Nahariya, Israel: R. Reshef, I. Rosen-Leibowitz; Clinical Hospital, #33, Nizhny Novgorod, Russia: O. Alexeeva, S. Krishopenko, E. Galperin, O. Chertkova; Military Medical Academy, St. Petersburg, Russia: V. Grinevich, I. Gubonina, A. Pershko; St. Petersburg Medical Academy, St. Petersburg, Russia: A. Baranovsky, O. Shchukina, E. Kondrashina; Medical Academy, St. Petersburg, Russia: V. Simanenkova, E. Poroshina, O. Solov'yeva; Moscow Regional Research Clinical Institute, Moscow, Russia: E. Bebusova, O. Kuzmina, I. Domareva; Regional Clinical Hospital, Yaroslavl, Russia: O. Khrustalev, O. Okhapkina, S. Tsybulko.

REFERENCES

- Akobeng AK, Gardener E. Oral 5-aminosalicylic acid for maintenance of medically-induced remission in Crohn's Disease. *Cochrane Database Syst Rev*. 2005; (1):CD003715.
- Candy S, Wright J, Gerber M, Adams G, Gerig M, Goodman R. A controlled double blind study of azathioprine in the management of Crohn's disease. *Gut*. 1995;37(5):674-678.
- Feagan BG, Fedorak RN, Irvine EJ, et al; North American Crohn's Study Group Investigators. A comparison of methotrexate with placebo for the maintenance of remission in Crohn's disease. *N Engl J Med*. 2000;342(22):1627-1632.
- Present DH, Korelitz BI, Wisch N, Glass JL, Sachar DB, Pasternack BS. Treatment of Crohn's disease with 6-mercaptopurine: a long-term, randomized, double-blind study. *N Engl J Med*. 1980;302(18):981-987.
- Hanauer SB, Feagan BG, Lichtenstein GR, et al. Maintenance infliximab for Crohn's disease: the ACCENT 1 randomised trial. *Lancet*. 2002;359(9317):1541-1549.
- Colombel JF, Sandborn WJ, Rutgeerts P, et al. Adalimumab for maintenance of clinical response and remission in patients with Crohn's disease: the CHARM trial. *Gastroenterology*. 2007;132(1):52-65.
- Schreiber S, Khaliq-Kareemi M, Lawrance I, et al. Maintenance therapy with certolizumab pegol for Crohn's disease. *N Engl J Med*. 2007;357(3):239-250.
- Sandborn WJ, Feagan BG, Stoinov S, et al. Certolizumab pegol for the treatment of Crohn's disease. *N Engl J Med*. 2007;357(3):228-238.
- Nurmohamed MT, Dijkmans BA. Efficacy, tolerability and cost effectiveness of disease-modifying antirheumatic drugs and biologic agents in rheumatoid arthritis. *Drugs*. 2005;65(5):661-694.
- Kirschner BS. Safety of azathioprine and 6-mercaptopurine in pediatric patients with inflammatory bowel disease. *Gastroenterology*. 1998;115(4):813-821.
- Keane J, Gershon S, Wise RP, et al. Tuberculosis associated with infliximab, a tumor necrosis factor alpha-neutralizing agent. *N Engl J Med*. 2001;345(15):1098-1104.
- Goldberg RJ, Katz J. A meta-analysis of the analgesic effects of omega-3 polyunsaturated fatty acid supplementation for inflammatory joint pain. *Pain*. 2007;129(1-2):210-223.
- Stamp LK, James MJ, Cleland LG. Diet and rheumatoid arthritis: a review of the literature. *Semin Arthritis Rheum*. 2005;35(2):77-94.
- Donadio JV Jr, Larson TS, Bergstralh EJ, Grande JP. A randomized trial of high-dose compared with low-dose omega-3 fatty acids in severe IgA nephropathy. *J Am Soc Nephrol*. 2001;12(4):791-799.
- Durrington PN, Bhatnagar D, Mackness MI, et al. An omega-3 polyunsaturated fatty acid concentrate administered for one year decreased triglycerides in simvastatin treated patients with coronary heart disease and persisting hypertriglyceridaemia. *Heart*. 2001;85(5):544-548.
- Bays H. Clinical overview of Omacor: a concentrated formulation of omega-3 polyunsaturated fatty acids. *Am J Cardiol*. 2006;98(4A):711-761.
- Stalenhoef AF, de Graaf J, Wittekoek ME, Bredie SJ, Demacker PN, Kastelein JJ. The effect of concentrated n-3 fatty acids versus gemfibrozil on plasma lipoproteins, low density lipoprotein heterogeneity and oxidizability in patients with hypertriglyceridemia. *Atherosclerosis*. 2000;153(1):129-138.
- Yokoyama M, Origasa H, Matsuzaki M, et al. Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomised open-label, blinded endpoint analysis. *Lancet*. 2007;369(9567):1090-1098.
- Marchioli R, Barzi F, Bomba E, et al. Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction: time-course analysis of the results of the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI)-Prevenzione. *Circulation*. 2002;105(16):1897-1903.
- Bucher HC, Hengstler P, Schindler C, Meier G. N-3 polyunsaturated fatty acids in coronary heart disease: a meta-analysis of randomized controlled trials. *Am J Med*. 2002;112(4):298-304.
- Belluzzi A, Brignola C, Campieri M, Pera A, Boschi S, Miglioli M. Effect of an enteric-coated fish-oil preparation on relapses in Crohn's disease. *N Engl J Med*. 1996;334(24):1557-1560.
- Lorenz-Meyer H, Bauer P, Nicolay C, et al; Study Group Members (German Crohn's Disease Study Group). Omega-3 fatty acids and low carbohydrate diet for maintenance of remission in Crohn's disease: a randomized controlled multicenter trial. *Scand J Gastroenterol*. 1996;31(8):778-785.
- Best WR, Beckett JM, Singleton J, Kern F Jr. Development of a Crohn's disease activity index: National Cooperative Crohn's Disease Study. *Gastroenterology*. 1976;70(3):439-444.
- Summers RW, Switz DM, Sessions JT Jr, et al. National Cooperative Crohn's Disease Study: results of drug treatment. *Gastroenterology*. 1979;77(4, pt 2):847-869.
- Greenberg GR, Feagan BG, Martin F, et al. Oral budesonide for active Crohn's disease. *N Engl J Med*. 1994;331(13):836-841.
- Ware JE Jr, Sherbourne CD. The MOS 36-Item Short-Form Health Survey (SF-36), I: conceptual framework and item selection. *Med Care*. 1992;30(6):473-483.
- International Federation of Pharmaceutical Manufacturers Associations. *Medical Dictionary for Regulatory Activities (MedDRA) Version 9.0*. Chantilly, VA: MedDRA MSSO; 2007.
- SAS Statistical Software Version 8.2. Cary, NC: SAS Institute Inc; 2001.
- StatXact Version 8.0 With Cytel Studio Version 8. Cambridge, MA: Cytel Inc; 2007.
- Sandborn WJ, Colombel JF, Enns R, et al. Natalizumab induction and maintenance therapy for Crohn's disease. *N Engl J Med*. 2005;353(18):1912-1925.
- Greenberg GR, Feagan BG, Martin F, et al; Canadian Inflammatory Bowel Disease Study Group. Oral budesonide as maintenance treatment for Crohn's disease: a placebo-controlled, dose-ranging study. *Gastroenterology*. 1996;110(1):45-51.
- Harmonised Tripartite Guideline ICH: Guideline for Good Clinical Practice E6(R1). International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use Web site. 1996. <http://www.ich.org/LOB/media/MEDIA482.pdf>. Accessibility verified March 12, 2008.
- Harvey RF, Bradshaw JM. A simple index of Crohn's-disease activity. *Lancet*. 1980;1(8167):514.
- Kjaergard LL, Villumsen J, Glud C. Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses. *Ann Intern Med*. 2001;135(11):982-989.
- Glud LL. Bias in clinical intervention research. *Am J Epidemiol*. 2006;163(6):493-501.
- Kjaergard LL, Nikolova D, Glud C. Randomized clinical trials in hepatology: predictors of quality. *Hepatology*. 1999;30(5):1134-1138.
- Hilsden RJ, Scott CM, Verhoef MJ. Complementary medicine use by patients with inflammatory bowel disease. *Am J Gastroenterol*. 1998;93(5):697-701.
- Heuschkel R, Afzal N, Wuerth A, et al. Complementary medicine use in children and young adults with inflammatory bowel disease. *Am J Gastroenterol*. 2002;97(2):382-388.
- Leong RW, Lawrance IC, Ching JY, et al. Knowledge, quality of life, and use of complementary and alternative medicine and therapies in inflammatory bowel disease: a comparison of Chinese and Caucasian patients. *Dig Dis Sci*. 2004;49(10):1672-1676.
- Li FX, Verhoef MJ, Best A, Otley A, Hilsden RJ. Why patients with inflammatory bowel disease use or do not use complementary and alternative medicine: a Canadian national survey. *Can J Gastroenterol*. 2005;19(9):567-573.
- Langmead L, Rampton DS. Review article: complementary and alternative therapies for inflammatory bowel disease. *Aliment Pharmacol Ther*. 2006;23(3):341-349.
- McClaskey EM, Michalets EL. Subdural hematoma after a fall in an elderly patient taking high-dose omega-3 fatty acids with warfarin and aspirin: case report and review of the literature. *Pharmacotherapy*. 2007;27(1):152-160.
- Knapp HR, Reilly IA, Alexandrini P, FitzGerald GA. In vivo indexes of platelet and vascular function during fish-oil administration in patients with atherosclerosis. *N Engl J Med*. 1986;314(15):937-942.