



US008680364B2

(12) **United States Patent**  
Hill et al.

(10) **Patent No.:** **US 8,680,364 B2**  
(45) **Date of Patent:** **\*Mar. 25, 2014**

(54) **SOYBEAN GENES FOR RESISTANCE TO  
APHIS GLYCINES**

(75) Inventors: **Curtis B. Hill**, Champaign, IL (US);  
**Glen L. Hartman**, Urbana, IL (US);  
**Yan Li**, Chicago, IL (US); **Brian Diers**,  
Urbana, IL (US); **Shawn Carlson**,  
Bondville, IL (US)

(73) Assignees: **The Board of Trustees of the  
University of Illinois**, Urbana, IL (US);  
**The United States of America, as  
represented by the Secretary of  
Agriculture**, Washington, DC (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 63 days.

This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **13/149,762**

(22) Filed: **May 31, 2011**

(65) **Prior Publication Data**

US 2011/0277177 A1 Nov. 10, 2011

#### Related U.S. Application Data

(62) Division of application No. 11/158,307, filed on Jun.  
21, 2005, now Pat. No. 7,994,389.

(60) Provisional application No. 60/581,501, filed on Jun.  
21, 2004.

(51) **Int. Cl.**  
**A01H 1/00** (2006.01)  
**A01H 1/04** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **800/266**; 800/267; 800/312

(58) **Field of Classification Search**  
None  
See application file for complete search history.

(56) **References Cited**

#### U.S. PATENT DOCUMENTS

5,491,081 A	2/1996	Webb	
5,689,035 A	11/1997	Webb	
5,948,953 A	9/1999	Webb	
6,054,300 A	4/2000	McKendree	
6,162,967 A	12/2000	Webb	
6,538,175 B1	3/2003	Webb	
7,994,389 B2 *	8/2011	Hill et al.	800/267
2004/0034890 A1	2/2004	St. Martin et al.	
2004/0261144 A1	12/2004	Martin et al.	

#### OTHER PUBLICATIONS

Sun et al (Soybean Genet News 17: 43-48, 1990).\*

Narvel et al (Crop Sci 41: 1931-1939, 2001).\*

Hill et al (Crop Science 44: 98-106, 2004).\*

Clark et al. Plant Dis. 86:1219-1222 2002.

Harrewijn, P. et al., "Integrated aphid management: general aspects," pp. 267-272, In Minks and Harrewijn (ed) Aphids: Their biology, natural enemies and control, vol. C. Elsevier. New York 1989.

Jeong et al. Abstract published by Plant & Animal Genome VIII Conference, Jan. 9-12, 2000 www.intl-pag.org/8/abstracts/pag8255.html.

Jeong et al. Plant Breeding 123:305-310 2004.

Rongwen et al. Theor Appl Genet 90:43-48 1995.

Tamulonis et al., Theor. Appl. Genet. 95:664-670 1997.

Yong et al., Proc. Nat. Acad. Sci. USA 933:11751-11755 1996.

Kim, K.S. et al., "Fine mapping the soybean aphid resistance gene Rag1 in soybean," Theor Appl Genet (2010) 120:1063-1071.

Zhu et al. (2006), "Fine Mapping of a Major Insect Resistance QTL in Soybean and its Interaction with Minor Resistance QTLs," *Crop Sci.* 46:1094-1099.

Bissonnette, S. (2002) "Good News on Aphids: First Resistance Found," *Champaign-Urbana News Gazette* Aug. 4.

Auclair, J.L. (1989) "Host Plant Resistance," In: *Aphids: Their Biology, Natural Enemies, and Control*, vol. C., Harrewijn et al. ed., Elsevier, New York, pp. 225-265.

Bell-Johnson et al. (1998) "Biotechnology Approaches to Improving Resistance to SCN and SDS: Method for High Throughput Marker Assisted Selection," *Soybean Genet. Newlett.* 25:115-117.

Bissonnette, S. (2002) "Good News on Aphids: First Resistance Found," *Champaign-Urbana News Gzette* Aug. 4.

Clark et al. (2002) "Transmissibility of Field Isolates of Soybean Viruses by *Aphis glycines*," *Plant Dis.* 86:1219-1222.

Cregan et al. (1999) "An Integrated Genetic Linkage Map of the Soybean Genome," *Crop Sci.* 39:1464-1490.

Du Toit, F. (1987) "Resistance in Wheat (*Triticum aestivum*) to *Diuraphis noxia* (Homoptera: Aphididae)," *Cereal Res. Commun.* 15:175-179.

Harrewijn et al. (1989) "Integrated Aphid Management: General Aspects," In: *Aphids: Their Biology, Natural Enemies, and Control*, vol. C., Harrewijn et al. ed., Elsevier, New York, pp. 267-272.

Hartman et al. (2001) "Occurrence and Distribution of *Aphis glycines* on Soybeans in Illinois in 2000 and its Potential Control," <http://www.plantmanagementnetwork.org/php/default.asp>.

Hill et al. (2002) "Resistance to the Soybean Aphid in Soybean and Other Legumes," *APS Annual Meeting*, Jul. 27-31.

Hill et al. (2004) "Resistance to the Soybean Aphid in Soybean Germplasm," *Crop Sci.* 44:981-106.

Hill et al. (2004) "Resistance of Glycine Species and Various Cultivated Legumes to the Soybean Aphid (Homoptera: Aphididae)," *J. Econ. Entomol.* 97(3):1071-1077.

Hill et al. (1996) "First Report of Transmission of *Soybean Mosaic Virus* and *Alfalfa Mosaic Virus* by *Aphis glycines* in the New World" *Plant Dis.* 85:561.

Hill et al. (2002) "Resistance to the Soybean Aphid in Soybean and Other Legumes," *Phytopathol.* 92:S36.

(Continued)

Primary Examiner — Shubo (Joe) Zhou

Assistant Examiner — Keith Robinson

(74) Attorney, Agent, or Firm — Ellen P. Winner; e winner & associates, PLLC

(57) **ABSTRACT**

*Aphis glycines* resistance (RAG) genes are provided by this invention, along with methods for identifying their presence using marker-assisted selection. Varieties of *G. max* and *G. soja* having resistance to *A. glycines* have been identified. The RAG genes, as well as the methods, aphid-resistant varieties, and markers disclosed herein may be used to breed new elite lines expressing soybean aphid resistance.

**10 Claims, 6 Drawing Sheets**

(56)

## References Cited

## OTHER PUBLICATIONS

- Hill et al. (2002) "Resistance to the Soybean Aphid in Soybean and Other Legumes," Poster, National Meeting of the American Phytopathological Society, Milwaukee, Wisconsin.
- Hill et al. (Nov. 2002) Oral Research Report, North Central Soybean Virus Workers Conference, Iowa State University, Ames, IA.
- Hill et al. (2003) "Resistance to the Soybean Aphid (*Aphis glycines*) has Been Discovered in Ancestral Soybean Germplasm," <http://www.soydiseases.uiuc.edu/projects.php?id=1>.
- Hill et al. (2004) "Resistance to the Soybean Aphid in Soybean Germplasm and other Legumes," Abstract and poster, World Soybean Research Conference, Iguatua Falls, Brazil.
- Hirano et al. (1996) "Effects of Temperature on Development, Longevity and Reproduction of the Soybean Aphid, *Aphis glycines* (Homoptera: Aphididae)," *Appl. Entomol. Zool.* 31(1):178-180.
- Hymowitz, T. (1970) "On the Domestication of the Soybean", *Econ. Bot.* 24:408-421.
- Iwaki et al. (1980) "A Persistent Aphid Borne Virus of Soybean, Indonesian Soybean Dwarf Virus Transmitted by *Aphis glycines*," *Plant Dis.* 64:1027-1030.
- Kaloshian et al (1997) "The impact of the *Meu1*-Mediated Resistance in Tomato on Longevity, Fecundity and Behavior of the Potato Aphid, *Macrosiphum euphorbiae*," *Entomol. Exp. Appl.* 83:181-187.
- Keim et al. (1988) "Construction of a Random Recombinant DNA Library that is Primary Single Copy Sequences," *Soybean Genet. Newslett.* 151:147-148.
- Kline, G. (Jun. 8, 2003) "Out of UI Vaults: Bug Resistance," *Champaign-Urbana News-Gazette*.
- Klinger et al. (2001) "Mapping of Cotton-Melon Aphid Resistance in Melon," *J. Am. Soc. Hortic. Sci.* 136:56-63.
- Li et al. (2004) "Effect of Three Resistant Soybean Genotypes on the Fecundity, Mortality, and Maturation of Soybean Aphid (Homoptera: Aphididae)," *J. Econ. Entomol.* 37(3):1106-1111.
- Luginbill, J.P. (1969) "Developing Resistant Plants—The Ideal Method of Controlling Insects," USDA, ARS. Prod Res. Rep. 111, USGPO, Washington, D.C.
- Mansur et al. (1996) "Genetic Mapping of Agronomic Traits Using Recombinant Inbred Lines of Soybean," *Crop Sci.* 36:1327-1336.
- Michelmore et al. (1991) "Identification of Markers Linked to Disease-Resistance Genes by Bulk Segregant Analysis: A Rapid Method to Detect Markers in Specific Genomic Regions by Using Segregating Populations," *Proc. Natl. Acad. Sci. USA* 88:9828-9832.
- Narvel et al. (2001) "A Retrospective DNA Marker Assessment of the Development of Insect Resistant Soybean," *Crop Sci.* 41:1931-1939. NCBI website, <http://www.ncbi.nlm.nih.gov/>, 2004.
- Ostlie, K. (2002) "Managing Soybean Aphid," [http://www.soybeans.umn.edu/crop/insects/aphid/aphid\\_publication\\_managingsba.htm](http://www.soybeans.umn.edu/crop/insects/aphid/aphid_publication_managingsba.htm).
- Patterson et al. (2002) "Assessing and Managing Risk from Soybean Aphids in the North Central States," <http://www.planthealth.info/soyaphid/aphid02.htm>.
- Sama et al. (1974) Research Reports 1968-1974, In, Varietal Screening for Resistance to the Aphid, *Aphis glycines*, in Soybean, Argicultural Cooperation, Indonesia, Netherlands, pp. 171-172.
- Sun et al. (1990) "Study on the Uses of Aphid-Resistant Character in Wild Soybean. I. Aphid-Resistance Performance of F<sub>2</sub> Generation from Crosses Between Cultivated and Wild Soybeans," *Soybean Genet. News* 17:43-48.
- Tyler et al. (1985) "Biotype E Greenbug Resistance in Wheat Streak Mosaic Virus-Resistant Wheat Germplasm Lines," *Crop Sci.* 25:686-688.
- Van Ooijen et al. (2001) "Software for the Calculation of Genetic Linkage Maps," JoinMap 3.0, Plant Research International, Wageningen, the Netherlands.
- Wang et al. (1996) "Effects of Soybean Aphid, *Aphis glycines* on Soybean Growth and Yield," *Soybean Sci.* 15:243-247.
- Wang et al. (2003) "A Low-Cost, High-Throughput Polyacrylamide Gel Electrophoresis System for Genotyping With Microsatellite DNA Markers," *Crop Sci.* 43:1828-1832.
- Wu et al. (2004) "A BAC and BIBAC-based Physical Map of the Soybean Genome," *Genome Res.* 14(2):319-326.
- Zhang et al. (2004) "QTL Mapping of Ten Agronomic Traits on the Soybean (*Glycine max* L. Merr.) Genetic Map and Their Association with EST Markers," *Theor. Appl. Genet.* 108:1131-1139.
- Zhuang et al. (1996) "A Study on Resistance to Soybean Mosaic Virus and *Aphis glycine* of Perennial Wild Soybean," *Soybean Genet. Newslett.* 23:66-69.

\* cited by examiner




FIG. 1A





FIG 1B






FIG 1C




FIG 1D




FIG. 2



FIG. 3



# SOYBEAN GENES FOR RESISTANCE TO APHIS GLYCINES

## CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a divisional application of U.S. Ser. No. 11/158,307 filed Jun. 21, 2005, which claims priority to U.S. Provisional Application No. 60/581,501 filed Jun. 21, 2004, both of which are incorporated by reference herein to the extent not inconsistent herewith.

## BACKGROUND OF THE INVENTION

This invention relates to a soybean gene for resistance to *Aphis glycines*, to soybean plants possessing this gene, which maps to a novel chromosomal locus, and to methods for identifying and breeding these plants, the methods involving marker-assisted selection.

Soybeans (*Glycine max* L. Merr.) are a major cash crop and investment commodity in North America and elsewhere. Soybean oil is one of the most widely used edible oils, and soybeans are used worldwide both in animal feed and in human food production.

A native of Asia, the soybean aphid was first found in the Midwest in 2000 (Hartman, G. L. et al., "Occurrence and distribution of *Aphis glycines* on soybeans in Illinois in 2000 and its potential control," (1 Feb. 2001 available at a website address beginning with the usual http and www prefixes, followed by plantmanagementnetwork.org/php/default, followed by the suffix .asp.) It rapidly spread throughout the region and into other parts of North America (Patterson, J. and Ragsdale, D., "Assessing and managing risk from soybean aphids in the North Central States," (11 Apr. 2002) available at a website address beginning with the usual http and www prefixes, followed by planthealth.info/soyaphid/aphid02, followed by the suffix .htm.) High aphid populations can reduce crop production directly when their feeding causes severe damage such as stunting, leaf distortion, and reduced pod set (Sun, Z. et al., "Study on the uses of aphid-resistant character in wild soybean. I. Aphid-resistance performance of F<sub>2</sub> generation from crosses between cultivated and wild soybeans," (1990) *Soybean Genet. News.* 17:43-48). Yield losses attributed to the aphid in some fields in Minnesota during 2001, where several thousand aphids occurred on individual soybean plants, were >50% (Ostlie, K., "Managing soybean aphid," (2 Oct. 2002) available at a website having an address beginning with the usual http and www, followed by soybeans.umn.edu/crop/insects/aphid/aphid\_publication\_managingsba, and having a suffix .htm) with an average loss of 101 to 202 kg ha<sup>-1</sup> in those fields (Patterson and Ragsdale, supra). In earlier reports from China, soybean yields were reduced up to 52% when there was an average of about 220 aphids per plant (Wang, X. B. et al., "A study on the damage and economic threshold of the soybean aphid at the seedling stage," (1994) *Plant Prot.* (China) 20:12-13) and plant height was decreased by about 210 mm after severe aphid infestation (Wang, X. B. et al., "Study on the effects of the population dynamics of soybean aphid (*Aphis glycines*) on both growth and yield of soybean," (1996) *Soybean Sci.* 15:243-247). An additional threat posed by the aphid is its ability to transmit certain plant viruses to soybean such as Alfalfa mosaic virus, Soybean dwarf virus, and Soybean mosaic virus (Sama, S. et al., "Varietal screening for resistance to the aphid, *Aphis glycines*, in soybean," (1974) *Research Reports* 1968-1974, pp. 171-172; Iwaki, M. et al., "A persistent aphid borne virus of soybean, Indonesian Soybean dwarf virus transmitted by

*Aphis glycines*," (1980) *Plant Dis.* 64:1027-1030; Hartman, G. L. et al., supra; Hill, J. H. et al., "First report of transmission of Soybean mosaic virus and Alfalfa mosaic virus by *Aphis glycines* (Homoptera: Aphididae)," (1996) *Appl. Entomol. Zool.* 31:178-180; Clark, A. J. and Perry, K. L., "Transmissibility of field isolates of soybean viruses by *Aphis glycines*," (2002) *Plant Dis.* 86:1219-1222).

Because *A. glycines* is a recent pest in the USA, a comprehensive integrated management approach to control the aphid has yet to be developed. Research to evaluate the efficacy of currently-available insecticides and other control measures has just begun.

An integral component of an integrated pest management (IPM) program to control aphids is plant resistance (Auclair, J. L., "Host plant resistance," pp. 225-265 In P. Harrewijn (ed.) *Aphids: Their biology, natural enemies, and control*, Vol. C., Elsevier, New York (1989); Harrewijn, P. and Minks, A. K., "Integrated aphid management: General aspects," pp. 267-272, In A. K. Minks and P. Harrewijn (ed.) *Aphids: Their biology, natural enemies, and control*, Vol. C., Elsevier, New York (1989). Insect resistance can significantly reduce input costs for producers (Luginbill, J. P., "Developing resistant plants—The ideal method of controlling insects," (1969) USDA, ARS. Prod. Res. Rep. 111, USGPO, Washington, D.C. Resistance was reported in *G. soja* (Sun, Z. et al., "Study on the uses of aphid-resistant character in wild soybean. I. Aphid-resistance performance of F<sub>2</sub> generation from crosses between cultivated and wild soybeans," (1990) *Soybean Genet. News* 17:43-48), a close relative of *G. max* (Hymowitz, T., "On the domestication of the soybean," (1970) *Econ. Bot.* 24:408-421), and other wild relatives (Zhuang, B. et al., "A study on resistance to soybean mosaic virus and *Aphis glycines* of perennial wild soybean," (1996) *Soybean Genet. News.* 23:66-69). There are no reports of resistance in *G. max*. A report from Indonesia indicated that there was no resistance in a test of 201 soybean cultivars and breeding lines (Sama, S. et al. (1974) *Research Reports* 1968-1974, p. 171-172. In Varietal screening for resistance to the aphid, *Aphis glycines*, in soybean. Agricultural Cooperation, Indonesia, the Netherlands).

There are numerous examples of the discovery and use of resistance genes to control aphids in crops other than soybean. Examples include Russian wheat aphid (Du Toit, F. (1987), "Resistance in wheat (*Triticum aestivum*) to *Diuraphis noxia* (Homoptera: Aphididae)," *Cereal Res. Commun.* 15:175-179; wheat greenbug (Tyler, J. M., et al. (1985), "Biotype E greenbug resistance in wheat streak mosaic virus-resistant wheat germplasm lines," *Crop Science* 25:686-688), potato aphid on tomato (Kaloshian, I., et al. (1997), "The impact of Meu-1-mediated resistance in tomato on longevity, fecundity and behavior of the potato aphid," *Macrosiphum euphorbiae*," *Entomol. Exp. Appl.* 83:181-187), and cotton-melon aphid on melon (Klinger, J. et al. (2001), "Mapping of cotton-melon aphid resistance in melon," *J. Am. Soc. Hortic. Ci.* 136:56-63).

A number of soybean markers have been mapped and linkage groups created, as described in Cregan, P. B., et al., "An Integrated Genetic Linkage Map of the Soybean Genome" (1999) *Crop Science* 39:1464-1490.

All publications referred to herein are incorporated herein by reference to the extent not inconsistent herewith.

Methods and molecular tools are needed to allow breeding of *A. glycines* resistance into high-yielding *G. max* soybean varieties.

## SUMMARY OF THE INVENTION

A novel method is provided for determining the presence or absence in a soybean germplasm of a gene for resistance to

the soybean aphid, *Aphis glycines*. The aphid resistance trait has been found to be closely linked to a number of molecular markers that map to linkage group M. Genes found on soybean linkage group M conferring the resistance trait are designated Rag1. The Rag1 gene was originally discovered in the resistance sources Dowling (PI548663) and Jackson (PI548657). ("PI" stands for "plant introductions" and these PI numbers refer to USDA depositary accession numbers.) The trait of resistance to *Aphis glycines* is also found in other varieties as described hereafter.

In accordance with the present invention, the gene for resistance to *Aphis glycines* (the RAG gene) co-segregates with molecular markers with which it is linked on linkage group M, most preferably, Satt435, Satt463, Satt245, and DOP\_H14. The Rag1 gene found on Dowling and Jackson, has been found to map to a locus that lies between the markers Satt435 and Satt463. Other markers of linkage group M may also be used to identify the presence or absence of the gene. Preferably flanking markers are used for identifying the presence of a RAG gene or for marker-assisted breeding. Most preferably, the markers used map within about 20 cM, and more preferably within about 10 cM of a RAG locus (which contains the Rag1 gene), or within about 20 cM and more preferably within about 10 cM of Satt435 or Satt463.

The information disclosed herein regarding RAG loci is used to aid in the selection of breeding plants, lines and populations containing *Aphis glycines* resistance for use in introgression of this trait into elite soybean germplasm, or germplasm of proven genetic superiority suitable for variety release.

Also provided is a method for introgressing a soybean *Aphis glycines* resistance gene into non-resistant soybean germplasm or less resistant soybean germplasm. According to the method, nucleic acid markers linked to a RAG gene are used to select soybean plants containing a RAG locus. Plants so selected have a high probability of expressing the trait *Aphis glycines* resistance. Plants so selected can be used in a soybean breeding program. Through the process of introgression, the RAG locus is introduced from plants identified using marker-assisted selection to other plants. According to the method, agronomically desirable plants and seeds can be produced containing the RAG locus from germplasm containing a RAG gene.

Particular examples of sources of Rag1 resistance to *A. glycines* are the following *G. max* varieties: Dowling (PI548663) and its grandparent CNS (PI548445), Jackson (PI548657), and its parent Palmetto (PI548480). PI071506 is also a source of *A. glycines* resistance.

Other sources of *A. glycines* resistance are disclosed below.

Also provided herein is a method for producing an inbred soybean plant adapted for conferring, in hybrid combination, *Aphis glycines* resistance. First, donor soybean plants for a parental line containing a RAG gene are selected. According to the method, selection can be accomplished via nucleic acid marker-associated selection as explained herein. Selected plant material may represent, among others, an inbred line, a hybrid, a heterogeneous population of soybean plants, or simply an individual plant. According to techniques well known in the art of plant breeding, this donor parental line is crossed with a second parental line. Preferably, the second parental line is high yielding. This cross produces a segregating plant population composed of genetically heterogeneous plants. Plants of the segregating plant population are screened for the RAG locus. Those plants having the RAG locus are selected for further breeding until a line is obtained that is homozygous for resistance to *Aphis glycines* at the RAG locus. This further breeding may include, among other tech-

niques, additional crosses with other lines, hybrids, backcrossing, or self-crossing. The result is an inbred line of soybean plants that are resistant to *Aphis glycines* and also have other desirable traits from one or more other inbred lines.

Soybean plants, seeds, tissue cultures, variants and mutants having *Aphis glycines* resistance produced by the foregoing methods are also provided in this invention.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows a published soybean genetic linkage M composite map and anchored markers, in which SSR markers used to create the linkage map with the RAG gene indicated by horizontal lines. The map has been broken into four consecutive vertical sections, FIG. 1A through FIG. 1D.

FIG. 2 is a linkage map of soybean linkage group M (LGM) showing the locations of the soybean aphid resistance genes Rag1 gene A, mapped in a Dowling×Loda mapping population; B, mapped in a Dowling×Williams 82 mapping population; and C. Integrated map from Dowling×Loda and Dowling×Williams 82 mapping populations.

FIG. 3. SSR markers showed co-segregation patterns with Rag1. The ideal PCR amplified band pattern of a co-segregated marker would show A, B, B, B as corresponding to R (resistant parent Dowling, D), S (susceptible parent Loda, L), S (susceptible bulk 1, S1), S (susceptible bulk 2, S2) phenotypes. Among the screened markers, four markers on soybean linkage group M showed potential co-segregation with Rag1. Satt245 showed A, B, B, B pattern, Satt150, Satt220, and Satt323 showed A, B, B, H patterns.

#### DETAILED DESCRIPTION

"Allele" is any of one or more alternative forms of a gene, all of which alleles relate to one trait or characteristic. In a diploid cell or organism, the two alleles of a given gene occupy corresponding loci on a pair of homologous chromosomes. The RAG genes in Dowling and Jackson may be allelic to each other.

"Backcrossing" is a process through which a breeder repeatedly crosses hybrid progeny back to one of the parents (recurrent parent), for example, a first generation hybrid F<sub>1</sub> with one of the parental genotypes of the F<sub>1</sub> hybrid.

"Cultivar" and "variety" are used synonymously and mean a group of plants within a species (e.g., *Glycine max*) that share certain genetic traits that separate them from the typical form and from other possible varieties within that species. Soybean cultivars are inbred lines produced after several generations of self-pollination. Individuals within a soybean cultivar are homogeneous, nearly genetically identical, with most loci in the homozygous state.

"Gene" means a specific sequence of nucleotides in DNA that is located in the germplasm, usually on a chromosome, and that is the functional unit of inheritance controlling the transmission and expression of one or more traits by specifying the structure of a particular polypeptide or controlling the function of other genetic material. In the present instance, RAG genes for resistance to *Aphis glycines* (RAG) have been found on RAG loci flanked by markers Satt435 and Satt463. The RAG gene is referred to as Rag1 when derived from or identical to the Dowling variety and when derived from or identical to the Jackson variety. RAG genes may be isolated by one skilled in the art without undue experiments by means known to the art including PCR cloning utilizing the adjacent Satt435 and Satt463 primer sequences, or primer sequences from other markers flanking the gene as described herein, by

positional cloning using BACs (bacterial artificial chromosomes), or other methods. See, e.g., Wu, et al., "A BAC and BIBAC-based Physical Map of the Soybean Genome" (2004) *Genome Res.* February; 14(2):319-26, which describes the use of BACs in mapping the soybean genome. Contiguous BACs that have been found to be anchored to Satt435 and in which the Rag1 gene may be found include B03124<sup>-</sup>, B52J11\*, B431224<sup>-</sup>, H57B23, H03008, B36M08\*, H62M17, H75H01, and E71J17. Information on these contiguous BACs is known to the art. Certain information is publicly available at the National Center for Biotechnology Information (NCBI) and GenBank web sites. The end sequence for H03008 is set forth below:

```
H03008:
[SEQ ID NO: 1]
AAGCTTCTAT CAAGTGGTAA TCAGAGCACA AGATCTTCAA
GTAGGTGATC CTTAAACCTC CATTAAATTTT TTGCTTTACC
TTCTCTTCTA TTGTTGTTTC TTCATTTTTC TCCATGTATC
TCCTCACATG TCTTGTGCTA AATGTTTTTA ACATGATTCT
TTAGAGTTTC CACCGATTAA ACTTGCTATA GAAGCTAGAT
TTGATTTTCT ATGGTTCAAA TTTCTTGTTT TTGTTCTTGA
TCCATGAATT GTGTTGAGTT TAGGTTCCCTT TGAGTTTGTG
CTTGTTATTT TTTGTGGCTG AACCTAAAC CATAAAATTC
TTACAAAAAT ATTAAAGTAG AGGAAAACCT CAAAAATCTA
GAGTGACTTG TTCACCTATT ATAGTTTGTG CATAGAAGTC
ATGTCCTAGTC ATGAAACTTG TCACATAAGA TTTCTTATGT
TGTGCTGAAT TTTATTTTCT TGTTTCTTTG TCTAACTCAT
TTGTTTATGA GTGTATGAAG TTATTTTAGC CTATTATTTG
ATTGGAGTCA AATCTTTCAT GTTAATTAGT CCTTAACATG
TTCATGCAAA ATTCTTAGAG AGTCTTTGAT TGTGAACCTT
TTCTTGAAGT TTTAGGTTTC CTTATGATTG TGTCTATTGT
GAATTTAAGT TTTGGTGATT GAATTGCTGG TTGAAATGTT
GATCCTAAGT GAATATTGAA CTCCTAAAC TGTGGTAAAC
AATCCTAGTG AGTTCAACAT ACATAGGAAG GTTGAAAGTA
AGCCCAAGGC AATCAATATA GCATGCTTAA AAAAAAATC
GCTGGTGCTG GCAGCTTGGA CATACAAACT TGTAATAAAT
ACTGAAAATT GGTACTTCG AATTTGAAC TGAATTTTTA
CTTAATTTGC TAGA
```

"Germplasm" means the genetic material with its specific molecular and chemical makeup that comprises the physical foundation of the hereditary qualities of an organism. As used herein, germplasm includes seeds and living tissue from which new plants may be grown; or, another plant part, such as leaf, stem, pollen, or cells; that may be cultured into a whole plant. Germplasm resources provide sources of genetic traits used by plant breeders to improve commercial cultivars.

"Hybrid plant" means a plant offspring produced by crossing two genetically dissimilar parent plants.

"Inbred plant" means a member of an inbred plant strain that has been highly inbred so that all members of the strain are nearly genetically identical.

"Introgression" means the entry or introduction by hybridization of a gene or trait locus from the genome of one plant into the genome of another plant that lacks such gene or trait locus.

"Molecular marker" is a term used to denote a nucleic acid or amino acid sequence that is sufficiently unique to characterize a specific locus on the genome. Examples include restriction fragment length polymorphisms (RFLPs) and single sequence repeats (SSRs). RFLP markers occur because any sequence change in DNA, including a single base change, insertion, deletion or inversion, can result in loss (or gain) of a restriction endonuclease recognition site. The size and number of fragments generated by one such enzyme is therefore altered. A probe that hybridizes specifically to DNA in the region of such an alteration can be used to rapidly and specifically identify a region of DNA that displays allelic variation between two plant varieties. SSR markers occur where a short sequence displays allelic variation in the number of repeats of that sequence. Sequences flanking the repeated sequence can serve as polymerase chain reaction (PCR) primers. Depending on the number of repeats at a given allele of the locus, the length of the DNA segment generated by PCR will be different in different alleles. The differences in PCR-generated fragment size can be detected by gel electrophoresis. Other types of molecular markers are known. All are used to define a specific locus on the soybean genome. Large numbers of these have been mapped. Each marker is therefore an indicator of a specific segment of DNA, having a unique nucleotide sequence. The map positions provide a measure of the relative positions of particular markers with respect to one another. When a trait is stated to be linked to a given marker it will be understood that the actual DNA segment whose sequence affects the trait generally co-segregates with the marker. More precise and definite localization of a trait can be obtained if markers are identified on both sides of the trait. By measuring the appearance of the marker(s) in progeny of crosses, the existence of the trait can be detected by relatively simple molecular tests without actually evaluating the appearance of the trait itself, which can be difficult and time-consuming, requiring growing up of plants to a stage where the trait can be expressed.

Another type of molecular marker is the random amplified polymorphic DNA (RAPD) marker. Chance pairs of sites complementary to single octa- or decanucleotides may exist in the correct orientation and close enough to one another for PCR amplification. With some randomly chosen decanucleotides no sequences are amplified. With others, the same length products are generated from DNAs of different individuals. With still others, patterns of bands are not the same for every individual in a population. The variable bands are commonly called random amplified polymorphic DNA (RAPD) bands.

Another type of molecular marker is the target region amplification polymorphism (TRAP) marker. The TRAP technique employs one fixed primer of known sequence in combination with a random primer to amplify genomic fragments.

A further type of molecular marker is the single nucleotide polymorphism (SNP) marker, in which DNA sequence variations that occur when a single nucleotide (A, T, C, or G) in the genome sequence is altered are mapped to sites on the soybean genome.

Other molecular markers known to the art, as well as phenotypic traits may be used as markers in the methods of this invention.

"Linkage" is defined by classical genetics to describe the relationship of traits that co-segregate through a number of

generations of crosses. Genetic recombination occurs with an assumed random frequency over the entire genome. Genetic maps are constructed by measuring the frequency of recombination between pairs of traits or markers. The closer the traits or markers lie to each other on the chromosome, the lower the frequency of recombination, the greater the degree of linkage. Traits or markers are considered herein to be linked if they generally co-segregate. A 1/100 probability of recombination per generation is defined as a map distance of 1.0 centiMorgan (1.0 cM). Preferably markers useful for screening for the presence of *Aphis glycines* resistance (RAG) map to within 20 cM of the trait, and more preferably within 10 cM of the trait.

A second marker that maps to within 20 cM of a first marker that co-segregates with the RAG trait and generally co-segregates with the RAG trait is considered equivalent to the first marker. Any marker that maps within 20 cM and more preferably 10 cM of the RAG trait belongs to the class of preferred markers for use in screening and selection of soybean germplasm having the RAG *Aphis glycines* resistance trait. A number of markers are known to the art to belong to linkage group M on which the RAG trait is found. A number of markers are proprietary markers known only to certain of those skilled in the art of soybean plant breeding. A proprietary marker mapping within 20 cM, and preferably within 10 cM, of any publicly known marker specified herein is considered equivalent to that publicly-known marker.

“Linkage group” refers to traits or markers that generally co-segregate. A linkage group generally corresponds to a chromosomal region containing genetic material that encodes the traits or markers.

“Locus” means a chromosomal region where a polymorphic nucleic acid or trait determinant or gene is located.

“Polymorphism” means a change or difference between two related nucleic acids. A “nucleotide polymorphism” refers to a nucleotide that is different in one sequence when compared to a related sequence when the two nucleic acids are aligned for maximal correspondence. A “genetic nucleotide polymorphism” refers to a nucleotide that is different in one sequence when compared to a related sequence when the two nucleic acids are aligned for maximal correspondence, where the two nucleic acids are genetically related, i.e., homologous, for example, where the nucleic acids are isolated from different strains of a soybean plant, or from different alleles of a single strain, or the like.

“Marker assisted selection” means the process of selecting a desired trait or desired traits in a plant or plants by detecting one or more nucleic acids from the plant, where the nucleic acid is linked to the desired trait.

“Plant” means plant cells, plant protoplast, plant cell or tissue culture from which soybean plants can be regenerated, plant calli, plant clumps and plant cells that are intact in plants or parts of plants, such as seeds, pods, flowers, cotyledons, leaves, stems, buds, roots, root tips and the like.

“Probe” means an oligonucleotide or short fragment of DNA designed to be sufficiently complementary to a sequence in a denatured nucleic acid to be probed and to be bound under selected stringency conditions.

“Rag1-derived resistance” means resistance in a soybean germplasm to *Aphis glycines* that is provided by the heterozygous or homozygous expression of the Rag1 gene within the RAG locus mapped between the SSR markers Satt435 and Satt463. “RAG-derived resistance” means *Aphis glycines* resistance conferred by a RAG gene on a RAG locus, the use of which is enabled by the disclosure herein.

“RAG phenotype” means resistance to *Aphis glycines* by soybean germplasm, as demonstrated by resistance to *Aphis glycines* after inoculation with same according to the methods described herein.

“RAG soybean plant” means a plant having resistance to *Aphis glycines* that is derived from the presence and expression of at least one RAG gene, or that is shown to have a RAG gene at the RAG locus described herein.

“Self-crossing or self-pollination” is a process through which a breeder crosses hybrid progeny with itself, for example, a second generation hybrid F<sub>2</sub> with itself to yield progeny designated F<sub>2,3</sub>.

As used herein, the terms “segregate,” “segregants,” “co-segregate,” “hybrid,” “crossing,” and “selfing” refer to their conventional meanings as understood in the art (see, for instance, Briggs, F. N. and Knowles, P. F. and, *Introduction to Plant Breeding* (Reinhold Publication Corp., New York, N.Y., 1967).

Markers that “flank” the RAG genes are markers that occur one to either side of a RAG gene. Flanking marker DNA sequences may be part of the gene or may be separate from the gene.

The method for determining the presence or absence of a RAG gene, which confers resistance to the soybean aphid *Aphis glycines* in soybean germplasm, comprises analyzing genomic DNA from a soybean germplasm for the presence of at least one molecular marker, wherein at least one molecular marker is linked to the RAG trait locus, and wherein the RAG trait locus preferably maps to soybean major linkage group M and is associated with resistance to the soybean aphid *Aphis glycines*. The term “is associated with” in this context means that the RAG locus containing the RAG gene has been found, using marker-assisted analysis, to be present in soybean plants showing resistance to *Aphis glycines* in live aphid bioassays as described herein.

The Rag1 gene occurs in the following varieties CNS (PI548445), and Dowling (PI548663), Jackson (PI548657), and Palmetto (PI548480), among others.

Other sources of *A. glycines* resistance include the *G. max* varieties: Moyashimame (PI87059), Sato (PI548409), Showa No. 1-4 (PI88508), Sugao Zarai (PI200538), T26OH (PI548237), PI71506, and PI230977 of *G. max*, and G3, JS1, I4, S12 Taichung 38 (PI518282 and Taichung 37 (PI518281), of *G. soja*, and progeny of these varieties.

Table 1 lists *Glycine max.* varieties that are sources of resistance to the soybean aphid. Progeny of these varieties also containing a RAG gene are also sources of resistance to the soybean aphid.

TABLE 1

SOURCES OF RESISTANCE TO SOYBEAN APHID		
PI#	Name	
71506		
87059	Moyashimame	
88508	Showa No. 1-4	
200538	Sugao Zarai	
230977		
417084A	Kumaji 1	
437696	San-haj-hun-mao-huan-dou	
499955		
507298	Sokoshin (Kamigoumura)	
508294		

TABLE 1-continued

SOURCES OF RESISTANCE TO SOYBEAN APHID	
PI#	Name
518726	Bao jiao huang
548237	T260H
548409	Sato
548445	CNS
548480	Palmetto
548657	Jackson
548663	Dowling
567391	Jiang se huang dou
567541B	
567543C	
567597C	
567598B	
587552	Nan jing da ping ding huang yi 1
587553A	
587559B	(Dan tu he shang tou jia)
587617	Jin tan qing zi
587656	Huang dou
587663	Zhong chun huang dou
587664B	(Shan zi bai)
587666	Er dao zao
587668A	Hui mei dou
587669	Zan zi bai
587674A	Ba yue bai
587677	Xiao li huang
587682A	Da li huang 1
587684A	Ai jiao huang
587685	Da li huang 2
587686A	Xi li huang 1
587687A	Xiao li dou 1
587693	Yu shan dou
587700A	Da qing dou
587702	Qing pi dou
587717	Xiang yang ba yue zha
587723A	Ying shan ji mu wo
587732	Ying shan ji mu wo
587759	Song zi ba yue cha
587763	Jing huang 36
587775	Tong shan si ji dou
587800	Ying shan da li huang
587816	Bai mao dou
587824	Ying shan qing pi cao
587840	Du wo dou
587844C	(Tong cheng hei se dou)
587861	Da qing dou
587863B	(Liu yue bai)
587870	Huang pi dou
587871	Bao mao dou
587873	Feng wo dou
587876	Xi mao dou
587877A	Jiu yue zao
587891A	Qi yue ba
587897	Qing pi dou
587899	Ba yue bai
587905	Xiao huang dou
587972	Chang zi dou
588000	Shi yue huang
588040	Shan xing dou
594421	Da du huang dou
594425	Xiao cao huang dou
594426A	Tie jiao huang
594426B	(Tie jiao huang)
594427A	Ba yue mang
594431	Chang pu qing dou
594499	Luo ma aluo
594503	Mu gu hei chi huang dou
594514	Hua lian dou
594554	Huang pi tian dou
594557B	(Lao shu dou)
594560B	(Xia shui huang)
594573	Lu pi dou
594586A	
594592	Shi yue xiao huang dou
594595	Ba yue da huang dou (jia)
594666B	(Liu yue mang -5)
594703	Qing pi dou -1
594707	Da hei dou

TABLE 1-continued

SOURCES OF RESISTANCE TO SOYBEAN APHID	
PI#	Name
594711B	(Qing huang za dou -3)
594751A	Long zhou dong feng dou
594822	Xi huang dou
594864	Yang yan dou
594868	Huang dou
594879	Huo shao dou
603521	
603530A	
603538A	
603640	
603644	
603650	
605771	
605823	
605855	
605902	

25 The following *G. soja* varieties are also sources of *A. gly-*  
*cines* resistance: PI441008, PI573059, and PI573071, and  
progeny of these varieties.

30 Any one of the foregoing varieties or their progeny bearing  
a RAG gene may be used in the methods of this invention, and  
any combination thereof is considered to be a class of varieties  
useful in the methods of this invention.

35 Preferably a marker used to determine the presence or  
absence of a RAG gene is Satt435, Satt463, Satt245, S04309,  
S01623, or DOP\_H14, or a marker that maps to within at least  
about 10 to about 20 cM of any of said markers.

40 Any marker assigned to soybean linkage group M may be  
useful for this purpose. Exemplary markers of linkage group  
M include Sat\_389, Satt404, Sat\_391, GMSC514, Satt636,  
Satt590, GM177, GM175, Satt201, Satt150, Sat\_316,  
A351\_1, Mng339\_1, Ts, S01256, S02020, Satt567,  
45 Satt540, Bng222\_1, RGA\_2b, RGA5b, GM260, S04309,  
Satt435, R079\_1, A060-2, DOP\_H14, GM260, A131\_1,  
Sat\_244, S01623, Satt463, Sat\_253, S03544, Satt245,  
GM284, A946\_2, GM256, GMS057, Satt220, A584\_3,  
Satt626, OP\_N04, Satt323, Sat\_258, Satt702, L204\_4,  
50 GMS003b, Satt536, Sat\_003, OM11\_1100, Sat\_148,  
Bng179\_2, K417\_3, Sat\_226, Satt175, K024\_1, A226\_1,  
GM230, ACCAGC315, Satt494, B157\_2, A715\_1, Sct\_  
147, Sat\_256, Satt677, Sat\_288, Satt655, Satt680,  
55 AF186183, BE823543, Satt306, A458\_4, Satt728, Sat\_422,  
E043\_1, Satt697, Mng186\_1, GM163, K070\_1, AC\_1,  
BLT025\_2, Cr326\_3, Satt551, M121\_1, Satt551, Sat\_131,  
Satt250, Satt618, Satt210, Satt346, K227\_1, Sat\_147,  
A064\_1, GM141, GM209b, GM035a, A504\_1, Sat\_276,  
60 Satt308, Satt336, Mng381\_1, Sat359, Sat\_330, and A504.1.

Updated information regarding markers assigned to soy-  
bean linkage group M may be found on the USDA's Soybase  
website. Table 2 provides current information on the genbank  
location and allele size of markers useful in this invention.  
Table 3 provides upper and lower primer sequences.

TABLE 2

MARKERS					
Name	Type	GenBank gi #	cM Position in linkage group	GenBank Accession #	Allele Size in Williams
Sat_389	SSR	31044744	0.00	CC453914	
Satt404	SSR	14970089	0.84	BH126586	181
Sat_391	SSR	31044746	1.02	CC453916	
GMSC514	SSR	18745	3.05	X56139	160
Satt636	SSR	31044825	5.00	CC453995	172
Satt590	SSR	14970259	7.84	BH126756	318
Satt201	SSR	14969911	13.56	BH126408	282
Satt150	SSR	14969865	18.58	BH126362	201
Sat_316	SSR	31044677	21.00	CC453847	298
A351_1	RFLP		22.394		
Mng339_1	RFLP		27.325		
Ts	UNKNOWN		30.251		
Satt567	SSR	14970236	33.493	BH126733	110
Satt540	SSR	14970211	35.85	BH126708	152
Bng222_1	RFLP		38.504		
RGA2B	RFLP		38.679		
Satt435	SSR	14970116	38.94	BH126613	286
R079_1	RFLP		40.354		
A0560_2	RFLP		40.354		
DOP-H14	RAPD	41.836	41.836		
A131_1	RFLP		47.12		
Sat_244	SSR	31044612	48.86	CC453782	224
Satt463	SSR	14970139	50.10	BH126636	226
Sat_253	SSR	31044619	51.60	CC453789	275
Satt245	SSR	14969948	53.54	BH126445	211
A946_2	RFLP		55.492		
Satt220	SSR	14969926	56.29	BH126423	245
A584.3	RFLP		58.501		
Satt626	SSR	31044818	58.60	CC453988	238
OP_N04	RAPD		59.11		
Satt323	SSR	14970017	60.05	BH126514	156
Sat_258	SSR	31044623	60.47	CC453793	193
Satt702	SSR	31044881	61.04	CC454051	
L204_4	RFLP		61.26		
Satt536	SSR	14970207	62.14	BH126704	162
Sat_003	SSR	14969756	62.31	BH126253	161
Sat_148	SSR	31044530	63.93	CC453700	162
Bng179_2	RFLP		65.213		
K417_3	RFLP		65.694		
Sat_226	SSR	31044595	65.79	CC453765	212
Satt175	SSR	14969887	66.99	BH126384	163
K024_1	RFLP		71.05		
A226_1	RFLP		71.094		
Satt494	SSR	14970168	71.71	BH126665	218
A715_1	RFLP		73.373		
Sct_147	SSR	14970282	73.88	BH126779	
Sat_256	SSR	31044622	74.53	CC453792	253
Satt677	SSR	31044860	75.57	CC454030	157
Sat_288	SSR	31044651	76.41	CC453821	215
Satt655	SSR	31044840	76.41	CC454010	287
Satt680	SSR	31044863	77.19	CC454033	304
AF186183	SSR	6671123	77.24	AF186183	
BE823543	SSR	10255728	78.38	BE823543	
Satt306	SSR	14970000	80.02	BH126497	212
Satt728	SSR	31044900	80.90	CC454070	
Sat_422	SSR	31044776	80.97	CC453946	
E043_1	RFLP		82.645		
Satt697	SSR	31044876	85.35	CC454046	302
Mng186_1	RFLP		85.433		
K070_1	RFLP		90.921		
BLT025_2	RFLP		93.941		
Satt551	SSR	14970221	95.45	BH126718	238
M121_1	RFLP		96.222		
Sat_121	SSR	14969794	103.98	BH126291	189
Satt250	SSR	14969951	107.70	BH126448	202
Satt618	SSR	31044812	111.06	CC453982	117
Satt210	SSR	14969919	112.08	BH126416	260
Satt346	SSR	14970039	112.79	BH126536	208
K227_1	RFLP		120.373		
Sat_147	SSR	31044529	122.37	CC453699	265
A064_1	RFLP		124.212		
Sat_276	SSR	31044640	128.48	CC453810	271
Satt308	SSR	14970002	130.76	BH126499	170
Satt336	SSR	14970030	133.83	BH126527	170
Mng381_1	RFLP		139.46		

TABLE 2-continued

MARKERS					
Name	Type	GenBank gi #	cM Position in linkage group	GenBank Accession #	Allele Size in Williams
Sat_359	SSR	31044715	139.81	CC453885	
Sat_330	SSR	31044687	140.69	CC453857	265
A504_1	RFLP		142.184		

TABLE 3

MARKER SEQUENCES		
Name	Upper primer sequence (5'-->3')	Lower primer sequence (5'-->3')
Sat_389	GCGGGTAGCCATATTCATATAT TGCTG [SEQ ID NO: 2]	GCGAAGGCTTATAAGGAGATA CGATTTA [SEQ ID NO: 3]
Satt404	TCATCCGCCATTGATTTT [SEQ ID NO: 4]	GCCCGGAACATACAAAA [SEQ ID NO: 5]
Sat_391	GCGTAGGCATCGTCAATATTT T [SEQ ID NO: 6]	GCGTTAGCGAGTGGATCAAGA TCA [SEQ ID NO: 7]
GMSC514	TACCTTTCTGTGAGTCGTA [SEQ ID NO: 8]	TATTGAGATGGATATTGTAGAT C [SEQ ID NO: 9]
Satt636	GTCATGACTCATGAGTCACGTA AT [SEQ ID NO: 10]	CCCAAGACCCCATTTTATGT CT [SEQ ID NO: 11]
Satt590	GCGCGCATTTTTTAAGTTAATGT TCT [SEQ ID NO: 12]	GCGCGAGTTAGCGAATTATTTG TC [SEQ ID NO: 13]
Satt201	GCGTTGATACTTTCCTAAGACA AT [SEQ ID NO: 14]	GGGAGAGAAGGCAATCTAA [SEQ ID NO: 15]
Satt150	AAGCTTGAGGTTATTCGAAAAT GAC [SEQ ID NO: 16]	TGCCATCAGGTTGTGTAAGTGT [SEQ ID NO: 17]
Sat_316	GCGCAACGCTTAAAGCACAAGG ATT [SEQ ID NO: 18]	GCGCGACTACGTTACAGTTCC AA [SEQ ID NO: 19]
Satt567	GGCTAACCCGCTCTATGT [SEQ ID NO: 20]	GGGCCATGCACCTGCTACT [SEQ ID NO: 21]
Satt540	CTGGCGAATCAAGCTTTGTAAC [SEQ ID NO: 22]	CCGTGATTGCGAAGAGGATAT T [SEQ ID NO: 23]
Satt435	GCGGTGAAACGGCTCTCTTTGA TAGTGA [SEQ ID NO: 24]	GCGTTGGATTAATTAATTAAAT TATTTT [SEQ ID NO: 25]
Sat_244	GCGTCAACCGGTGAAAAACCT A [SEQ ID NO: 26]	GCGTGGCTGGCAGTAGTCTAT ATCA [SEQ ID NO: 27]
Satt463	TTGGATCTCATATTCAAACCTTC AAG [SEQ ID NO: 28]	CTGCAAATTGATGCACATGTG TCTA [SEQ ID NO: 29]
Sat_253	GCGATTGGTTGGGTGTTTAATT TTAAGAT [SEQ ID NO: 30]	GCGTGTGATGGTATAAAGATC GCTACTCT [SEQ ID NO: 31]
Satt245	AACGGGAGTAGGACATTTTATT [SEQ ID NO: 32]	GCGCCTCCTGAATTTCAAAGAA TGAAGA [SEQ ID NO: 33]
Satt220	GAGGAGGATCCCAAGGTAATAA T [SEQ ID NO: 34]	GCGCATGGAGAAAAGAAGAG [SEQ ID NO: 35]
Satt626	GCGGATGGAGACGGGGGAC GGACGA [SEQ ID NO: 36]	GCGCATAGCTAATTTTATATCA ATTAT [SEQ ID NO: 37]
Satt323	GCGGTCGTCCTATCTAATGAAG AG [SEQ ID NO: 38]	TGTGCGTTTAAATTGCAGCTAA AT [SEQ ID NO: 39]
Sat_258	GCGCAATAGATAATCGAAAAAC ATACAAGA [SEQ ID NO: 40]	GCGGGGAAAGTAAAACAAGA TCAAATA [SEQ ID NO: 41]
Satt702	GCGGGGTCTGTGGCTTCAAC [SEQ ID NO: 42]	GCGCATTGGAATAACGTCAAA [SEQ ID NO: 43]

TABLE 3-continued

MARKER SEQUENCES		
Name	Upper primer sequence (5'-->3')	Lower primer sequence (5'-->3')
Satt536	GCGCCACAGAAATTCCTTTTTC TA [SEQ ID NO: 44]	GCGCCATAAGGTGGTTACCAA AAGA [SEQ ID NO: 45]
Sat_003	TGATTTTGGTGTAGAACTC [SEQ ID NO: 46]	CAAATTGGTTAGCTTACTCCA [SEQ ID NO: 47]
Sat_148	GCGGAGTTTCCCCTAATTAGAT [SEQ ID NO: 48]	GCGCAAGCTAGCTTCACCCAA AACTA [SEQ ID NO: 49]
Sat_226	GCGGAAACCCACCTATATGTGA TCAAAATG [SEQ ID NO: 50]	GCGCAATTCCAGATGAAACAG AAGAAGGAT [SEQ ID NO: 51]
Satt175	GACCTCGCTCTCTGTTTCTCAT [SEQ ID NO: 52]	GGTGACCACCCCTATTCCTTAT [SEQ ID NO: 53]
Satt494	GGCCGGTTCCTATTACAGGTCT CT [SEQ ID NO: 54]	GGATTTCATCTTGAATTTTATT A [SEQ ID NO: 55]
Sct_147	TCTCGACTCAGACTCA [SEQ ID NO: 56]	CCAAGGTCTCTCAGAGG [SEQ ID NO: 57]
Sat_256	GCGCGGAAAATTATTTTACTTTT TCAAT [SEQ ID NO: 58]	GCGCACGGATTGAGAGAAAGC AGAAAGA [SEQ ID NO: 59]
Satt677	CAACGACCAACTGACGAGACCT [SEQ ID NO: 60]	GGGAATTCAACATGTGATGGTT TT [SEQ ID NO: 61]
Sat_288	GCGACAGACTGCAAGAATTGAT GTAAATCT [SEQ ID NO: 62]	GCGGGAAGGTAGGTAAAGAAA ATTCAAATGA [SEQ ID NO: 63]
Satt655	GAAGACCAAACTTATTTTCAGAT C [SEQ ID NO: 64]	ATTTTAAGCACCAGCAAAGACT [SEQ ID NO: 65]
Satt680	GCGGGATATCGTGAGCATAGTT TTAC [SEQ ID NO: 66]	GCGGCCTGAATATTTTAGGTTT AGAGTT [SEQ ID NO: 67]
AF186183	GCGTATTTTGGGGGATTTTGAA CA [SEQ ID NO: 68]	GCGTTTCTCTTCTTATTCTTTCT CT [SEQ ID NO: 69]
BE823543	GCGAAATGCCGAAAGAG [SEQ ID NO: 70]	GCGGGGATAAGAAAAACAAT [SEQ ID NO: 71]
Satt306	GCGCTTAAGGACACGGATGTAA C [SEQ ID NO: 72]	GCGTCTCTTTCGATTGTTCTAT TAG [SEQ ID NO: 73]
Satt728	GCGTACCCCTATATGGATGTTT CTTCCT [SEQ ID NO: 74]	GCGTATGCAGCAACAAAAAAT ATATAAT [SEQ ID NO: 75]
Sat_422	GCGTTTTCTTAATGAAGATT [SEQ ID NO: 76]	GCGTGTAATAGTGATGGATGTA A [SEQ ID NO: 77]
Satt697	GCGTGCTTTAAATGATTGATTG A [SEQ ID NO: 78]	GCGTGCGAACATAACTAATACA T [SEQ ID NO: 79]
Satt551	GAATATCACGCGAGAATTTTAC [SEQ ID NO: 80]	TATATGCGAACCTCTTACAAT [SEQ ID NO: 81]
Sat_121	GACAAATGTAAAAAGTGACAGA TAGAATGT [SEQ ID NO: 82]	GTGTGGTGGTGGTACAGTTTTA TACTAA [SEQ ID NO: 83]
Satt250	CGCCAGCTAGCTAGTCTCAT [SEQ ID NO: 84]	AATTTGCTCCAGTGTTTAAAGT TT [SEQ ID NO: 85]
Satt618	GCGGTGATATTACCCCAAAAAA ATGAA [SEQ ID NO: 86]	GCGCTAGTTTCTAGTGGAAAG ATGAGT [SEQ ID NO: 87]
Satt210	GCGAAAAACGTCAGGTCAATGA CTGAAA [SEQ ID NO: 88]	GCGGGGCTTAGATATAAAAAAA AAGATG [SEQ ID NO: 89]
Satt346	GGAGGGAGGAAAGTGTGTGG [SEQ ID NO: 90]	GCGCATGCTTTTCATAAGTTT [SEQ ID NO: 91]
Sat_147	GTGCGACGTCATGCCTTACTCA AT [SEQ ID NO: 92]	GCGCTCCGTACACTTAAAAAAG AA [SEQ ID NO: 93]



TABLE 3-continued

MARKER SEQUENCES		
Name	Upper primer sequence (5'-->3')	Lower primer sequence (5'-->3')
Sat_276	GCGGAACCCATCTAGATATG AAAAACA [SEQ ID NO: 94]	GCGTTCTTCTCGAGGTGAGAT ACAATC [SEQ ID NO: 95]
Satt308	GCGTTAAGGTTGGCAGGGTGG AAGTG [SEQ ID NO: 96]	GCGCAGCTTTATACAAAATCA ACAA [SEQ ID NO: 97]
Satt336	AATTGGAGTGGGTACAC [SEQ ID NO: 98]	TTCCCGAAAGAAAGAAA [SEQ ID NO: 99]
Sat_359	GCGGGTCACGATTCTAGTCACT ATAACTTCA [SEQ ID NO: 100]	GCGCAACGTAAGAAATGTAAAT ACAATGGA [SEQ ID NO: 101]
sat_330	GCGTTAGGATTAGGATGAGGA TAGG [SEQ ID NO: 102]	GCGCAAATCAGTTGAGCAATG ACTTA [SEQ ID NO: 103]

The sequence of the RAPD marker DOP\_H14 is: 5' to 3': ACCAGGTTGG [SEQ ID NO:104].

Table 4 provides information on additional SNP markers that are useful in practicing the present invention, showing their relative locations with respect to the markers described in Tables 2 and 3.

TABLE 4

SNP MARKERS				
Locus	SNP ID	BARC Seq. ID	Type	GenBank source seq. Position in LG
S01256	BARC-GM-01256	13845	3'mRNAsequence	AW348751 33.493
S02020	BARC-GM-02020	15945	3'mRNAsequence	AW349790 33.493
Satt567				33.493
Satt540				35.879
Bng222_1				38.528
RGA_2b				38.703
S04309	BARC-GM-04309	22289	3'mRNAsequence	AW351227 38.964
Satt435				38.964
R079_1				39.004
A060_2				40.378
DOP_H14				41.858
A131_1				47.142
Sat_244				48.876
S01623	BARC-GM-01623	14705	3'mRNAsequence	AW349229 50.117
Satt463				50.117
Sat_253				51.617
S03544	BARC-GM-03544	18283	From subclone of BAC identified with Satt245	53.558
Satt245				53.558

S01256 is available through: Genome Systems, Inc. 4633 World Parkway Circle St. Louis, Mo. The sequence of S01256 and equivalent markers is taken from the 3' end of the following sequence: TATCATTATA TTGCAGGCTA CNNAATTTTC CAGTNNNTAAT ACAGTATAAT TAAG-CAGAGT GTGGTATCTA CAAAATCTCA ATCCAAA-CAC ATAATTACAA AACTCTAGAA CAGCAGAACA CATATAGCAT TTGATTGAA GTATTCATTC ACTAAT-TGAT TAGCCTTAGA AATTCAAATG ATATAATCTG ACCACTCAGA GATAAAGGAA GTATGGTCCA TGGACTCCCC AGGAACATCC TCGTGCTTAG AGGGCTTCTC CTTCACCA ACCAACCTGG CTGGGTTCCC AACAGCTGTT GTCTGTGGTG GCA-CATCGAT TAAAACCACC GAGCCAGCAC AAC-CTTTGC ACCTTCCCCG ATCTTAATAT TCCCCAGAAT GGTAGCACCG GCACCAATAA GCACCCCATC CCCAATCTTG GGATGCCGGT CCCCACCAAC TTGC-

CAGTC CCACCCAGCG TAACGTGGTG CAGGATC-GAC ACATTGTTCC GATCACTGC CGTCTCCCCC ACCACCACCC CGGTGGCATG GTCGAACAGA TCCCCITCC CGATCCTCGC CGCAGGGTGA ATGTC-CACCG CGAACACATC GCGATGCGA GAGTGCAGTG CNAAAGCCAA TGGCTGCCGC GATTGNCNC CAA-

CAGATG CGCCACACGG TGCGCCTGCA AAATCA-CAAT CACACACAAC TAATCCTAAG ATTCAATAAT CAAAAAAGAG TNNACTNNNC ATACACTGTC ATC-NCNNNTA TAGTCATGTT TCATNNNAAT CTNGNNNNAC AATGCATATA AATTAAACTC AAT [SEQ ID NO:105]

S02020 is available through: Genome Systems, Inc. 4633 World Parkway Circle St. Louis, Mo. The sequence of S02020 and equivalent markers is taken from the 3' end of the following sequence: AAAGNNAACA TTTTGTGTTA TAT-GACNNNA ACAAACTGCA AAGAAAAATT GTTAAAAACC AGAAGCAATT TAGGTGATCA CAAATACCAC ATGCTTACAC CTCCAGTGA CAAG-TACAGT ATGTTGTGGC ACCAGCCGTT TCAGT-TGATG CAAACTTGCT TCGTGCCAAA ATTCTAACAA CAAACTACC TAAGCTATCA AACAAGAGAA GCCCTTTTGT CCTTGTGTCG ACCTATCAAA GGT-

19

CATCAGA TCACACTAGT CCTACCCTTT TAA-  
GAAAACC TACTATCAAC AGTCATATGT ATCTCAT-  
GAA AAGCACATAA AAACATGTCA CTTTGCCTCT  
TCACCATCTC CACTGTTATG AGCAGCCGCG GAGCT-  
GCCTT GGCCGTCTCC ACCAGCTGTT CCAGCCTCAG  
AGGCATCTTG CTTGCTTCCA CCACGTGCAT  
CGTTTGGACC AGTAGCCGAA GGTGGACCAC CGCT-  
GTTTCC CCTCCAAGA GCAGCCTCAC TGTGCATTGG  
ATGCATGCCA TTTATTTAT CTCCAGGTCT AAGT-  
CCATT TGACCTTGA TGGCTGCTG GTGTAGCTGC  
TGTGTTGTT CTTGCAATTTG ATGTGGATTG  
CCAAATTGCA ATGGCATTCT CTGGGGGAAC ANNC-  
CTTGCT GCTGCTGNNN NATTGCTGCT  
GCAGCNNNN GNNNATNNNN NATATANNNN NC  
[SEQ ID NO:106]

S04309 is available through: Genome Systems, Inc. 4633  
World Parkway Circle St. Louis, Mo. The sequence of  
S04309 and equivalent markers is taken from the 3' end of the  
following sequence: TATCATTATA TTGCAGGCTA  
CNNAATTTT CAGTNNTAAT ACAGTATAAT TAAG-  
CAGAG GTGGTATCTA CAAATCTCA ATCCAAA-  
CAC ATAATTACAA AACTCTAGAA CAGCAGAACA  
CATATAGCAT TTGATTGAA GTATTCATTC ACTAAT-  
TGAT TAGCCTTAGA AATTCAAATG ATATAATCTG  
ACCACTCAGA GATAAAGGAA GTATGGTCCA  
TGGACTCCCC AGGAACATCC TCGTGCTTAG  
AGGGCTTCTC CTTCACCA ACCAACCTGG  
CTGGGTTCCC AACAGCTGTT GTCTGTGGTG GCA-  
CATCGAT TAAAACCACC GAGCCAGCAC CAAC-  
CTTTC ACCTTCCCCG ATCTTAATAT TCCCCAGAAT  
GGTAGCACCG GCACCAATAA GCACCCCATC  
CCCAATCTTG GGATGCCGGT CCCCACCAAC  
CTTGCCAGTC CAGCAGCG TAACGTGGTG CAG-  
GATCGAC ACATTGTTCC CGATCACTGC CGTCTC-  
CCCC ACCACCACC CGGTGGCATG GTCGAACAGA  
ATCCCCTTCC GATCCTCGC CGCAGGGTGA ATGTC-  
CACCG CGAACACATC AGCGATGCGA AGTGCAGTG  
CNAAGCCAA TGGCTGCCGC GATTGNCGNC  
ACAACAGATG GCCACACGG TGCCTGCA AAAT-  
CACAAT CACACACAAC TAATCCTAAG ATTCAATAAT  
CAAAAAAGAG TNNACTNNNC ATACACTGTC ATC-  
NCNNNTA TAGTCATGTT TCAATNNNAAT  
CTNGNNNNAC AATGCATATA AATTAACTC AAT  
[SEQ ID NO:107]

S01623 is available through: Genome Systems, Inc. 4633  
World Parkway Circle St. Louis, Mo. The sequence of  
S01623 and equivalent markers is taken from the 3' end of the  
following sequence: AAGACANNNN CGTTACATAA  
TCCTCACATA TAGTCATCCA ATCAGAACTG AATAG-  
GAAAA AAAAATACAC AATATTAATG AAATTTAATT  
TATCATCTGC ATGTTTGGAT AAGCGTCAAA  
GGTAAACCTA CTATTAGTAG CTTTCTTGTC TTTCCT-  
TCAA TTTGACGTGA TTTTAGTTTG AGACGTGCAT  
GTATAAAGTG GATCCAAACA CACTATTATG GTATG-  
CAGAG TGAAGTAAAA ACTTAAAAAT CAGAG-  
CAGCG ACCATTGCGT TCCCAGTCAC CATACTAGT  
GGGCTCAGG CTTGGGTC CACCAATCTC ACCT-  
GTTTCT TTGTTAATAC TGTCACCATC TTCGTGGTCT  
TCTTCGGGCT CATGGCTTTG TTTGTTCTCA TCATG-  
GAGAG ATTCTTGAGG TGGTGTCTGT GCTTGTTCCT  
TGAGNGGGTT TTCGTGTTGT GGCTGAGTTG  
AAGAGCAGNN GAGCCGTGTC ACTGTGTTTG AAA-  
CAAAATG GTAAACTGC TCGGATTGG TCGCGT-  
GANN NNCNNTGTTG GCTACACAAG CAGTGAGCG  
AGGGAANNNG GTGGTCATTG TTGTTTGTTA ATGAT-  
GTAAG GCAGATGATC AGAAANNAGA AAACCTCG-  
TAN CNNNACGAAC AAAACCTGA AATGGTTTAA

20

AGCTNNNCCT TGGATTTTGA TTCTTGTGTC  
TGCGCGTTNG NNTGC [SEQ ID NO:108]

Markers that map closer to the RAG locus are preferred  
over markers that map farther from the RAG locus for use in  
this invention. A more preferred set of markers includes:  
Satt150, Sat\_316, A351\_1, Mng339\_1, Ts, S01256,  
S02020, Satt567, Satt540, Bng222\_1, RGA\_2b, RGA5b,  
GM260, S04309, Satt435, R079\_1, A060-2, DOP\_H14,  
GM260, A131\_1, Sat\_244, S01623, Satt463, Sat\_253,  
S03544, Satt245, GM284, A946\_2, GM256, GMS057,  
Satt220, A584\_3, Satt626, OP\_N04, Satt323, Sat\_258,  
Satt702, L204\_4, GMS003b, Satt536, Sat\_003, OM11\_1,  
Sat\_148, Bng179\_2, K417\_3, Sat\_226, Satt175,  
K024\_1, A226\_1, GM230, ACCAGC315, Satt494, B157\_1,  
A715\_1, and Sct\_147.

A most preferred set of markers from which to choose at  
least one marker for use in this invention includes Satt435,  
Satt463, Satt245, S04309, S01623, and DOP\_H14.

The markers may be any type of mapped molecular marker  
or phenotypic trait known to the art, including restriction  
fragment length polymorphism (RFLP) markers, target  
region amplification polymorphism (TRAP) markers, ran-  
dom amplified polymorphic (RAPD) markers, single  
sequence repeat (SSR) markers, single nucleotide polymor-  
phism (SNP) markers, and isozyme markers.

In one embodiment of the invention, markers flanking the  
RAG locus are used in the marker-assisted selection pro-  
cesses of this invention. The genomic DNA of soybean ger-  
mplasm is preferably tested for the presence of at least two of  
the foregoing molecular markers, one on each side of the  
RAG locus. Most preferably, the two markers are Satt435 and  
Satt463. Markers that map close to Satt435 and Satt463 can  
also be used, provided they fall to either side of the RAG  
locus. Preferably, one of said at least two molecular markers  
is within at least about 10 to about 20 cM of Satt435 and  
another of said at least two molecular markers is within at  
least about 10 to about 20 cM of Satt463, and to ensure that  
the markers used flank the RAG locus, one of said at least two  
molecular markers within at least about 10 to about 20 cM of  
Satt435 should be farther than that distance from Satt463, and  
another of said at least two molecular markers within at least  
about 10 to about 20 cM of Satt463 should be farther than that  
distance from Satt435.

The method of this invention for reliably and predictably  
introgressing soybean *Aphis glycines* resistance into non-  
resistant soybean germplasm or less resistant soybean ger-  
mplasm comprises: providing a first soybean germplasm that  
has RAG-gene-derived resistance to *Aphis glycines*; provid-  
ing a second soybean germplasm that lacks RAG-gene-de-  
rived resistance to *Aphis glycines*; crossing the first soybean  
germplasm with the second soybean germplasm to provide  
progeny soybean germplasm; screening said progeny ger-  
mplasm to determine the presence of RAG-gene-derived resis-  
tance to *Aphis glycines*; and selecting progeny that tests posi-  
tive for the presence of RAG-gene-derived resistance to *Aphis  
glycines* as being soybean germplasm into which germplasm  
having RAG-gene-derived resistance to *Aphis glycines* has  
been introgressed.

Preferably, the screening and selection are performed by  
using marker-assisted selection using a marker on major link-  
age group M as described above.

The screening and selection may also be performed by  
exposing plants containing said progeny germplasm to aphids  
of the species *Aphis glycines* in a live aphid assay and select-  
ing those plants showing resistance to aphids as containing  
soybean germplasm into which germplasm having RAG-  
gene-derived resistance to *Aphis glycines* has been intro-

gressed. The live aphid assay may be any such assay known to the art, e.g., as described in Hill, C. B., et al., "Resistance to the soybean aphid in soybean germplasm" (2004) Crop Science 44:98-106, Hill, C. B., et al., "Resistance of *Glycine* species and various cultivated legumes to the soybean aphid (Homoptera: Aphididae)" (2004) J. Economic Entomology 97(3):1071-1077, or "Li, Y. et al., "Effect of three resistant soybean genotypes on the fecundity, mortality, and maturation of soybean aphid (Homoptera: Aphididae)" (2004) J. Economic Entomology 97(3):1106-1111, or as described in the Examples hereof. A preferred method includes placing aphid-infested plant parts on vegetative cotyledon (VC) stage plants and rating aphid population and plant damage weekly. As described herein, a 0-5 rating scale in which 0=no aphids present, 1=a few solitary and transient aphids present, 2=small scattered colonies, 3=dense colonies, 4=dense colonies with plant damage, and 5=dense colonies with severe plant damage, may be used.

The screening and selection may also be done directly by hybridizing nucleic acid from plants containing progeny germplasm to a nucleic acid fragment comprising a RAG gene, and selecting those plants having germplasm that hybridizes to the nucleic acid fragment as having RAG-gene-derived resistance to *Aphis glycines*.

The method of this invention for breeding a soybean plant homozygous for an *Aphis glycines* resistance gene that is a cultivar adapted for conferring, in hybrid combination with a suitable second inbred, resistance to *Aphis glycines*, comprises selecting a first donor parental line possessing the desired *Aphis glycines* resistance, said first donor parental line comprising an *Aphis glycines* resistance gene that is located on major linkage group M; crossing the first donor parental line with a second parental line that is high yielding in hybrid combination to produce a segregating plant population of genetically heterogenous plants; screening the plants of the segregating plant population for the gene; selecting plants from the population having the gene; and breeding by self-crossing the plants containing the gene until a line is obtained that is homozygous for the locus containing the gene and adapted for conferring, in hybrid combination with a suitable second inbred, resistance to *Aphis glycines*.

The screening and selection are preferably performed by using marker-assisted selection as described above, but may also be performed by live aphid bioassay as described above, selecting those plants showing resistance to aphids as containing soybean germplasm having a RAG gene. The screening and selection may also be done by hybridizing nucleic acid from plants containing said progeny germplasm to a nucleic acid fragment comprising a RAG gene and selecting those plants whose germplasm hybridizes to the nucleic acid fragment as having the gene.

As the parental line having soybean aphid resistance, any line known to the art or disclosed herein, as described above, may be used.

Also included in this invention are soybean plants produced by any of the foregoing methods:

Isolated nucleic acid fragments comprising a nucleic acid sequence coding for soybean resistance to *Aphis glycines*, are also included in this invention. The nucleic acid fragment comprises at least a portion of nucleic acid belonging to linkage group M, and further comprises nucleotide sequences falling between molecular markers Satt435 and Satt463. It is capable of hybridizing under stringent conditions to nucleic acid of a soybean cultivar resistant to *Aphis glycines*.

Vectors comprising such nucleic acid fragments, expression products of such vectors expressed in a host compatible therewith, antibodies to the expression product (both poly-

clonal and monoclonal), and antisense nucleic acid to the nucleic acid fragment are also included within this invention.

This invention also includes soybean plants having resistance to *Aphis glycines* comprising a RAG gene and produced by introgression of DNA containing the gene into a soybean germplasm lacking the gene in its genome, and progeny of said soybean plant.

Seed of a soybean germplasm produced by crossing a soybean variety having *Aphis glycines* resistance in its genome with a soybean variety lacking the RAG gene in its genome, and progeny thereof, is also included in this invention. Such seed, from BC3 or BC4 generations derived from crosses with aphid resistant Dowling×Loda F2 plants using as recurrent parents other soybean lines adapted to Illinois, is also included in this invention.

## EXAMPLES

### Example 1

#### Genetic Analysis Identifying the Aphid Resistance Gene Rag1 in Dowling

Crosses were made between the ancestral soybean cultivar Dowling and two susceptible cultivars, Loda and Williams 82. The parents, F<sub>1</sub>, and F<sub>2</sub> plants were tested in a choice test in the greenhouse using the methods described in Hill, C. B., et al., "Resistance to the soybean aphid in soybean germplasm" (2004) Crop Science 44:98-106. Three weeks after infestation, aphid colonization was visually rated using the following scale: 0=no aphids present, 1=few solitary and transient aphids present, 2=small scattered non-established colonies, 3=dense colonies, and 4=dense colonies with plant damage. Plants were considered resistant with a rating of 0, 1, or 2 and susceptible with a rating of 3 or 4. F<sub>1</sub> plants were all resistant to the soybean aphid, indicating that resistance was dominant over susceptibility.  $\chi^2$  analyses on the segregation of resistance phenotypes of F<sub>2</sub> plants from different F<sub>1</sub> plants (families) indicated that a single dominant gene, called Rag1 pending approval by the Soybean Genetics Committee, conditioned resistance (Tables 5 and 6). Evaluation of the segregation of aphid resistance in F<sub>2:3</sub> families confirmed the monogenic dominant inheritance of resistance from Dowling (Table 8).

TABLE 5

REACTIONS OF DOWLING × LODA  
F<sub>2</sub> PLANTS AND PARENTS 21 DAYS AFTER  
INFESTATION BY THE SOYBEAN APHID

Family	Number of plants	Observed		Expected (3:1)		$\chi^2$	P
		R <sup>1</sup>	S	R	S		
4021	19	14	5	14.25	4.75	0.018	0.89
4281	14	11	3	10.5	3.5	0.095	0.76
4301	16	13	3	12	4	0.333	0.56
4302	11	11	0	8.25	2.75	3.667	0.06
4303	11	9	2	8.25	2.75	0.273	0.6
4304	12	8	4	9	3	0.444	0.5
4306	15	8	7	11.25	3.75	3.756	0.05
4307	8	5	3	6	2	0.667	0.41
4308	6	2	4	4.5	1.5	5.556	0.02
4309	13	9	4	9.75	3.25	0.231	0.63
4310	10	8	2	7.5	2.5	0.133	0.72
4343	8	8	0	6	2	2.667	0.1

23

TABLE 5-continued

REACTIONS OF DOWLING × LODA F <sub>2</sub> PLANTS AND PARENTS 21 DAYS AFTER INFESTATION BY THE SOYBEAN APHID								
Family	plants	Number of		Expected (3:1)				P
		Observed		R	S	$\chi^2$		
4344	15	11	4	11.25	3.75	0.022	0.88	
4531	19	15	4	14.25	4.75	0.158	0.69	
Totals						18.023	0.32	
Pooled	177	132	45	132.75	44.25	0.017	0.9	
Heterogeneity						18.006	0.26	
Dowling	32	32	0					
Loda	32	12	31					

<sup>1</sup>R (resistant) = 0, 1, 2 aphid colonization rating; S (susceptible) = 3, 4 rating.<sup>2</sup>One Loda plant had an aphid colonization rating of 2.

TABLE 6

REACTIONS OF DOWLING × WILLIAMS 82 F <sub>2</sub> PLANTS AND PARENTS 21 DAYS AFTER INFESTATION BY THE SOYBEAN APHID								
Population	Number of plants	Number Observed		Expected (3:1)				P
		R <sup>1</sup>	S	R	S	$\chi^2$		
4041	179	135	44	134.25	44.75	0.002	0.89	
Dowling	19	19	0					
Williams 82	20	0	20					

<sup>1</sup>R (resistant) = 0, 1, 2 aphid colonization rating; S (susceptible) = 3, 4 rating.

## Example 2

## Genetic Analysis Identifying the Resistance Gene in Jackson

Crosses were made between the ancestral soybean cultivar Jackson and Loda. The parents, F<sub>1</sub>, and F<sub>2</sub> plants were tested in a choice test in the greenhouse using the methods described in Hill, C. B., et al., "Resistance to the soybean aphid in soybean germplasm" (2004) Crop Science 44:98-106. Three weeks after infestation, aphid colonization was visually rated using the following scale: 0=no aphids present, 1=few solitary and transient aphids present, 2=small scattered colonies, 3=dense colonies, and 4=dense colonies with plant damage. Plants were considered resistant with a rating of 0, 1, or 2 and susceptible with a rating of 3 or 4. F<sub>1</sub> plants were all resistant to the soybean aphid, indicating that resistance was dominant over susceptibility.  $\chi^2$  analyses on the segregation of resistance phenotypes of F<sub>2</sub> plants from different F<sub>1</sub> plants (families) indicated that a single dominant gene (Table 7) was present. Evaluation of the segregation of aphid resistance in F<sub>2,3</sub> families indicated that the segregation of families did not fit a monogenic dominant inheritance model (Table 8). The unexpected F<sub>2,3</sub> family segregation ratio may have been due to differential seed production between resistant and susceptible F<sub>2</sub> plants. Progeny of F<sub>2</sub> plants that produced at least 12 seeds were evaluated so that number plants tested exceeded the minimum required (10 plants) to have high confidence (95%) in detecting double recessive susceptible plants in segregating families with a monogenic dominant gene model. About 80% of the resistant F<sub>2</sub> plants produced at least 12 seeds, whereas about 17% of the susceptible F<sub>2</sub> plants produced 12 seeds or more.

24

TABLE 7

REACTIONS OF JACKSON × LODA F <sub>2</sub> PLANTS AND PARENTS 21 DAYS AFTER INFESTATION BY THE SOYBEAN APHID.								
Family	plants	Number of		Expected (3:1)				P
		Observed		R	S	$\chi^2$		
4123	38	28	10	28.5	9.5	0.04	0.85	
4124	40	28	12	30	10	0.53	0.47	
4201	39	29	10	29.25	9.75	0.01	0.93	
4202	38	30	8	28.5	9.5	0.32	0.57	
4203	40	29	11	30	10	0.13	0.72	
4204	39	26	13	29.25	9.75	1.44	0.23	
4211	30	21	9	22.5	7.5	0.4	0.53	
4212	40	38	2	30	10	8.53	0	
4213	40	25	15	30	10	3.33	0.07	
4214	40	28	12	30	10	0.53	0.47	
4215	40	25	15	30	10	3.33	0.07	
4216	40	28	12	30	10	0.53	0.47	
4432	19	9	10	14.25	4.75	7.74	0.01	
Totals						26.87	0.01	
Pooled	483	344	139	362.25	120.75	3.68	0.06	
Heterogeneity						23.2	0.02	
Jackson	24	24	0					
Loda	51	0	51					

<sup>1</sup>R (resistant) = 0, 1, 2 aphid colonization rating; S (susceptible) = 3, 4 rating

TABLE 8

REACTIONS OF DOWLING × LODA, DOWLING × WILLIAMS 82, AND JACKSON × LODA F <sub>2,3</sub> FAMILIES AND 21 DAYS AFTER INFESTATION BY THE SOYBEAN APHID									
F <sub>2,3</sub> population	Number of families <sup>1</sup>	Number Observed		Expected (1:2:1)				$\chi^2$	P
		R <sup>2</sup>	H	S	R	H	S		
Dowling × Loda	146	31	73	42	36.5	73	36.5	1.65	0.44
Dowling × Williams 82	128	35	63	30	32	64	32	0.42	0.81
Jackson × Loda	206	86	96	24	51.5	103	51.5	38.27	0

<sup>1</sup>12 seeds of each F<sub>2</sub> plant were sown.<sup>2</sup>R = all plants in an F<sub>2,3</sub> family were resistant, H = plants in a family segregated for resistance, S = all plants in a family were susceptible.

## Example 3

## Molecular Markers Linked to Rag1

A soybean F<sub>2</sub> population developed from a cross between Dowling×Loda was used for mapping the location of Rag1. A total of 90 F<sub>2</sub> individuals and the two parents were included in the mapping work. The phenotypic data (aphid colonization on F<sub>2</sub> plants) was scored as described above in the genetic analysis.

For genotypic data, DNA was isolated from individual plants and polymerase chain reaction (PCR) was carried out using simple sequence repeat (SSR) markers developed by Dr. Perry Cregan, USDA-ARS (See Table 2). The PCR products were evaluated on gels as previously described in Wang, D. J. et al., "A low-cost, high-throughput polyacrylamide gel electrophoresis system for genotyping with micro satellite DNA markers," (2003) Crop Science 43:1828-1832. Initial screening was done using the parents and two bulked DNA

samples to identify polymorphic simple sequence repeat (SSR) markers. Each bulk consisted of pooled DNA samples from five susceptible F<sub>2</sub> individuals. A total of about 342 SSR markers were screened against the bulks to identify polymorphic markers potentially associated with aphid resistance. Markers showing strong association with Rag1 were further screened using the entire mapping population to determine linkage relationships and map locations. Joinmap 3.0 was used to create a genetic map. As shown in FIG. 1, Rag1 mapped to Linkage Group M where it is flanked by the SSR markers Satt435 and Satt463 that are 3 cM and 6 cM from the Rag1 locus, respectively.

#### Example 4

##### Molecular Markers Linked to Rag1 in Jackson

A soybean F<sub>2</sub> population developed from a cross between Jackson and Loda was used for mapping the location of the resistance gene to *Aphis glycines*. A total of 92 F<sub>2</sub> individuals and the two parents were included in the mapping work.

The phenotypic data (aphid colonization on F<sub>2</sub> plants) was scored as described above in the genetic analysis.

For genotypic data, DNA isolation, PCR, and gel electrophoresis were done as described in Wang, D. J. et al., "A low-cost, high-throughput polyacrylamide gel electrophoresis system for genotyping with micro satellite DNA markers," (2003) Crop Science 43:1828-1832). Three SSR markers, Satt435, Satt463, and Satt245, which are mapped 3 cM, 6 cM, and 13 cM from Rag1 in Dowling (FIG. 2), respectively, showed polymorphism between Jackson and Loda and are associated with aphid resistance in Jackson based on 14 F<sub>2</sub> individuals. These three markers were further screened using the entire mapping population to determine linkage relationships and map locations.

Joinmap 3.0 was used to create a genetic map. The *Aphis glycines* resistance gene locus mapped to Linkage Group M where the SSR marker Satt435 is 9 cM away from the gene locus.

#### Example 5

##### Location of Rag1 in the Soybean Genetic Map

F<sub>2:3</sub> populations from the cross between Dowling and the two susceptible soybean cultivars, Loda and Williams 82, were used to map Rag1 in Dowling using linked SSR markers Satt150, Satt540, Satt435, Satt463, Satt245, Satt220 and Satt323. See Tables 2 and 3.

One hundred and forty nine F<sub>2</sub> plants and their F<sub>2:3</sub> families from Dowling×Loda were used for initial marker screening and initial mapping of Rag1 in Dowling. One hundred and twenty one F<sub>2:3</sub> families from Dowling×Williams 82 were used to confirm the Rag1 map location and to construct an integrated map for Rag1 in Dowling.

In the integrated map from Dowling×Loda and Dowling×Williams 82 populations, Rag1 was mapped to soybean linkage group M flanked by the SSR markers Satt435 and Satt463 4.0 cM and 8.0 cM from Rag1, respectively (Tables 9 and 10; FIGS. 1 and 2).

##### Plant Materials

Three F<sub>2:3</sub> populations from the crosses "Dowling" (PI 548663)×"Loda" (PI 614088), Dowling×"Williams 82" (PI 518671) and "Jackson" (PI 548657)×Loda, and one F<sub>2</sub> population of Dowling×Palmetto (PI 548480), were used in this study based on crosses made by Curt Hill. Dowling, Jackson and Palmetto are aphid resistant while Loda and Williams 82

are aphid susceptible. Palmetto was suggested as the origin of the resistance in Jackson because it is the only known resistant ancestor of Jackson (Hill, C. B., et al., "Resistance to the soybean aphid in soybean germplasm" (2004) Crop Science 44:98-106). One hundred and forty nine F<sub>2</sub> plants and their F<sub>2:3</sub> families from Dowling×Loda were used for initial marker screening and initial mapping of Rag1 in Dowling. One hundred and twenty one F<sub>2:3</sub> families from Dowling×Williams 82 were used to confirm the Rag1 map location and to construct an integrated map for Rag1 in Dowling. One hundred and forty F<sub>2</sub> plants and their F<sub>2:3</sub> families from Jackson×Loda were used to map the gene in Jackson. Sixty-five F<sub>2</sub> plants from Dowling×Palmetto were used to test allelism indirectly between Rag1 in Dowling and Jackson. Dowling×Jackson crosses were not made because there are no known polymorphic markers known that could be used to distinguish F1 hybrids from selfs in crosses.

##### Aphid Clone

The aphid clone was collected from Urbana, Ill. by Dr. Les Domier (USDA-ARS and Department of Crop Sciences, University of Illinois, Urbana, Ill. 61801) and reared on the seedlings of soybean cultivar Williams 82 in a plant growth chamber at 22° C. under continuous 200 μmol m<sup>-2</sup> s<sup>-1</sup> PAR irradiation.

##### Soybean Aphid Resistance Phenotyping

The parents, F<sub>2</sub> plants, and susceptible checks were screened for aphid resistance under semi-controlled conditions (22-25° C. under continuous 24-h illumination (160-200 μmol m<sup>-2</sup> s<sup>-1</sup>) in the greenhouse. In a randomized complete block design, seeds were grown in plastic multi-pot inserts within plastic trays without holes. One week later, soybean aphids were transferred from the infested Williams 82 cut stems and leaves to the young test seedlings. Each individual plant was evaluated for aphid score twice at 14 days and 21 days after infestation. Aphid score was rated as index based on aphid population density and plant damage: 0-4, where 0=no aphids observed, 1=few number of aphids scattered on the plant, 2=limited colonization of aphids observed, 3=high aphid density and colonization, 4=high aphid density and colonization plus leaf distortion and plant stunting. After the 21-day rating, insecticide (imidacloprid) was applied. After one week, leaf tissue from the F<sub>2</sub> plants was sampled for DNA extraction. All F<sub>2</sub> plants were transplanted to 5-inch diameter plastic pots and were grown in the greenhouse under a 12 h photoperiod to produce F<sub>3</sub> seeds. In the progeny test, 10-12 F<sub>3</sub> seeds per F<sub>2</sub> family were evaluated for aphid resistance in a randomized complete block design with three replicates (four F<sub>3</sub> plants per F<sub>2</sub> family per replicate). F<sub>2</sub> genotypes (homozygous resistant, heterozygote, or homozygous susceptible) were inferred from the segregation of the F<sub>3</sub> plants.

##### DNA Isolation, PCR Reaction and Gel Electrophoresis

Young trifoliolate leaves were harvested from the new growth of each individual plant after the aphids were killed. Soybean DNA was extracted from either an individual F<sub>2</sub> plant or pooled 10-12 F<sub>3</sub> plants, by using either the CTAB method (Keim, P. and Shoemaker, R. C., "Construction of a random recombinant DNA library that is primarily single copy sequence" (1988) Soybean Genet. Newslet. 15:147-148), or DNA quick extraction method (Bell-Johnson, B. et al., "Biotechnology approaches to improving resistance to SCN and SDS: methods for high throughput marker assisted selection" (1998) Soybean Genet. Newslet. 25:115-117).

The PCR amplification was performed in a PTC-220 Thermalcycler manufactured by MJ Research (Waltham, Mass.). PCR reactions were done in 15 μl volumes with 50-250 ng of template DNA, 2 μM primer, 30 mM MgCl<sub>2</sub>, 3 mM each

dNTP, 2.5 unit of Taq polymerase, and 1×PCR buffer. The PCR was performed with an initial denaturing at 94° C. for 4 min, followed by 34 cycles of 25 s of denaturing at 94° C., 25 s of annealing at 47° C., and 25 s of extension at 68° C., with a final 7-min extension at 72° C.

The gel electrophoresis was done using non-denaturing polyacrylamide gels as described before (Wang, D. J. et al., "A low-cost, high-throughput polyacrylamide gel electrophoresis system for genotyping with micro satellite DNA markers," (2003) Crop Science 43:1828-1832). After electrophoresis the gels were photographed and the polymorphic bands were scored as described below.

#### SSR Marker Screening and Bulk Segregant Analysis

Soybean simple sequence repeat (SSR) markers developed by Dr. Cregan (Cregan, RB., et al., "An Integrated Genetic Linkage Map of the Soybean Genome" (1999) Crop Science 39:1464-1490) were used in this study. Bulk segregant analysis (Michelmore, R.W., et al., "Identification of markers linked to disease resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions using segregating populations" (1991) Proc. Natl. Acad. Sci. (USA) 88:9828-9832 in Dowling×Loda F<sub>2</sub> population was used to screen for aphid-resistance associated markers. Since at the time of screening, only F<sub>2</sub> individuals were available, the resistant F<sub>2</sub> plants could be either heterozygous or homozygous, therefore susceptible bulks were used to look for linkage. Two pools of DNA from five susceptible F<sub>2</sub> individuals each, bulk A and bulk B were prepared. DNA from resistant parent Dowling and susceptible parent Loda, along with the DNA pools bulk A and B were used to identify polymorphic SSR markers with potential association with aphid resistance. 342 SSR markers were tested for polymorphism between two parents Dowling and Loda, and the segregation patterns for the aphid resistance versus susceptibility. The polymorphic markers with putative linkage with Rag1 were first identified by contrasting bulk segregant analysis, and then were further screened in the whole Dowling×Loda mapping population.

The SSR markers that were determined to map close to Rag1 in the Dowling×Loda mapping population were used to test the polymorphism between Dowling and Williams 82 and between Jackson and Loda. The polymorphic markers were then further screened in the entire population of Dowling×Williams 82 and Jackson×Loda. The data from Dowling×Loda and Dowling×Williams 82 was integrated together to map Rag1 in Dowling, and the data from Jackson×Loda was used to map the gene in Jackson.

The two SSR markers that flanked the Rag1 gene in Dowling and Jackson were found to be monomorphic in Palmetto also, the parent of Jackson.

#### Genetic Mapping

Joinmap 3.0 (Van Ooijen, J. W. and Voorrips, V. E., Joinmap 3.0, Software for the calculation of genetic linkage maps. Plant Research International, Wageningen, the Netherlands (2001)) was used for linkage analysis to create a genetic map using the Kosambi mapping function. A LOD of 3.0 was used as the threshold to group markers into linkage groups. Chi square ( $\chi^2$ ) test at P=0.05 was used to verify the segregation ratio of each locus in the F<sub>2</sub> population. The genotypes of the SSR markers were scored as either co-dominant (A=RR, H=Rr, B=rr) or dominant (D=A+H, B or A, C=B+H). Genotypes of the F<sub>2</sub> aphid population were scored as co-dominant (A, H, B) after confirmation with the F<sub>3</sub> progeny test. Those that had less than 10 F<sub>3</sub> seeds available from the F<sub>2</sub> plant were scored as dominant (D, B). All of the SSR markers and aphid resistance genes were set to the cor-

responding  $\chi^2$ -test classification as described in Joinmap 3.0, 1:2:1 (A: H: B) or 3:1 (A+H+D: B=3:1) segregation ratio.

TABLE 9

Locus	a	h	b	c	d	—	$\chi^2$	Classes
Rag1	26	72	44	1	5	1	1.9	[a + h + d:b]
Satt150	25	61	35	1	8	19	0.3	[a + h + d:b]
Satt220	22	24	31	4	33	35	0.6	[a + h + d:b]
Satt245	24	76	43	0	3	3	1.5	[a + h + d:b]
Satt435	17	68	46	1	13	4	3.7	[a + h + d:b]
Satt463	35	57	39	0	11	7	0.5	[a + h + d:b]
Satt540	19	81	38	1	6	4	0.1	[a + h + d:b]

TABLE 10

Locus	a	h	b	c	d	—	$\chi^2$	Classes
Rag1	25	69	27	0	0	0	2.5	[a:h:b]
Satt150	31	42	25	2	13	8	0.4	[a + h + d:b]
Satt245	22	67	22	3	5	2	2.3	[a + h + d:b]
Satt323	27	59	28	4	2	1	0.1	[a + h + d:b]
Satt435	21	62	27	1	10	0	0.4	[a + h + d:b]
Satt463	31	55	22	3	10	0	2.5	[a + h + d:b]
Satt540	19	63	25	8	5	1	0.4	[a + h + d:b]

#### Example 6

##### Location of *Aphis glycines* Resistance Gene in the Soybean Genetic Map

One hundred and forty F<sub>2</sub> plants and their F<sub>2:3</sub> families from Jackson×Loda were used to map the RAG gene in Jackson using linked SSR markers (Tables 2 and 3).

The RAG gene was mapped to linkage group M flanked by markers Satt435 and Satt463 1.9 cM and 7.7 cM, respectively (Table 11; FIGS. 1 and 2).

TABLE 11

Locus	a	h	b	c	d	—	$\chi^2$	Classes
RAG	40	58	26	0	16	0	3.1	[a + h + d:b]
Satt150	29	53	20	4	14	20	3.7	[a + h + d:b]
Satt220	46	56	14	10	7	7	12.2*	[a + h + d:b]
Satt245	47	42	24	0	9	18	1.9	[a + h + d:b]
Satt435	43	65	26	0	6	0	3.1	[a + h + d:b]
Satt463	40	57	32	0	8	3	0.2	[a + h + d:b]
Satt540	38	68	24	7	2	1	3.3	[a + h + d:b]

Although the foregoing invention has been described in detail for purposes of clarity and understanding, it will be clear to those skilled in the art that equivalent cultivars, markers, and methods may be practiced within the scope of the claims hereof.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 108

<210> SEQ ID NO 1

<211> LENGTH: 894

<212> TYPE: DNA

<213> ORGANISM: Glycine max

<400> SEQUENCE: 1

```

aagcttctat caagtggtaa tcagagcaca agatcttcaa gtaggtgata cttaaacctc      60
cattaatttt ttgctttacc ttctcttcta ttgttggttc ttcatttttc tccatgtatc      120
tcctcacatg tcttggtgta aatgttttta acatgattct ttagagtttc caccgattaa      180
acttgctata gaagctagat ttgattttct atgggttcaaa tttcttggtc ttgttcttga      240
tccatgaatt gtgttgagtt taggttcctt tgagttttgt cttgttatct tttgtggctg      300
aaacctaaac cataaaattc ttacaaaaat attaaagtag aggaaaacct caaaaatcta      360
gagtgaactg ttcacctatt atagttttgt catagaagtc atgtctagtc atgaaacttg      420
tcacataaga tttcttatgt tgtgtgaat tttattttct tgtttctttg tctaaactcat      480
ttgttcataa gtgtatgaag ttatttttagc ctattatttg attggagtca aatctttcat      540
gttaattagt ccttaacatg ttcatgcaaa attcttagag agtctttgat tgtgaacctt      600
ttcttgaact tttaggtttc cttatgattg tgtctattgt gaatttaagt tttggtgatt      660
gaattgctgg ttgaaatggt gatcctaagt gaattatgaa ctctaaaac tgtggtaaac      720
aatcctagtg agttcaacat acataggaag gttgaaagta agcccaaggc aatcaatata      780
gcatgcttaa aaaaaaaaac gctgggtgctg gcagcttgga catacaaact tgtaaaaatt      840
actgaaaatt ggttactctg aattttgaac tgaattttta cttaatttgc taga          894

```

<210> SEQ ID NO 2

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 2

```

gcgggtagcc atattcatat attgctg      27

```

<210> SEQ ID NO 3

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 3

```

gcgaaggcctt ataaggagat acgattta      28

```

<210> SEQ ID NO 4

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 4

```

tcatccgcca ttgatttt      18

```

<210> SEQ ID NO 5

-continued

---

<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 5  
  
gcccggaaca tacaaaat 18  
  
<210> SEQ ID NO 6  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 6  
  
gcgtaggcat cgtcaatat ttt 23  
  
<210> SEQ ID NO 7  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 7  
  
gcgttagcga gtggatcaag atca 24  
  
<210> SEQ ID NO 8  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 8  
  
tacctttctt gtgagtcgta 20  
  
<210> SEQ ID NO 9  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 9  
  
tattgagatg gatattgtag atc 23  
  
<210> SEQ ID NO 10  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 10  
  
gtcatgactc atgagtcacg taat 24  
  
<210> SEQ ID NO 11  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 11



-continued

---

cccaagaccc ccatttttat gtct 24

<210> SEQ ID NO 12  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 12

gcgcgcattt ttaagttaa tgttct 26

<210> SEQ ID NO 13  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 13

gcgcgagtta gcgaattatt tgtc 24

<210> SEQ ID NO 14  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 14

gcgttgatac tttcctaaga caat 24

<210> SEQ ID NO 15  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 15

gggagagaag gcaatctaa 19

<210> SEQ ID NO 16  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 16

aagcttgagg ttattcgaaa atgac 25

<210> SEQ ID NO 17  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 17

tgccatcagg ttgtgtaagt gt 22

<210> SEQ ID NO 18  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 18

gcgcaacgctc taaagcaciaa ggatt 25

<210> SEQ ID NO 19

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 19

gcgcgactac gttacagtgc caa 23

<210> SEQ ID NO 20

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 20

ggctaaccgc ctctatgt 18

<210> SEQ ID NO 21

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 21

ggcccatgca cctgctact 19

<210> SEQ ID NO 22

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 22

ctggcgaatc aagctttgta ac 22

<210> SEQ ID NO 23

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 23

ccgtgattgc gaagaggata tt 22

<210> SEQ ID NO 24

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 24

gcggtgaaac ggctctcttt gatagtga 28

<210> SEQ ID NO 25

-continued

---

<211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 25

gcgttgatt aattaattaa attatttt

28

<210> SEQ ID NO 26  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 26

gcgtcaaccg gtgaaaaaac cta

23

<210> SEQ ID NO 27  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 27

gcgtggctgg cagtagtcta tatca

25

<210> SEQ ID NO 28  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 28

ttgatctca tattcaaaact ttcaag

26

<210> SEQ ID NO 29  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 29

ctgcaaattt gatgcacatg tgtcta

26

<210> SEQ ID NO 30  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 30

gcgattgggtt ggggtgttaa tttaagat

29

<210> SEQ ID NO 31  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 31

-continued

---

gcgtgttgat ggtataaaga tcgctactct	30
 <210> SEQ ID NO 32 <211> LENGTH: 22 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer  <400> SEQUENCE: 32  aacgggagta ggacatttta tt	
	22
 <210> SEQ ID NO 33 <211> LENGTH: 28 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer  <400> SEQUENCE: 33  gcgcctcctg aatttcaaag aatgaaga	
	28
 <210> SEQ ID NO 34 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer  <400> SEQUENCE: 34  gaggaggatc ccaaggtaat aat	
	23
 <210> SEQ ID NO 35 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer  <400> SEQUENCE: 35  gcgcatggag aaaagaagag	
	20
 <210> SEQ ID NO 36 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer  <400> SEQUENCE: 36  gcggatggag acggggggga cggacga	
	27
 <210> SEQ ID NO 37 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer  <400> SEQUENCE: 37  gcgcatagct aattttatat caattat	
	27
 <210> SEQ ID NO 38 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE:	

-continued

---

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 38

gcggctcgtcc tatctaataga agag 24

<210> SEQ ID NO 39

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 39

tgtgcggttta aattgcagct aaat 24

<210> SEQ ID NO 40

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 40

gcgcaataga taatcgaaaa acatacaaga 30

<210> SEQ ID NO 41

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 41

gcgggggaaag tgaaaacaag atcaaata 28

<210> SEQ ID NO 42

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 42

gcgggggttct gtggcttcaa c 21

<210> SEQ ID NO 43

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 43

gcgcattgga ataacgtcaa a 21

<210> SEQ ID NO 44

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 44

gcgccacaga aattcctttt tcta 24

<210> SEQ ID NO 45

-continued

---

<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 45  
  
gcgccataag gtggttacca aaaga 25  
  
<210> SEQ ID NO 46  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 46  
  
tgatttttgg tgtagaactc 20  
  
<210> SEQ ID NO 47  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 47  
  
caaattgggt agcttactcc a 21  
  
<210> SEQ ID NO 48  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 48  
  
gcggagtttc ccctaattag at 22  
  
<210> SEQ ID NO 49  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 49  
  
gcgcaagcta gcttcaccca aaacta 26  
  
<210> SEQ ID NO 50  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 50  
  
gcggaaaccc acctatatgt gatcaaatg 29  
  
<210> SEQ ID NO 51  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 51

-continued

---

gcgcaattcc agatgaaaca gaagaaggat 30

<210> SEQ ID NO 52  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 52

gacctogetc tctgtttctc at 22

<210> SEQ ID NO 53  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 53

ggtgaccacc cctattcctt at 22

<210> SEQ ID NO 54  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 54

ggccggttct cattacaggt ctct 24

<210> SEQ ID NO 55  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 55

ggatttccat cttgaatttt atta 24

<210> SEQ ID NO 56  
 <211> LENGTH: 17  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 56

tctcgactca cgactca 17

<210> SEQ ID NO 57  
 <211> LENGTH: 17  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 57

ccaaggtctc tcagagg 17

<210> SEQ ID NO 58  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 58

gcgcggaaaa ttattttact ttttcaat 28

<210> SEQ ID NO 59

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 59

gcgcacggat tgagagaaag cagaaaga 28

<210> SEQ ID NO 60

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 60

caacgaccaa ctgacgagac ct 22

<210> SEQ ID NO 61

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 61

gggaattcaa catgtgatgg tttt 24

<210> SEQ ID NO 62

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 62

gcgacagact gcaagaattg atgtaaactc 30

<210> SEQ ID NO 63

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 63

gcgggaaggt aggtaaagaa aattcaaag a 31

<210> SEQ ID NO 64

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 64

gaagaccaa acttatttca gatc 24

<210> SEQ ID NO 65



-continued

---

<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 65  
  
attttaagca ccagcaaaga ct 22  
  
<210> SEQ ID NO 66  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 66  
  
gcgggatatc gtgagcatag ttttac 26  
  
<210> SEQ ID NO 67  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 67  
  
gcggcctgaa tatttttaggt ttagagtt 28  
  
<210> SEQ ID NO 68  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 68  
  
gcgtattttg ggggattttg aaca 24  
  
<210> SEQ ID NO 69  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 69  
  
gcgtttctct tcttattctt tctct 25  
  
<210> SEQ ID NO 70  
<211> LENGTH: 17  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 70  
  
gcgaaatgcc gaaagag 17  
  
<210> SEQ ID NO 71  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer  
  
<400> SEQUENCE: 71

-continued

---

gcggggataa gaaaaacaat 20

<210> SEQ ID NO 72  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 72

gcgcttaagg acacggatgt aac 23

<210> SEQ ID NO 73  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 73

gcgtctcttt cgattgttct attag 25

<210> SEQ ID NO 74  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 74

gcgtaccctt atatggatgt ttcttcct 28

<210> SEQ ID NO 75  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 75

gcgtatgcag caaacaaaaa atatataat 29

<210> SEQ ID NO 76  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 76

gcgttttcct aatgaagatt t 21

<210> SEQ ID NO 77  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 77

gcgtgtaata gtgatggatg taa 23

<210> SEQ ID NO 78  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial sequence  
<220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 78

gcgtgcttta aatgattgat tga 23

<210> SEQ ID NO 79

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 79

gcgtgcgaac ataactaata cat 23

<210> SEQ ID NO 80

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 80

gaatatcacg cgagaatttt ac 22

<210> SEQ ID NO 81

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 81

tatatgcgaa ccctcttaca at 22

<210> SEQ ID NO 82

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 82

gacaaatgta aaaagtgaca gatagaatgt 30

<210> SEQ ID NO 83

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 83

gtgtggtggt ggtacagttt tatactaa 28

<210> SEQ ID NO 84

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 84

cgccagctag ctagtctcat 20

<210> SEQ ID NO 85

-continued

---

<211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 85

aatttgctcc agtgttttaa gttt 24

<210> SEQ ID NO 86  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 86

gcggtgatat taccctaaaa aaatgaa 27

<210> SEQ ID NO 87  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 87

gcgctagttt ctagtggaat gatgagt 27

<210> SEQ ID NO 88  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 88

gcgaaaaacg tcaggtcaat gactgaaa 28

<210> SEQ ID NO 89  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 89

gcggggctta gatataaaaa aaaagatg 28

<210> SEQ ID NO 90  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 90

ggagggagga aagtgttgtg g 21

<210> SEQ ID NO 91  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 91

-continued

---

 gcgcacgtctt ttcataagtt t 21

<210> SEQ ID NO 92  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic priemr

<400> SEQUENCE: 92

gtgcgacgtc atgccttact caat 24

<210> SEQ ID NO 93  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 93

gcgctccgta cacttaaaaa agaa 24

<210> SEQ ID NO 94  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 94

gcggaaaccc atctagaata tgaaaaaca 29

<210> SEQ ID NO 95  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 95

gcgttcttct cgagggtgaga tacaatc 27

<210> SEQ ID NO 96  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 96

gcgttaaggt tggcaggggtg gaagtg 26

<210> SEQ ID NO 97  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 97

gcgcagcttt atacaaaaat caacaa 26

<210> SEQ ID NO 98  
 <211> LENGTH: 18  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial sequence  
 <220> FEATURE:



-continued

---

```

<211> LENGTH: 797
<212> TYPE: DNA
<213> ORGANISM: Glycine max
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(797)
<223> OTHER INFORMATION: N is A, T, G or C.

<400> SEQUENCE: 105
tattcattata ttgcaggcta cnaaaatttc cagtnntaat acagtataat taagcagagt      60
gtggtatctca caaaatctca atccaaacac ataattacaa aactctagaa cagcagaaca      120
catatagcat ttgatttgaa gtattcattc actaattgat tagccttaga aattcaaattg      180
atataatctg accactcaga gataaaggaa gtatgggtcca tggactcccc aggaacatcc      240
tcgtgcttag agggccttctc ctccccacca accaacctgg ctggggtccc aacagctggt      300
gtctgtgggtg gcacatcgat taaaaccacc gagccagcac aacctttgca ccttccccga      360
tcttaaatatt cccagaatg gtagcaccgg caccaataag caccatcc ccaatcttgg      420
gatgccggtc cccaccaact tgccagtcac acccagcgta acgtgggtgca ggatcgacac      480
attgttccga tcaactgcgt cccccccacc accaccccggt tggcatgggc gaacagatcc      540
ccttcccgat cctgcgcgca ggggtgaatgt ccaccgcgaa cacatcgca tgcgagagt      600
cagtgcnaaa gccaatggct gccgcgattg ncnccaaca gatgcgccac acggtgcgcc      660
tgcaaaatca caatcacaca caactaatcc taagattcaa taatcaaaaa agagtnnact      720
nnncatacac tgtcatcnch nntatagtca tgtttcatnn naatctngnn nnacaatgca      780
tataaattaa actcaat                                         797

<210> SEQ ID NO 106
<211> LENGTH: 701
<212> TYPE: DNA
<213> ORGANISM: Glycine max
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(701)
<223> OTHER INFORMATION: N is A, T, G or C.

<400> SEQUENCE: 106
aaagnnaaca tttttgttta tatgacnna acaaactgca aagaaaaatt gttaaaaacc      60
agaagcaatt taggtgatca caaataccac atgcttacac cttccagtga caagtacagt      120
atgttgtggc accagccgtt tcagttgatg caaacttgct tcgtgccaaa attctaaca      180
cacaactacc taagctatca aacaagagaa gcccttttgt cctttggtcg acctatcaaa      240
ggtcacgaga tcacactagt cctacccttt taagaaaacc tactatcaac agtcatatgt      300
atctcatgaa aagcacataa aaacatgtca ctttgctctc tcaccatctc cactgttatg      360
agcagccgag gagctgcctt ggccgtctcc accagctggt ccagcctcag aggcattctg      420
cttgcttcca ccacgtgcat cgtttgacc agtagccgaa ggtggaccac cgctgtttcc      480
cctcaagag cagcctcact gtgcattgga tgcattgcat tatttatatc tccaggtcta      540
agtcacattt gaccttggat ggccgtgctg tgtagctgct gttgtgttcc ctgcatttga      600
tgtggattgc caaattgcaa tggcattttc tgggggaaca nnccttgcgt ctgctgnnnn      660
attgctgctg cagcnnntg nnnatnnnnn atatannnnn c                                         701

<210> SEQ ID NO 107
<211> LENGTH: 799
<212> TYPE: DNA
<213> ORGANISM: Glycine max
<220> FEATURE:

```

-continued

&lt;221&gt; NAME/KEY: misc\_feature

&lt;222&gt; LOCATION: (1) .. (799)

&lt;223&gt; OTHER INFORMATION: N is A, T, G or C.

&lt;400&gt; SEQUENCE: 107

tatcattata ttgcaggeta cnaaaatttc cagtnntaat acagtataat taagcagagt	60
gtggtatcta caaaatctca atccaaacac ataattacaa aactctagaa cagcagaaca	120
catatagcat ttgatttgaa gtattcattc actaattgat tagccttaga aattcaaagt	180
atataatctg accactcaga gataaaggaa gtatgggtcca tggactcccc aggaacatcc	240
tcgtgcttag agggcttctc cttcccacca accaacctgg ctgggttccc aacagctgtt	300
gtctgtggtg gcacatcgat taaaaccacc gagccagcac caacctttgc accttccccg	360
atcttaatat tccccagaat ggtagcaccg gcaccaataa gcaccccatc cccaatcttg	420
ggatgccggt ccccaaccaac cttgccagtc caccagcgt aacgtggtgc aggatcgaca	480
cattgttccc gatcactgcc gtctccccca ccaccacccc ggtggcatgg tcgaacagaa	540
tcccccttcg atcctcgccg cagggtgaat gtccaccgag aacacatcag cgatgcgaag	600
tgcagtgcna aagccaatgg ctgcccgat tgnccgcaca acagatggcc acacggtgcg	660
cctgcaaaat cacaatcaca cacaactaat cctaagattc aataatcaaa aaagagtnna	720
ctnnncatag actgtcatcn cnntatagt catgtttcat nnnaatctng nnnnacaatg	780
catataaatt aaactcaat	799

&lt;210&gt; SEQ ID NO 108

&lt;211&gt; LENGTH: 702

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Glycine max

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: misc\_feature

&lt;222&gt; LOCATION: (1) .. (702)

&lt;223&gt; OTHER INFORMATION: N is A, T, G or C.

&lt;400&gt; SEQUENCE: 108

aagacannnn cgttacataa tcctcacata tagtcatcca atcagaactg aataggaaaa	60
aaaaatacac aatattaatg aaatttaatt tatcatctgc atgtttggat aagcgtcaaa	120
ggtaaaccta ctattagtag ctttcttgtc tttccttcaa ttgacgtga ttttagtttg	180
agacgtgcat gtataaagtg gatccaaaca cactattatg gtatgcagag tgaagtaaaa	240
acttaaaaat cagagcagcg accattgctg tcccagtcac catacctagt gggctcaggc	300
ccttgggtcc accaatctca cctgtttctt tgtaataact gtcaccatct tcgtggtctt	360
cttcgggctc atggctttgt ttgttctcat catggagaga ttcttgaggt ggtgtctgtg	420
cttggtccct gagnggggtt tcgtgtgtgt gctgagttga agagcagnng agcgtgtca	480
ctgtgttgga aacaaaatgg taaactgtc ggatttgggt cggtgannnn cnntgttggc	540
tacacaagca gtgagcgagg gaannnggtg gtcattgttg tttgttaatg atgtaaggca	600
gatgatcaga aannagaaaa ctcgatcnnn nacgaacaaa accctgaaat ggtttaaagc	660
tnnncccttg attttgatc ttgttgctgc gcgttngnnt gc	702



The invention claimed is:

1. A method for reliably and predictably introgressing soybean *Aphis glycines* resistance into non-resistant soybean germplasm or less resistant soybean germplasm comprising:

- a. providing a first soybean germplasm that has resistance to *Aphis glycines*;
- b. providing a second soybean germplasm that lacks resistance to *Aphis glycines*;
- c. crossing the first soybean germplasm with the second soybean germplasm to provide progeny soybean germplasm;
- d. analyzing said progeny germplasm to determine the presence of resistance to *Aphis glycines* by identifying the presence or absence of a gene coding for resistance to *Aphis glycines* in said progeny germplasm comprising analyzing said germplasm by marker-assisted selection (MAS) to:

(1) detect a resistance to *Aphis glycines* (RAG) locus that maps to soybean linkage group M of said soybean germplasm,

- (a) wherein said RAG locus is flanked on opposite sides by SSR markers Satt435 and Satt463, which show allelic polymorphism between *Aphis glycines*-resistant and *Aphis glycines*-susceptible soybean genotypes and are linked to the RAG locus, and
- (b) wherein the RAG locus comprises allelic DNA sequences that control resistance to *Aphis glycines*; and

(2) determine the presence or absence of an allelic form of DNA linked to the gene coding for resistance to *Aphis glycines* in said germplasm; wherein the presence or absence of said allelic form of DNA linked to said gene is determined by a method comprising:

- (a) comparing a first PCR-amplified polymorphic marker fragment of said soybean germplasm to a second PCR-amplified polymorphic marker fragment of soybean germplasm from a plant having *Aphis glycines* resistance conferred by said Rag1 gene;
- i. wherein said second fragment is made using the same marker used to make said first fragment, and
- ii. wherein said second fragment has a size substantially the same as that of a PCR-amplified polymorphic marker fragment of germplasm of *Aphis glycines*-resistant soybean varieties Dowling and Jackson made using the same marker used to make said first and second fragments; and

- (b) determining that said gene coding for RAG resistance is present in said soybean germplasm when said first fragment is substantially the same size as said second fragment, and determining that said gene is not present in said germplasm when said first fragment is not substantially the same size as said second fragment, and

e. selecting progeny that tests positive for the presence of resistance to *Aphis glycines* as being soybean germplasm into which germplasm having resistance to *Aphis glycines* has been introgressed.

2. The method of claim 1 wherein said first and second marker fragments comprise the sequence of a primer sequence of a marker selected from the group consisting of Satt435, Satt463, and/or Satt245, and markers that map to within at least about 20 cM of any of these markers.

3. The method of claim 1 wherein said soybean germplasm that has resistance to *Aphis glycines* is germplasm of a variety having a Plant Introductions (PI) number selected from the

group consisting of 71506, 87059, 88508, 200538, 230977, 417084A, 437696, 499955, 507298, 508294, 518726, 548237, 548409, 548445, 548480, 548657, 548663, 567391, 567541B, 567543C, 567597C, 567598B, 587552, 587553A, 587559B, 587617, 587656, 587663, 587664B, 587666, 587668A, 587669, 587674A, 587677, 587682A, 587684A, 587685, 587686A, 587687A, 587693, 587700A, 587702, 587717, 587723A, 587732, 587759, 587763, 587775, 587800, 587816, 587824, 587840, 587844C, 587861, 587863B, 587870, 587871, 587873, 587876, 587877A, 587891A, 587897, 587899, 587905, 587972, 588000, 588040, 594421, 594425, 594426A, 594426B, 594427A, 594431, 594499, 594503, 594514, 594554, 594557B, 594560B, 594573, 594586A, 594592, 594595, 594666B, 594703, 594707, 594711B, 594751A, 594822, 594864, 594868, 594879, 603521, 603530A, 603538A, 603640, 603644, 603650, 605771, 605823, 605855, and 605902, and progeny of these varieties having *Aphis glycines* resistance.

4. The method of claim 1 wherein said soybean germplasm that has resistance to *Aphis glycines* is germplasm of a variety selected from the group consisting of CNS (PI548445) and Dowling (PI548663), and progeny of these varieties having *Aphis glycines* resistance.

5. The method of claim 1 wherein said soybean germplasm that has resistance to *Aphis glycines* is germplasm of a variety selected from the group consisting of Jackson (PI5478657) and Palmetto (PI548480), and progeny of these varieties having *Aphis glycines* resistance.

6. The method of claim 1 wherein said soybean germplasm that has resistance to *Aphis glycines* is germplasm of a variety selected from the group consisting of Moyashimame (PI87059), Sato (PI548409), Showa No. 1-4 (PI88508), Sugao Zarai (PI200538), T260H (PI200538), PI71506, and PI230977 of *G. max*, and G3, JS1, L4, S12 Taichung 38 (PI518282) and Taichung 37 (PI518281) of *G. soja*, and progeny of these varieties having *Aphis glycines* resistance.

7. The method of claim 1 for the production of an inbred soybean cultivar adapted for conferring, in hybrid combination with a suitable second inbred, resistance to *Aphis glycines*, wherein said second soybean germplasm that lacks resistance to *Aphis glycines* is high yielding in hybrid combination with said first soybean germplasm that has resistance to *Aphis glycines*; comprising the further step of self-crossing the plants that tested positive for the presence of *Aphis glycines* resistance until a line is obtained that is homozygous for resistance to *Aphis glycines* and adapted for conferring, in hybrid combination with a suitable second inbred, resistance to *Aphis glycines*.

8. The method of claim 1 wherein said marker fragments comprise the sequence of a primer sequence of a marker selected from the group consisting of Satt435, Satt463, and/or Satt245, and markers that map to within at least about 10 cM of any of these markers.

9. The method of claim 8 wherein said marker that maps to within 20 cM of said Satt435, Satt463 and/or Satt245 markers is a DNA marker selected from the group consisting of SSR, RFLR SNR and RAPD markers.

10. The method of claim 1 wherein said detecting step includes hybridizing at least one polymorphic marker, which is linked to the gene coding for resistance to *Aphis glycines* and maps between about 3 and about 20 cM from Satt435, Satt463, and/or Satt245 markers with nucleic acid of soybean linkage group M of said soybean germplasm.

\* \* \* \* \*