

2020 Fall EECS205002 Linear Algebra

2021/01/13 Quiz 5

1. (50%) Multiple choice questions. Each question may have 0, 1, or more correct choices. For each question, you need to choose all the correct items to get the credit.
 - (a) For an $n \times n$ real matrix A is symmetric, which of the following statements are true?
 - A. If (λ, x) is an eigenpair of A , $\lambda = x^T Ax$.
 - B. If λ is an eigenvalue of A , λ equals to its conjugate $\bar{\lambda}$.
 - C. The eigenvectors of A belonging to distinct eigenvalues are linearly independent.
 - D. There exists a set of n orthonormal eigenvectors of A .
 - E. Matrix A is always diagonalizable.
 - (b) For an Householder reflector $H = I - 2uu^T$, where $\|u\| = 1$, which of the following statements are true?
 - A. H is symmetric.
 - B. $\|Hx\| = \|x\|$ for any vector x that can be pre-multiplied by H .
 - C. $H^{-1} = H$.
 - D. The eigenvalue of H are either 1 or -1.
 - E. $(-1, u)$ is an eigenpair of H .
 - (c) For an $m \times n$ matrix A , $m \geq n$, which of the following statements are true?
 - A. A can be factorized as $U\Sigma V^T$ uniquely, where U and V are orthogonal matrices, and Σ is a diagonal matrix.
 - B. If $m = n$ and A has eigenvalue $\lambda_1, \dots, \lambda_n$, the singular values of A are $|\lambda_1|, \dots, |\lambda_n|$.
 - C. The row vectors in V are the eigenvectors of $A^T A$.
 - D. The column vectors in U are the eigenvectors of AA^T .
 - E. For singular values $\sigma_1, \dots, \sigma_n$, $Au_i = \sigma_i v_i$ for $i = 1, 2, \dots, n$.

- (d) If an $m \times n$ matrix A has rank r , and the SVD of A is $A = U\Sigma V^T$, where $U = [u_1, u_2, \dots, u_m]$ and $V = [v_1, v_2, \dots, v_n]$, which of the following statements are true?

- A. The number zero singular values is $m - r$.
- B. v_1, v_2, \dots, v_r form an orthonormal basis for $R(A)$.
- C. $v_{r+1}, v_{r+2}, \dots, v_n$ form an orthonormal basis for $N(A)$.
- D. u_1, u_2, \dots, u_r form an orthonormal basis for $R(A^T)$.
- E. $u_{r+1}, u_{r+2}, \dots, u_m$ form an orthonormal basis for $N(A^T)$.

- (e) The Frobenius norm of an $m \times n$ matrix A is defined as

$$\|A\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2}, \text{ which of the following statements are true?}$$

- A. If Q is an $m \times m$ orthogonal matrix, $\|Q\|_F = 1$.
- B. If Q is an $m \times m$ orthogonal matrix, $\|QA\|_F = \|A\|_F$.
- C. If A has singular values $\sigma_1, \sigma_2, \dots, \sigma_n$, $\|A\|_F = (\sigma_1^2 + \sigma_2^2 + \dots + \sigma_n^2)^{1/2}$.
- D. If A has rank r , for $0 < k < r$, the minimum of $\|A - X\|_F$ for all possible matrices X of rank k is $(\sigma_1^2 + \sigma_2^2 + \dots + \sigma_k^2)^{1/2}$, where $\sigma_1, \dots, \sigma_k$ are the k largest singular values of A .
- E. If A has rank r , for $0 < k < r$, the matrix of rank k that makes $\|A - X\|_F$ minimum has Frobenius norm $(\sigma_1^2 + \sigma_2^2 + \dots + \sigma_k^2)^{1/2}$, where $\sigma_1, \dots, \sigma_k$ are the k largest singular values of A .


- (f) Which operator L is a linear transformation?

- A. $L([x_1, x_2]) = [x_2, 0, x_1 + x_2]$ where x_1, x_2 are scalars.
- B. $L([x_1, x_2]) = \sqrt{x_1^2 + x_2^2}$ where x_1, x_2 are scalars.
- C. $L(v_1 + v_2) = L(v_1) + L(v_2)$ for all $v_1, v_2 \in V$.
- D. $L([x_1, x_2]) = [\cos(\theta)x_1 - \sin(\theta)x_2, \sin(\theta)x_1 + \cos(\theta)x_2]$ where x_1, x_2 are scalars, and $\theta \in [0, 2\pi)$.
- E. $L(x) = Ax$ for an $m \times n$ matrix A and a vector $x \in \mathbb{R}^n$.

- (g) If L is a linear transformation mapping a vector space V into a vector space W , which of the following statements are true?

- A. The kernel of L is $\{v \in V | L(v) = 0_W\}$.
- B. If S is a subspace of V , the image of S is also a subspace in V .
- C. The range of L is the image of V .
- D. If $L(x) = Ax$ for an $m \times n$ matrix A and a vector $x \in \mathbb{R}^n$, the kernel of L is the null space of A .

- E. If $L(x) = Ax$ for an $m \times n$ matrix A and a vector $x \in \mathbb{R}^n$, the range of L is the row space of A .
- (h) If L is a linear transformation from V into W , where V and W are vector spaces, which of the following statements are true?
- A. L is **one-to-one** if $L(v_1) \neq L(v_2)$ implies $v_1 \neq v_2$.
- B. L is **one-to-one** if and only if $\ker(L) = \{0_V\}$.
- C. L maps V **onto** W if for each $w \in W$ there exists at most one $v \in V$ that $L(v) = w$.
- D. L maps V **onto** W if the image of V is W .
- E. If $L(x) = Ax$ is **one-to-one** and maps V **onto** W , A is nonsingular.
- (i) The following figure illustrates the matrix representation theorem. Which explanations about the figure are correct?



- A. L is a linear transformation mapping \mathbb{R}^n into \mathbb{R}^m .
- B. $E = [e_1, e_2, \dots, e_n]$ is an ordered basis of V ;
 $F = [f_1, f_2, \dots, f_m]$ is an ordered basis of W .
- C. $x = [v]_E$ means $x = [x_1, x_2, \dots, x_n]^T \in \mathbb{R}^n$, $v = x_1 e_1 + \dots + x_n e_n$;
 $y = [w]_F$ means $y = [y_1, y_2, \dots, y_m]^T \in \mathbb{R}^m$, $w = y_1 f_1 + \dots + y_m f_m$.
- D. A is the matrix representing L with respect to \mathbb{R}^n and \mathbb{R}^m .
- E. L maps v into w if and only if $y = Ax$.
- (j) If L is a linear transformation mapping a vector space V into a vector space W , which of the following statements are true?
- A. If $V = \mathbb{R}^2$ and $W = \mathbb{R}^3$, L can be represented as $L(x) = Ax$ for $x \in \mathbb{R}^3$ and a 2×3 matrix A .
- B. If $L([x_1, x_2, x_3]^T) = [x_1 + x_2, x_2 + x_3]^T$ for $x_1, x_2, x_3 \in \mathbb{R}$, then

$$L \left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \right) = x_1 L \left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right) + x_2 L \left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right) + x_3 L \left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right).$$

- C. If V has an ordered basis $E = [v_1, v_2]$, and W has an ordered basis F , for a vector v in V , we can represent L as $[L(v)]_F = A[v]_E$, where $A = [[L(v_1)]_F, [L(v_2)]_F]$.
- D. If L is linear transformation that rotates an \mathbb{R}^2 vector \vec{x} by degree θ , L can be represented as $L(\vec{x}) = A\vec{x}$, where $A = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$.
- E. If L is a differential operator for polynomial of degree 2, L can be represented as $L(ax^2 + bx + c) = A \begin{bmatrix} a & b & c \end{bmatrix}^T$, where A is a 2×3 matrix with column vector $L(x^2), L(x), L(1)$.
2. (10%) If an $n \times n$ matrix A has an eigenvector $e_1 = [1, 0, \dots, 0]^T$, what kind of structure of A should be? Justify your answer.
3. (10%) If A is an $n \times n$ matrix which has eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ and singular values $\sigma_1, \sigma_2, \dots, \sigma_n$, show that $|\lambda_1 \lambda_2 \cdots \lambda_n| = \sigma_1 \sigma_2 \cdots \sigma_n$.
4. (10%) Let $L(x) = [x_1 + x_3, x_3 - x_2]$ for $x = [x_1, x_2, x_3]^T \in \mathbb{R}^3$.
- (a) What is the kernel of L ?
- (b) For $S = \text{span}([1, 1, 0]^T)$, what is the image of S , $L(S)$?
5. (10%) Let $L([x_1, x_2, x_3]) = [x_1 + x_2, x_3 - x_1]^T$, $E = [u_1, u_2, u_3]$ be an ordered basis for \mathbb{R}^3 and $F = [b_1, b_2]$ be an ordered basis for \mathbb{R}^2 , where

$$u_1 = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, u_2 = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, u_3 = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}, b_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, b_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}.$$

Find the matrix representation A of L with respect to E and F .

6. (10%) Matrix polynomial is a polynomial with square matrices as variables. For example, we can put an $n \times n$ matrix A into a polynomial $q(x) = 2x^2 + 3x + 4$, and get a matrix polynomial $q(A) = 2A^2 + 3A + 4I$. Let A be a symmetric matrix, and $p(x) = \det(A - xI)$ be A 's characteristic polynomial. Show that $p(A) = O$, a zero matrix.
7. (20%) There are four subspaces in an $m \times n$ matrix A , row space, column space, null space, and null space of A^T . Write down all the facts you have learned about those four subspaces.