



FULL TEXT LINKS



Meta-Analysis Arch Phys Med Rehabil. 2014 Jun;95(6):1162-73.
doi: 10.1016/j.apmr.2014.02.006. Epub 2014 Feb 26.

Prognostic importance of self-reported traits/problems/strengths and environmental barriers/facilitators for predicting participation outcomes in persons with traumatic brain injury: a systematic review

Mark Sherer ¹, Lynne C Davis ², Angelle M Sander ², Jerome S Caroselli ², Allison N Clark ², Nicholas J Pastorek ³

Affiliations

PMID: 24583024 DOI: 10.1016/j.apmr.2014.02.006

Abstract

Objective: To conduct a systematic review of the prognostic value of self-reported traits/problems/strengths and environmental barriers/facilitators for participation outcomes in persons with traumatic brain injury (TBI).

Data sources: Articles published through August 15, 2013, obtained by conducting electronic searches of PubMed, PsycINFO, and Cumulative Index to Nursing and Allied Health Literature databases and a review of reference lists of reviewed articles.

Study selection: Reviewed articles were written in English and presented findings on adult humans with TBI, participation outcomes, and ≥ 1 self-reported trait/problem/strength (eg, depression, pain, coping style) and/or ≥ 1 environment barrier/facilitator (eg, social support, family functioning, access to services).

Data extraction: Each of the 996 abstracts was examined by 2 reviewers, and those failing to meet all inclusion criteria were excluded. Data were extracted from the 63 retained articles by 2 independent reviewers, who met to resolve any differences in study quality rating or evidence recorded. Study quality was determined using American Academy of Neurology (AAN) criteria.

Data synthesis: Conclusions regarding prognostic importance of self-report and environmental barrier/facilitator variables were made using AAN criteria. Conclusions regarding barrier/facilitator variables indicated that access to transportation, access to services, and participation in social interaction were possibly predictive of employment outcome, whereas living arrangements and social support were possibly not predictive of employment outcome. Conclusions regarding self-report variables indicated that the number of postconcussive symptoms, fatigue, and physical competence were probably predictive of employment and need for supervision, whereas self-efficacy was probably not predictive of employment. Subjective well-being, pain, and social interaction were possibly predictive of employment, whereas coping style was possibly not predictive.

Conclusions: Although additional investigation is needed, self-report variables are likely to make important contributions to predicting participation outcomes. Future research should be guided by coherent conceptual models and use a consistent set of assessment instruments to facilitate comparisons between studies.

Keywords: Brain injuries; Community integration; Rehabilitation; Review [publication type]; Self report.

Copyright © 2014 American Congress of Rehabilitation Medicine. Published by Elsevier Inc. All rights reserved.

[PubMed Disclaimer](#)

Related information

[Cited in Books](#)

[MedGen](#)

LinkOut – more resources

Full Text Sources

[ClinicalKey](#)

[Elsevier Science](#)

Other Literature Sources

[scite Smart Citations](#)

Medical

[MedlinePlus Health Information](#)

Miscellaneous

[NCI CPTAC Assay Portal](#)