

Gaming Disorder Test (GDT)

Please cite. Pontes, H. M., Schivinski, B., Sindermann, C., Li, M., Becker, B., Zhou, M., & Montag, C. (2021). Measurement and conceptualization of gaming disorder according to the World Health Organization framework: The development of the Gaming Disorder Test. *International Journal of Mental Health and Addiction*, 19(2), 508–528. <https://doi.org/10.1007/s11469-019-00088-z>

Administration. Ask the screening question first and keep the 12-month time frame. Code answers 1 to 5; the total score is the sum of the four items (4 to 20). An item is endorsed when answered 4 or 5; endorsing all four items classifies a potentially disordered gamer for research purposes. The GDT measures severity and does not replace a clinical diagnosis. Instructions, items and response format are as published.

Instructions

The questions below are about your gaming activity during the past year (i.e., last 12 months). Here, gaming activity means any gaming-related activity that has been played either from a computer/laptop or from a gaming console or any other kind of device (e.g., mobile phone, tablet) both online and/or offline.

Part 1. Screening

Have you played any video game in the past 12 months?

☐ Yes ☐ No

For researchers: Respondents who answer “No” do not continue. This is the eligibility question used in the article.

Part 2. Gaming disorder

Please indicate how often the following issues occurred on average over the past twelve months until today.

		Never	Rarely	Sometimes	Often	Very often
GDT1	I have had difficulties controlling my gaming activity.	☐	☐	☐	☐	☐
GDT2	I have given increasing priority to gaming over other life interests and daily activities.	☐	☐	☐	☐	☐
GDT3	I have continued gaming despite the occurrence of negative consequences.	☐	☐	☐	☐	☐
GDT4	I have experienced significant problems in life (e.g., personal, family, social, education, occupational) due to the severity of my gaming behavior.	☐	☐	☐	☐	☐