
Check B(ii): predicted Lazuli per-bin precision against published HST and JWST precision
WASP-39 b Feinstein NIRISS
WASP-39 b Rustamkulov PRISM
GJ 1214 b Kreidberg WFC3
K2-18 b Benneke WFC3
K2-18 b Tsiaras WFC3
GJ 3470 b Benneke WFC3
WASP-107 b Spake WFC3
WASP-12 b Kreidberg WFC3
HD 209458 b Sing WFC3
HAT-P-11 b Fraine WFC3
WASP-107 b Kreidberg WFC3
WASP-121 b Evans WFC3
WASP-12 b Stevenson WFC3
WASP-39 b Wakeford WFC3
HD 189733 b Pont ACS
HD 209458 b Deming WFC3
HD 189733 b Sing STIS
WASP-39 b Fischer STIS
WASP-39 b Sing STIS
GJ 1214 b Berta WFC3
HD 209458 b Sing STIS
GJ 3470 b Benneke STIS
HD 189733 b Pont STIS
HD 189733 b Sing WFC3
WASP-12 b Sing STIS
WASP-12 b Mandell WFC3
GJ 3470 b Ehrenreich WFC3
HD 189733 b McCullough WFC3
HD 189733 b Pont NICMOS
equal precision
2.30
2.74
1.63
1.63
0.92
JWST dataset (single transit each)
HST dataset (indicative: N transits not in the archive)
area-only expectation vs JWST: 1.90
area-only expectation vs HST: 0.80
10^{-1}
10^{0}
Lazuli predicted σ (1 transit) / published σ, same bins