

Math 4460 Course Notes

August 29, 2024

Contents

1	Lecture 1	1
1.1	Administrativa	1
1.1.1	Canvas	1
1.2	Course Overview	2
1.2.1	Introduction	2
1.2.2	The Standard Diagram	2
1.2.3	Themes	2
1.3	Zeroth Assignment	3
1.4	Probability	3
1.4.1	Context	3
1.4.2	Intuition for Measure from Calculus	4
1.5	The Formalism: Axiomatizing Measure	5
2	Lecture 2	8
2.1	Random Variables	8
2.2	Expectation	10
2.3	Joint Measures	11
2.4	Marginalization	11
2.5	Conditional Probability	12
2.6	Independence	13
2.7	Data	14
3	Lecture 3	15
3.1	Independence	15
3.2	Data	16
3.3	Concentration Inequalities	16
3.3.1	Law of Large Numbers (LLN)	17
3.4	Monkeys on a Keyboard: Intuition in High Dimension	18
4	Lecture 4	20
4.1	Setting: Detection / Binary Classification	20
4.2	Interpreting the Score	20
4.2.1	Optimal Threshold	21
4.3	Constructing the Score	22
4.4	Metrics	23

1 Lecture 1

1.1 Administrativa

1.1.1 Canvas

Assignments, starter code, and data will be housed in canvas. Please check weekly for newly posted assignments.

1.2 Course Overview

1.2.1 Introduction

From the syllabus:

Machine Learning describes a mishmash of computational techniques for “finding patterns in data.” The scope of use, analytic tools, algorithms, and results are almost too numerous to meaningfully batch all such applications under a common appellation. Still, we try. This course focuses on *supervised* machine learning (sml) which roughly deals with using historical *labeled* data to construct a predictor which will correctly label future data.

Formally, we will be operating in space $\mathcal{X} \times \mathcal{Y}$, where \mathcal{X} denotes “input” space, and \mathcal{Y} “output.” While these notions are heuristic, they well frame the situation that we may easily sample data at will from \mathcal{X} , while sampling from \mathcal{Y} may be difficult or expensive, and often we would like to decision according to how we believe $x \in \mathcal{X}$ is associated with label $y \in \mathcal{Y}$.

In this course, you will learn how to formulate the supervised learning problem in mathematical terms, how to describe a measure of performance, restrict search space for constructing models for prediction, optimize performance measure in search space, and how to check for generalization. You will learn, also, how to implement some of these methods in code, from the ground up, as well as incorporating pre-built libraries (such as pytorch) for such tasks. Finally, you will learn how to articulate learning guarantees, and understand some of the limits of learning claims. While this course is primarily theory-centric, there will be no dearth of opportunity for employing concrete computational techniques.

1.2.2 The Standard Diagram

Describing our problem space in more detail, consider the following diagram

$$\begin{array}{ccc} \mathcal{X} \times \mathcal{Y} & \xrightarrow{\pi_{\mathcal{Y}}} & \mathcal{Y} \\ \downarrow \pi_{\mathcal{X}} & \nearrow \tilde{y} & \\ \mathcal{X} & & \end{array} \quad (1)$$

We will refer to this diagram often, and to do so give it the somewhat non-descriptive, but in our context wholly unambiguous, name ‘the standard diagram.’

Traditionally, $\pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathcal{X}$, defined by mapping $(x, y) \mapsto x$ (read: $\pi_{\mathcal{X}}(x, y) := x$), is taken to be “easy, efficient, or cheap” to evaluate or sample while $\pi_{\mathcal{Y}} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathcal{Y}$, defined by mapping $(x, y) \mapsto y$, is computationally expensive, expensive otherwise, difficult for other reasons, or altogether infeasible. The original space $\mathcal{X} \times \mathcal{Y}$ is itself inaccessible, except for some (finite) *labeled* data $S = \{(x_1, y_1), \dots, (x_m, y_m)\} \subset \mathcal{X} \times \mathcal{Y}$ which provides a proxy (and incomplete!) illustration of what $\mathcal{X} \times \mathcal{Y}$ looks like. The map $\tilde{y} : \mathcal{X} \dashrightarrow \mathcal{Y}$ is a critter we’d like to construct from data S so that both $\tilde{y}(x) \approx y$ for $(x, y) \in S$ and for (arbitrary) $(x, y) \in \mathcal{X} \times \mathcal{Y}$. What “ \approx ” means, how to construct \tilde{y} , conditions on S which are needed to make this problem feasible, etc. are all aspects of the supervised machine learning problem which we will explore in this course.

As an example, suppose $\mathcal{X} = \mathbb{R}$ denotes credit score and $\mathcal{Y} = \{0, 1\}$ loan repayment (say ‘1’ corresponds to repayment of loan, ‘0’ to default). Then a *point* $(x, y) \in \mathbb{R}$ represents data corresponding to a loan whose account holder has credit score x and for which the loan was either paid in full ($y = 1$) or not ($y = 0$). The reason we say $\pi_{\mathcal{X}}$ is “easy” to sample is that you may ask any person what their credit score is (more realistically: as creditor, you would see this information *at the time of application*), while loan repayment information (the “label”) would not be observed until potentially many years later when the loan is finally repaid or defaults.

It is worth noting, and perhaps lingering upon the observation, that the “input-output” relation (x, y) is not necessarily functional, i.e. for two points $(x, y), (x', y') \in \mathcal{X} \times \mathcal{Y}$, $x = x'$ does not imply that $y = y'$. The stand-in for determinism is probability, i.e. we will presume that there is some joint probability measure $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ on $\mathcal{X} \times \mathcal{Y}$, and often that $\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y = f(x)|x) \neq 1$ for any function $f : \mathcal{X} \rightarrow \mathcal{Y}$.

1.2.3 Themes

Generalization Given model $\tilde{y} : \mathcal{X} \rightarrow \mathcal{Y}$, how well does \tilde{y} match the (finite) data we have $S \subseteq \mathcal{X} \times \mathcal{Y}$ —i.e. $\tilde{y}(x) \approx y$ for $(x, y) \in S$ —and the data we don’t have, $(x, y) \in \mathcal{X} \times \mathcal{Y}$?

Dimensionality Computation in high dimensions becomes harder, in part because computation is more expensive, and because there are more “corners” for data to hide in (which exacerbates the computational problem). The geometry of high dimensionality will be a recurring theme; for now, we simply observe sources of high dimensionality:

1. The data “set” itself $S = \{(x_1, y_1), \dots, (x_m, y_m)\} \subset \mathcal{X} \times \mathcal{Y}$. Properly speaking, this data will be presumed to be sampled $(x_i, y_i) \sim_{\text{iid}} \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ which *means* that the data “set” is a *point* in the (higher dimensional) space $(\mathcal{X} \times \mathcal{Y})^m$. A reasonable question to ask, then, is: what is the measure $\mathbb{P}_{(\mathcal{X} \times \mathcal{Y})^m}$? Independence tells us that it is $\prod_{j=1}^m \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$. More on this later.
2. Size of space itself. This could include high dimensionality of \mathcal{X} and/or \mathcal{Y} . Examples abound of high dimensional input data: numerous columned tabular data, imagery data, audio data, video data.
3. Parameter space for model \tilde{y} . In the case of linear regression, a model $\tilde{y}(x) = \sum_{j=0}^n a_j x^j$ may have arbitrarily large degree n . Or a fully connected neural network with many nodes and many layers. And so on. More generally, the (full) space of functions $\mathcal{Y}^{\mathcal{X}} := \{\tilde{y} : \mathcal{X} \rightarrow \mathcal{Y}\}$ is even bigger.

Trade-offs Assumptions must be made and compromises allowed for in order to gain tractability in the learning problem. There is no universal solution—“no free lunch,” and as you may imagine, there’s a theorem for that—and formulating the setup to address one challenge may introduce other ones elsewhere (ML can sometimes feel like one giant game of whackamole).

The famed bias-variance trade-off is one example: a high complexity model may well represent the data S , which in one sense is good, but in another is bad if said model represents data *too well*, i.e. at the exclusion of modeling ‘from where the data comes.’

1.3 Zeroth Assignment

Programming Assignment You may find the first programming assignment under pa0 on Canvas,¹ along with starter code and data. I suggest you follow the tutorial at Real Python [real python](#)² which shows you how to spin up a logistic regression model using sklearn. Scikit-Learn (also known as sklearn) is an open source ML library for python, and contains functionality for constructing numerous models. This is perhaps the only time in this course you will be asked to use this library, and if you have a preferred alternative library, you are more than welcome to use it for this assignment.

The purpose of the assignment is threefold:

1. Gain initial exposure to the *structure* of machine learning code, including object oriented programming and the typical methods included.
2. Shake off any residual rust using Python.
3. To gain deeper appreciation for the *aim* of building a model $\tilde{y} : \mathcal{X} \rightarrow \mathcal{Y}$ as in diagram (1), and metrics that illustrate success.

1.4 Probability

1.4.1 Context

Recall the Standard Diagram

$$\begin{array}{ccc} \mathcal{X} \times \mathcal{Y} & \xrightarrow{\pi_{\mathcal{Y}}} & \mathcal{Y} \\ \downarrow \pi_{\mathcal{X}} & \nearrow \tilde{y} & \\ \mathcal{X} & & \end{array} \quad (2)$$

¹I index pythonically.

²You may need to sign up in order to view this content, but it is not behind a paywall.

This diagram provides formalism for talking about “approximating” y with model $\tilde{y} : \mathcal{X} \rightarrow \mathcal{Y}$ when $y \neq y(x)$ is not necessarily functionally determined by x .

Consider a concrete example to illustrate the problem: suppose that $\mathcal{X} = \mathbb{R}$ denotes credit score and $\mathcal{Y} = \{0, 1\}$ denotes repayment on loan, 1 denotes full repayment and 0 denotes default. A data point $(x, y) \in \mathcal{X} \times \mathcal{Y}$ corresponds to a credit score-loan repayment pair, and in real life a loan account would have these attributes associated with it, i.e. the debtor would have some credit score and their loan will (eventually) be repaid or not. (Note that “eventuality” is what, in this case, makes $\pi_{\mathcal{Y}}$ hard or expensive to evaluate.) It is possible for two different loans, belonging to two different people, to agree on credit score but disagree on outcome $y \in \mathcal{Y}$. In fact, we will likely observe both outcomes $y = 0$ and $y = 1$ associated to *any* credit score. Presumably, there should be some relation between the relative *counts* of $\#y = 1$ and credit score; in other words, one might suppose that lower credit scores correspond to accounts which in actual fact get repaid less frequently than those with high credit scores. We need mathematical language to describe and work with this phenomenon. The language is probability.

Thus, we suppose that there is some joint probability measure $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ on $\mathcal{X} \times \mathcal{Y}$. One way of contextualizing supervised machine learning is as a study of probability on the standard diagram. In this lecture, we will review probability, give intuition for probability as measure, as well as notation $\mathbb{P}_{\mathcal{X}} = \int d\mathbb{P}_{\mathcal{X}}(x)$, and define expectation $\mathbb{E}(f)$ of a random variable $f : \mathcal{X} \rightarrow \mathbb{R}$ as a *Lebesgue integral* $\mathbb{E}(f) := \int_{\mathcal{X}} f(x) d\mathbb{P}_{\mathcal{X}}(x)$. I hesitate in bringing in the Lebesgue integral, if it weren’t for that expectation of discrete random variables just *is* defined as a Lebesgue integral. We shall bite the bullet and lean on the concept in discussing arbitrary random variables as well.

We kick this lecture off by asserting that probability has nothing to do with randomness... at least not yet. When we return to admitting randomness into our lexicon, it will be as a *result*. For now, we forget any association between probability and chance, stochasticity, randomness, or any other (for now) anathema word affiliated with the notion of uncertainty. If you find yourself tempted to rely on the association, make an honest attempt to circle back and fight the temptation.

1.4.2 Intuition for Measure from Calculus

We start reviewing integration in calculus to preview notation for measure. One interpretation of the integral is as “area under the curve.” Another (what amounts to similar) is as measure. And one interpretation of dQ is as an infinitesimal Q element, whatever Q is. Another is as an indicator of what kind of measurement we are taking. We then express length ℓ as $\int d\ell$, area a as $\int da$, volume v as $\int dv$, measure m as $\int dm$, and so on. Note that the expression $\int dm$ is defined in terms of measure m , i.e. we suppose prior (at the very least conceptual) knowledge of the measure, irrespective of whether or not given a particular object to measure A , we are actually able to evaluate its measure $m(A)$.

Example: length We’ve learned that the length $\ell([a, b])$ of an interval $[a, b]$ is the difference of endpoints $b - a$. We can write this with an integral as $\ell([a, b]) = \int_a^b dx$ or more in line with notation we will use, as $\ell = \int_{[a, b]} d\ell(x)$. In this notation, the subscript is the object we are measuring, ℓ is the type of measure, and x is a dummy variable. Until we start integrating functions, we won’t need it, and we could have just written $\int_{[a, b]} d\ell$. The dummy variable was kept only to clearly delineate that $d\ell$ is not the same thing as dx : we are using calculus intuition only for loose guidance.

General Formula For measure m and object to be measured A , we write $m(A) = \int_A dm$. Right now, do not put too much stock in the—what in calculus would be thought of as infinitesimal— dm term: it is *defined* as a *phrase* with the integral \int : neither in isolation makes sense, and the definition goes this

way: $\int_A dm := m(A)$. Thus, you need to first know the measure. We'll return to this momentarily.

Example: Area




Figure 1: Area

In our notation area A of region R is $A(R) = \int_R dA$. If you want to import calculus intuition, when we interpret dA as a differential area element, we may express it likewise as the *product* $dA = f(x)dx$, where $f(x)$ represents height and dx the differential width. As area is equal to height times width (or length) a differential area is a length element times a differential length element and we recover the standard form $A(R) = \int_a^b f(x)dx$.

Example: Volume




Figure 2: Volume

Using the above picture, $V(S) = \int_S dV$, where $V(S)$ is the volume. Therefore, by generalizing the above idea we can write $\mathbb{P}(A) = \int_A d\mathbb{P}$.

1.5 The Formalism: Axiomatizing Measure

Now we formalize what we mean by 'probability measure.' The first thing to say is that it is a measure.

Definition 1.1. Let \mathcal{X} be a set. We define a *probability measure*

$$\mathbb{P}_{\mathcal{X}} : ([\text{Some}] \text{ Subsets of } \mathcal{X}) \rightarrow [0, 1]$$

on \mathcal{X} to be a map from (a subset of)³ the power set of \mathcal{X} to the closed interval $[0, 1]$ satisfying the following two properties:

³This subtle point is a technicality beyond the scope of this course, and quite frankly unnecessary for reasonably understanding measure. Measure measures subsets. But it's possible that it may not be able to measure *all* subsets. So the domain of the measure may not be *all* subsets. A lot of work goes into specification of the structure of the collection of subsets you can measure, and like topology is characterized by closure operations, e.g., if you can measure $\{A_j\}_{j \in \mathbb{N}}$ then you can measure its union $\bigcup_{j \in \mathbb{N}} A_j$. For the curious, you may look into sigma algebras for more detail.

1. $\mathbb{P}_{\mathcal{X}}(\mathcal{X}) = 1$ (space is measure finite and normalized), and
2. $\mathbb{P}_{\mathcal{X}}\left(\bigsqcup_{j=1}^{\infty} A_j\right) = \sum_{j=1}^{\infty} \mathbb{P}_{\mathcal{X}}(A_j)$ where $\bigsqcup_{j=1}^{\infty} A_j$ denotes disjoint union, i.e. as a set $\bigsqcup_{j=1}^{\infty} A_j = \bigcup_{j=1}^{\infty} A_j$ and $A_i \cap A_j = \emptyset$ for $i \neq j$ (countable additivity).

Recall that the power set $2^{\mathcal{X}}$ of a set \mathcal{X} is defined to be the set of all subsets of \mathcal{X} , including \emptyset and \mathcal{X} itself. For example, when $\mathcal{X} = \{1, 2, 3\}$,

$$2^{\mathcal{X}} = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \mathcal{X}\}.$$

When $\mathcal{X} = \mathbb{R}$, the power set includes any reasonable combination (e.g. unions) of intervals (a, b) or $[a, b]$, but also many many more (see [Cantor Set](#) for a fun, but not particularly relevant, excursion into the wonders of measure theory).

The second condition is called ‘countable additivity’ (informally: a conservation of stuff principle) and represents the intuitive idea that if you slice and dice an object for measurement, measure each constituent piece without double counting, and add your results, you’ll end up with the same result as if you just measured the whole original unadulterated tamale.

You should check that countable additivity implies *finite* additivity $\mathbb{P}_{\mathcal{X}}\left(\sum_{j=1}^n A_j\right) = \sum_{j=1}^n \mathbb{P}_{\mathcal{X}}(A_j)$.

You will need the fact that $\mathbb{P}_{\mathcal{X}}(\emptyset) = 0$, which itself is implied by conditions 1. and 2. Indeed, $\mathcal{X} = \mathcal{X} \sqcup \bigsqcup_{j=2}^{\infty} \emptyset$. Also check a very useful second fact, the Union Bound: $\mathbb{P}(\cup_{j \in \mathbb{N}} A_j) \leq \sum_{j \in \mathbb{N}} \mathbb{P}(A_j)$. You may find it worthwhile to verify that $A \subset B$ implies that $\mathbb{P}_{\mathcal{X}}(A) \leq \mathbb{P}_{\mathcal{X}}(B)$.

Remark 1.1. We are being fairly blasé about the *domain* “some subsets of \mathcal{X} ” of $\mathbb{P}_{\mathcal{X}}$. Perhaps we shouldn’t be. Much of the scaffolding around constructing measure depends on the fact that some structure must be placed on the set of subsets of \mathcal{X} which are suitable for measurement; in particular, that such a set comprises a so-called σ -algebra, is not necessarily (and in many cases in fact is not) the entire power set $2^{\mathcal{X}} = \{A \subset \mathcal{X}\}$, and so on. We pay lip-service to this nuance, but fret little over the possibility that we will accidentally run across both a measure $\mathbb{P}_{\mathcal{X}}$ and subset $A \subset \mathcal{X}$ for which $\mathbb{P}_{\mathcal{X}}(A)$ does not (and fundamentally cannot) make sense (read: which $\mathbb{P}_{\mathcal{X}}$ is “incapable” of measuring). One must try very hard—you might find such a question on a measure theory qualifying exam—to come up with such an example. Therefore you may reasonably suppose that any set you’d come across in real life is in fact measurable. Still, know *that* there is a potential problem: if you can construct a non-measurable set, then you can also cut an apple into finitely many pieces and reassemble those finitely many pieces into *two* apples of the same size (see [Banach Tarski](#) for more information). In other words, weird things can happen with things that aren’t measurable.

Definition 1.2. We define a *probability space* to be a pair $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ where \mathcal{X} is a set and

$$\mathbb{P}_{\mathcal{X}} : ([\text{Some/Many}] \text{ Subsets of } \mathcal{X}) \rightarrow [0, 1]$$

is a probability measure (c.f. definition 1.1).

In probabilistic terminology, measurable subsets $A \subset \mathcal{X}$ are often called ‘events’ and individual points $x \in \mathcal{X}$ called ‘outcomes.’ An outcome $x \in \mathcal{X}$ defines the *event* $\{x\} \subset \mathcal{X}$ with the single outcome $x \in \{x\}$.

Example 1 Let $(\mathcal{X} = [0, 1], \mathbb{P}_{\mathcal{X}}([a, b]) := b - a)$ for $0 \leq a \leq b \leq 1$. We define notation $\int_{[a, b]} d\mathbb{P}(x) = \mathbb{P}([a, b])$.

Example 2 Let $(\mathcal{X} = \mathbb{R}, \mathbb{P}_{\mathcal{X}}([a, b]) := \frac{1}{\sqrt{2\pi}} \int_a^b e^{-x^2/2})$, the so-called normal distribution with zero mean and unit variance. Observe that we use a Riemann integral to *compute* or give the rule for realizing the probability measure. The integrand $\frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ of the Riemann integral is called a *probability*

density function. The integral $\int_{[a,b]} d\mathbb{P}_{\mathcal{X}}(x)$, by contrast, is not a Riemann integral; it is *defined* by the measure $\mathbb{P}_{\mathcal{X}}([a,b])$. (When you ask: but what is the measure?, we gave the rule for how to calculate it!)

Example 3 Let $\mathcal{X} = \mathcal{Y} = [0, 1]$, and define $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}([x_1, x_2] \times [y_1, y_2]) := (x_2 - x_1) \cdot (y_2 - y_1)$. This example is a preliminary look into independence as $\mathbb{P}_{\mathcal{X}}([x_1, x_2]) = x_2 - x_1$, $\mathbb{P}_{\mathcal{Y}}([y_1, y_2]) = y_2 - y_1$ so the measure on product space as defined is also $\mathbb{P}_{\mathcal{X}}([x_1, x_2])\mathbb{P}_{\mathcal{Y}}([y_1, y_2])$. We will return to this example when we discuss independence, which we'll want to situate as a notion which properly at home in high dimension: independence really is a high dimensional phenomenon and it is clear that we can generalize the picture in 3 to n dimensions.




Figure 3: The area measure

Example 4 : Bernouli Random Variable (flipping a fair or unfair coin): Let $\mathcal{X} = \{0, 1\}$ where $\mathbb{P}_{\mathcal{X}}(\{1\}) = p$ and $\mathbb{P}_{\mathcal{X}}(\{0\}) = 1 - p$. This is a probability space with two outcomes that clearly satisfies the axioms stated earlier since the events or subsets of the space \mathcal{X} , $\{0\}$ and $\{1\}$ are disjoint and the sum of their probabilities is 1:

$$\mathbb{P}_{\mathcal{X}}(1) + \mathbb{P}_{\mathcal{X}}(0) = p + (1 - p) = 1 = \mathbb{P}_{\mathcal{X}}(\{0\} \sqcup \{1\})$$

2 Lecture 2

2.1 Random Variables

Recall that a probability space $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ consists of a set \mathcal{X} and a measure $\mathbb{P}_{\mathcal{X}}$ (see definition 1.2). We said that probability is a theory of measure, which really means that it's a theory of integration, and integration deals with numbers. So far, we've said nothing about the nature of the set \mathcal{X} , and we don't need to. What we do need is a way to associate numerical values to outcomes $x \in \mathcal{X}$.

Definition 2.1. Let $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ be a probability space. We define a *random variable* to be a function $f : \mathcal{X} \rightarrow \mathbb{R}$, whose codomain is \mathbb{R} .




Figure 4: Random variable f

Such a function induces a measure $\mathbb{P}_{\mathbb{R}}$ on \mathbb{R} , defined by

$$\mathbb{P}_{\mathbb{R}}([a, b]) := \mathbb{P}_{\mathcal{X}}(f^{-1}([a, b])) = \mathbb{P}_{\mathcal{X}}(\{x \in \mathcal{X} : f(x) \in [a, b]\}). \quad (3)$$

One should check that this defines an honest probability measure on \mathbb{R} .

Remark 2.1. It is worth noting that certain conditions must be met by the map $f : \mathcal{X} \rightarrow \mathbb{R}$ in order to ensure that the induced measure $\mathbb{P}_{\mathbb{R}}$ is well-defined, i.e. that it is a true (probability) measure. In essence, we require a condition known as ‘measurability,’ that f must be a *measurable* function (which really just means: f is such that the induced measure is a measure—this isn’t circular!, it all comes down to saying, you cannot with impunity claim that any function whatsoever will induce a measure). Just as with the domain of $\mathbb{P}_{\mathcal{X}}$, where we suppose with little guilt that “all subsets” we encounter are measurable, we will also suppose that the functions we come across in practice are measurable. Again, there is nuance to be appreciated and after pausing to make the quick remark will proceed to not appreciating it: the supposition that “everything is measurable” (now including functions) will very unlikely harm any of our day to day calculations.

(For the ultra-curious, the condition of measurability stipulates that any potentially measurable set in \mathbb{R} has preimage (by f^{-1}) which is measurable in \mathcal{X} , so really everything comes down to the notion of measurability of events in the first place!)

Remark 2.2. We noted that $f : \mathcal{X} \rightarrow \mathbb{R}$ induces a measure. In fact, this holds more generally for any (measurable) map $f : \mathcal{X} \rightarrow \mathcal{Y}$, $\mathbb{P}_{\mathcal{Y}}(B) := \mathbb{P}_{\mathcal{X}}(f^{-1}(B))$.

Remark 2.3. While we have in definition 2.1 defined a random variable to be a *function*, we will often (also) call the codomain $(\mathbb{R}, \mathbb{P}_{\mathbb{R}})$ with its induced measure a ‘random variable,’ without concern for the domain which produced it.

Definition 2.2. Let $f : \mathcal{X} \rightarrow \mathbb{R}$ be a random variable. We define *expectation* of f , denoted $\mathbb{E}(f)$, to be

$$\mathbb{E}(f) := \int_{\mathcal{X}} f(x) d\mathbb{P}_{\mathcal{X}}(x). \quad (4)$$

Equation (4) defines expectation, but we have some odd sort of critter we’ve not seen before on the right hand side. We must define *it*. We will do this in steps, 1. first for discrete random variable (taking finitely many values) and 2. by extension to arbitrary random variables.

Definition 2.3. Let $f : \mathcal{X} \rightarrow \mathbb{R}$ be a *simple* random variable, i.e. taking *finitely* many values a_1, \dots, a_k . Then we define the *Lebesgue* integral of f , denoted $\int_{\mathcal{X}} f(x) d\mathbb{P}_{\mathcal{X}}(x)$, to be

$$\int_{\mathcal{X}} f(x) d\mathbb{P}_{\mathcal{X}}(x) := \sum_{j=1}^k a_j \mathbb{P}_{\mathcal{X}}(f = a_j) = \sum_{j=1}^k a_j \mathbb{P}_{\mathcal{X}}(\{x \in \mathcal{X} : f(x) = a_j\}).$$




Figure 5: Simple random variable f

One may recognize this definition as the *expectation of a discrete random variable*. Indeed, that is exactly what it is. It is the Lebesgue integral! The Lebesgue integral extends to continuous random variables, but it requires some abstraction. Before doing so, observe that for continuous random variable $f : \mathcal{X} \rightarrow \mathbb{R}$ (not necessarily taking finitely many values), we can *approximate* $\mathbb{E}(f)$ using the Lebesgue integral of a simple random variable (expectation of discrete r.v.), see fig. 6.




Figure 6: Approximation via simple functions

Remark 2.4. In the following, we will continue the definition of Lebesgue integral for continuous random variable. While there are theoretical reasons for insisting on the use of this abstraction, ours are more practically oriented. In fact, one may (very) often compute expectation *using* instead a Riemann integral. For example, the expectation of a mean 0 unit variance normally distributed random variable is $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} x e^{-x^2/2} dx$ (which of course is zero). In other words, if push comes to shove and you're asked to *compute* the expectation for a continuous random variable, very often you'll have a density function in your pocket and can just multiply it by the random variable, and integrate, business as usual. We introduce Lebesgue integration for two reasons:

1. To articulate the fact that the extension from discrete to continuous random variables may *seem* confusing, and that is due in part to the nauseating head-spinning move from Lebesgue to

who knows (but usually Riemann) integration without even lip-service paid to the fact that expectation of discrete r.v.s itself is a non-trivial extension of our conceptual apparatus.

2. Ease of notation: $d\mathbb{P}_{\mathcal{X}}$ always makes sense and makes immediately obvious what our measure is. In other words, I don't always want to say: suppose that the density of a probability space exists, and anyway it might not and that doesn't matter!, for $\int_{\mathcal{A}} d\mathbb{P}_{\mathcal{X}}(x)$ makes sense regardless of whether we can integrate (Riemann-wise) as $\int \varphi(x)dx$ (for density $\varphi(x)$). Related: the notation $d\mathbb{P}_{\mathcal{X}}$ collapses the distinction between continuous and discrete random variables. The distinction, in my view, is convoluted and confusing, especially when we have mixed discrete-continuous nonsense going on (e.g. in the case of binary classification $\mathcal{X} = \mathbb{R}$, $\mathcal{Y} = \{0, 1\}$). Of course, successfully working with the collapse requires an comfort with the abstraction, and that may by itself be initially confusing as well.

Remark 2.5. Observe also that the notation $\mathbb{E}(f)$ is somewhat uninformative, in a way that $\int_{\mathcal{X}} f(x)d\mathbb{P}_{\mathcal{X}}(x)$ is not. At the moment the added notational baggage of the latter may seem inconvenient, but once we start squinting at joint probability spaces $\mathcal{X} \times \mathcal{Y}$, turning them upside down, and so on, it will be imperative to be clear on how we are integrating. For example, we will see

$$\int_{\mathcal{X} \times \mathcal{Y}} f(x, y) d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(x, y) = \int_{\mathcal{X}} \int_{\mathcal{Y}|\mathcal{X}} f(x, y) d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) d\mathbb{P}_{\mathcal{X}}(x).$$

This is sort of an extension on the notational point above. I will be relying on this notation aggressively, so it's crucial to anticipate eventually becoming comfortable with standard manipulations, and recalling (any time there is lingering confusion) that $d\mathbb{P}_{\mathcal{X}}$ is defined, not in isolation, but as a package $\int_{\mathcal{A}} d\mathbb{P}_{\mathcal{X}} := \mathbb{P}_{\mathcal{X}}(A)$.

2.2 Expectation

Recall definition 2.2 for which $\mathbb{E}(f) = \int_{\mathcal{X}} f(x) d\mathbb{P}_{\mathcal{X}}(x)$ for a random variable $f : \mathcal{X} \rightarrow \mathbb{R}$ on the probability space $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$. While this definition lays out the concept of expectation, it leaves us pondering what defines the integral itself. We can draw insight from discrete (or simple) random variables like in definition 2.3.

Definition 2.4. Let $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ be a probability space and $f : \mathcal{X} \rightarrow \mathbb{R}^{\geq 0}$ a nonnegative random variable. Then we define the *Lebesgue integral* of f as

$$\int_{\mathcal{X}} f(x) d\mathbb{P}_{\mathcal{X}}(x) := \sup \{ \mathbb{E}(\bar{f}) : 0 \leq \bar{f} \leq f \text{ is simple random variable} \}. \quad (5)$$

A visual illustration is provided in ??.⁴

We provide this definition because it is the definition given to us of expectation. The move to continuous random variables may seem abstract—indeed, we discretize the codomain \mathbb{R} instead of domain \mathcal{X} as we are used to doing with Riemann integration from calculus—but one may rest assured that in many instances expectation *may* be computed as a Riemann integral instead. In particular, when a *density* function exists, $\rho : \mathcal{X} \rightarrow \mathbb{R}$ such that $\mathbb{P}_{\mathcal{X}}([a, b]) = \int_a^b \rho(x)dx$, we may compute expectation using the following procedure:

1. multiply the random variable f by density ρ ; this will give us a function $f \cdot \rho : \mathcal{X} \rightarrow \mathbb{R}$, and
2. compute the Riemann integral $\int_{-\infty}^{\infty} f(x)\rho(x)dx$.

When the density does not exist, the definition of expectation still makes sense, and one cannot resort to this procedure for computation. Thus for the sake of understanding, one may choose to delve into the nuance of Lebesgue integration, or pretend, as we've been pretending about measurable sets and measurable functions, that “everything” can be computed as a Riemann integral.

⁴An Introduction to the Lebesgue Integral, Ikhlas Adi, 2017

Example 2.1. We point out that probability may be computed as expectation:

$$\mathbb{P}_{\mathcal{X}}(A) = 1 \cdot \mathbb{P}_{\mathcal{X}}(A) + 0 \cdot \mathbb{P}_{\mathcal{X}}(\mathcal{X} \setminus A) = \int_{\mathcal{X}} \mathbb{1}_{x \in A} d\mathbb{P}_{\mathcal{X}}(x) = \mathbb{E}(\mathbb{1}_{x \in A}).$$

Observe that this Lebesgue integral uses the definition from definition 2.3; we do not need to rely on any limiting procedure (as in e.g. definition 2.4).

For the sake of completeness, we define expectation for arbitrary r.v. $f : \mathcal{X} \rightarrow \mathbb{R}$.

Definition 2.5. Let $f : \mathcal{X} \rightarrow \mathbb{R}$ be a random variable and suppose that $\mathbb{E}(f \cdot \mathbb{1}_{f \geq 0}) < \infty$ and $\mathbb{E}(-f \cdot \mathbb{1}_{f < 0}) < \infty$ (both expectations of non-negative random variables). Then we define

$$\mathbb{E}(f) := \mathbb{E}(f \cdot \mathbb{1}_{f \geq 0}) - \mathbb{E}(-f \cdot \mathbb{1}_{f < 0})$$

Remark 2.6. Expectation is already defined as $\mathbb{E}(f) = \int_{\mathcal{X}} f(x) d\mathbb{P}_{\mathcal{X}}(x)$. This definition is defining the right hand side.

Definition 2.6. Let $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ and $(\mathcal{Y}, \mathbb{P}_{\mathcal{Y}})$ be probability spaces. A map $f : (\mathcal{X}, \mathbb{P}_{\mathcal{X}}) \rightarrow (\mathcal{Y}, \mathbb{P}_{\mathcal{Y}})$ of probability spaces is a map from \mathcal{X} to \mathcal{Y} that respects the measure i.e.

$$\mathbb{P}_{\mathcal{Y}}(B) = \mathbb{P}_{\mathcal{X}}(f^{-1}(B))$$

for all $B \subseteq \mathcal{Y}$.

With expectation (integration) in hand, we now consider various kinds of decompositions of measure.

2.3 Joint Measures

A joint probability space $(\mathcal{X} \times \mathcal{Y}, \mathbb{P}_{\mathcal{X} \times \mathcal{Y}})$ is a probability space whose base set is a product of (other) sets \mathcal{X}, \mathcal{Y} . In principle, you could imagine that each individually has its own measure $\mathbb{P}_{\mathcal{X}}, \mathbb{P}_{\mathcal{Y}}$, but do not need to suppose so apriori. Still: we obtain separate measure by marginalization (section 2.4); indeed, there are some relations among the joint measure $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ and $\mathbb{P}_{\mathcal{X}}, \mathbb{P}_{\mathcal{Y}}$. Before diving into what they are, keep in mind that a joint measure $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ is *not*, in general, constructed from measures $\mathbb{P}_{\mathcal{X}}, \mathbb{P}_{\mathcal{Y}}$. Therefore, you should think of the joint measure (simply and primarily) as a measure on the joint space; any connection to the marginal spaces is secondary.

2.4 Marginalization

We now turn to induced probability measures on the standard diagram

$$\begin{array}{ccc} \mathcal{X} \times \mathcal{Y} & \xrightarrow{\pi_{\mathcal{Y}}} & \mathcal{Y} \\ \downarrow \pi_{\mathcal{X}} & \searrow \tilde{\gamma} & \\ \mathcal{X} & & \end{array} \quad (6)$$

Let $(\mathcal{X} \times \mathcal{Y}, \mathbb{P}_{\mathcal{X} \times \mathcal{Y}})$ be a joint probability measure. The projection maps $\mathcal{X} \times \mathcal{Y} \xrightarrow{\pi_{\mathcal{X}}} \mathcal{X}$ and $\mathcal{X} \times \mathcal{Y} \xrightarrow{\pi_{\mathcal{Y}}} \mathcal{Y}$ induce probability measures on \mathcal{X} and \mathcal{Y} defined as:

$$\mathbb{P}_{\mathcal{X}}(A) := \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(\pi_{\mathcal{X}}^{-1}(A)) = \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(A \times \mathcal{Y}). \quad (7)$$

The second equality follows from the fact that $\pi_{\mathcal{X}}^{-1}(A) := \{(x, y) \in \mathcal{X} \times \mathcal{Y} : \pi_{\mathcal{X}}(x, y) \in A\}$. Of course, the marginalization for $\mathbb{P}_{\mathcal{Y}}$ is defined analogously.

Quite literally, marginalization is projection: it's a mechanism for putting a probability measure on the projected space assuming the existence of probability measure upstairs. To measure event $A \subset \mathcal{X}$ downstairs, you look at the tube $A \times \mathcal{Y}$ upstairs and measure it (fig. 7).




Figure 7: Geometry of Marginalization

2.5 Conditional Probability

For a probability space $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ you've likely seen the definition of conditional probability as $\mathbb{P}_{\mathcal{X}}(A|B) := \mathbb{P}_{\mathcal{X}}(A \cap B) / \mathbb{P}_{\mathcal{X}}(B)$ provided that the denominator is nonzero. It is easier to visualize this notion with joint probability. We continue with supposing that $(\mathcal{X} \times \mathcal{Y}, \mathbb{P}_{\mathcal{X} \times \mathcal{Y}})$ is a joint probability space. We've already defined marginal probability, so we can define the following conditional probability.

Definition 2.7. The conditional probability $\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(B|A)$ is defined to be

$$\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(B|A) := \frac{\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(A \times B)}{\mathbb{P}_{\mathcal{X}}(A)} \quad (8)$$

provided that the marginal $\mathbb{P}_{\mathcal{X}}(A) \neq 0$.

The proviso in this definition raises a question on what conditional probability means when $\mathbb{P}_{\mathcal{X}}(A) = 0$. We define it implicitly, with notation, $d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}$, that you will eventually become comfortable manipulating mechanically:

$$\int_A \int_{B|A} d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) d\mathbb{P}_{\mathcal{X}}(x) := \int_{A \times B} d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(x, y), \quad (9)$$

or for random variable $f : \mathcal{X} \times \mathcal{Y}$ we have

$$\int_{\mathcal{X}} \int_{\mathcal{Y}|\mathcal{X}} f(x, y) d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) d\mathbb{P}_{\mathcal{X}}(x) := \int_{\mathcal{X} \times \mathcal{Y}} f(x, y) d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(x, y). \quad (10)$$

Starting with (9), you should read this as: the inner integral $\int_{B|A} d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}$ defines a random variable on \mathcal{X} , of which we may evaluate expectation with respect to measure $\mathbb{P}_{\mathcal{X}}$. What is the random variable? Whatever it is, it makes the equality (9)! At the level of formalism, this implicit definition may not seem very illuminating, but momentarily we will interpret it geometrically in a manner reminiscent of concepts from multivariable calculus which you *are* familiar with.

You may recognize the original definition in eq. (8) as secretly appearing here, since $d\mathbb{P}_{\mathcal{Y}|\mathcal{X}} d\mathbb{P}_{\mathcal{X}} = d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ may be "solved" for $d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}$. But remember, in addition, that this 'dP' notation itself is defined as a package $\int_A d\mathbb{P} = \mathbb{P}(A)$.

Remark 2.7. There were many questions on interpreting the critter $A|B$. In pictures I draw the rectangle $A \times B$. It is fine to think of $A|B$ pictorially as $A \times B$, but you must concomitantly think of the

ambient space in which $A|B$ lives: $A \times B$, $A|B$, $B|A$ are all “the same” rectangle, but $A \times B \subset \mathcal{X} \times \mathcal{Y}$, $A|B \subset \mathcal{X}|B$ (which you may visualize as the rectangle $\mathcal{X} \times B$ which rectangle has probability one, *not* $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(\mathcal{X} \times B)$ which may be less than one). Similarly, $B|A$ is the rectangle $A \times B$, but instead of living in $\mathcal{X} \times \mathcal{Y}$ it lives in $\mathcal{Y}|A$ (or visually the rectangle $A \times \mathcal{Y}$ with unit probability).




Figure 8

In math, hand-waiving can sometimes be dangerous. On the other hand, it can sometimes allow us to think reasonably about, and operationalize our intuition of, notions whose formalism is “beyond the scope of this course,” all with the aim of performing computations and symbolic manipulations. If you are interested in more rigorously making sense of $d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}$, I recommend diving into the measure theory.⁵

The level of rigor we need is: we may fluidly manipulate integral expressions involving joint probability, relying on our multivariable intuition from calculus that decomposes high dimensional integration as sequence of one-dimensional integrals. What changes with probability, is that the (measure used for the) inner one-dimensional integral may depend on the outer value.

Now for the geometric intuition: in multivariable calculus, to integrate a function $f(x, y)$, $\int_{\mathcal{R}} f$, over some region $\mathcal{R} \subset \mathcal{X} \times \mathcal{Y}$ (imagine e.g. that $\mathcal{X} = \mathcal{Y} = \mathbb{R}$, so this function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$), we decompose the high-dimensional integral into two *one-dimensional* integrals $\int_{\mathcal{X}} \int_{\mathcal{Y}} f(x, y) dy dx$, i.e. we integrate the function $f(x, y)$ with respect to y , holding x fixed, then integrate the what is left with respect to x . That means we slice \mathbb{R}^2 at $\{x\} \times \mathbb{R}$, integrate in \mathbb{R} (y ’s copy of \mathbb{R}), which returns a *value*, then integrate those values again with respect to \mathbb{R} (x ’s copy of \mathbb{R}).

2.6 Independence

Forget, for now, our reliance on joint probability space $(\mathcal{X} \times \mathcal{Y}, \mathbb{P}_{\mathcal{X} \times \mathcal{Y}})$, and suppose that we have two separate probability spaces $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ and $(\mathcal{Y}, \mathbb{P}_{\mathcal{Y}})$. From these two spaces, one may reasonably ask if we can “go the other direction” and construct a (joint) probability measure $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ on $(\mathcal{X} \times \mathcal{Y})$. The answer is yes.

Before doing so, let us reason about the properties we would like this measure to have. The first thing we should expect is that projection $\pi_{\mathcal{X}} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathcal{X}$ *preserves measure*, i.e. that

$$\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(A \times \mathcal{Y}) = \mathbb{P}_{\mathcal{X}}(\pi_{\mathcal{X}}(A \times \mathcal{Y})) = \mathbb{P}_{\mathcal{X}}(A).$$

⁵See e.g. the [Radon-Nikodym Theorem](#).

Observe that in contrast to eq. (7), we do not define $\mathbb{P}_{\mathcal{X}}(A)$ in terms of $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$. Instead, the definition goes the other way:

$$\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(A \times \mathcal{Y}) := \mathbb{P}_{\mathcal{X}}(A). \quad (11)$$

Of course, we would like the analogous equality to hold for $\mathcal{X} \times B$ with $\mathbb{P}_{\mathcal{Y}}(B)$.

Now this condition by itself only allows us to define sets of the form $A \times \mathcal{Y}$ or $\mathcal{X} \times B$. For general rectangle $A \times B \subset \mathcal{X} \times \mathcal{Y}$, define

$$\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(A \times B) := \mathbb{P}_{\mathcal{X}}(A) \cdot \mathbb{P}_{\mathcal{Y}}(B). \quad (12)$$

Observe, in connection with one's first typical encounter with independence as $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$, we can recover this relation by observing that

$$A \times B = (A \times \mathcal{Y}) \cap (\mathcal{X} \times B), \quad (13)$$

and compute

$$\begin{aligned} \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(A \times B) &= \mathbb{P}_{\mathcal{X}}(A)\mathbb{P}_{\mathcal{Y}}(B) \\ &= \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(A \times \mathcal{Y}) \cdot \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(\mathcal{X} \times B). \end{aligned} \quad (14)$$

The intuition we extract from this construction is that independence is a most natural way to construct measure on high dimensional space using measure from its lower dimensional components. In exactly the same way that we construct area from length or volume from area and length, or volume from length, etc. Use your geometric intuition liberally!

Definition 2.8. In general, we say that $(\mathcal{X}^m, \mathbb{P}_{\mathcal{X}^m})$ is *independent* if

$$\mathbb{P}_{\mathcal{X}^m} = (\mathbb{P}_{\mathcal{X}})^m$$

or more generally that $\left(\prod_{j=1}^m \mathcal{X}_j, \mathbb{P}_{\prod \mathcal{X}_j} \right)$ independent if

$$\mathbb{P}_{\prod \mathcal{X}_j} = \prod_{j=1}^m \mathbb{P}_{\mathcal{X}_j}. \quad (15)$$

Remark 2.8. Be careful with this definition: from spaces $(\mathcal{X}_j, \mathbb{P}_{\mathcal{X}_j})$, the independent measure $\mathbb{P}_{\prod \mathcal{X}_j}$ in (15) is not the only possible measure on joint space $\prod_j \mathcal{X}_j$!

2.7 Data

We will discuss the connection between data and probability more later, but for now assert an easy way to interpret marginalization and conditioning in terms of data. Imagine a spreadsheet with n columns. Data in the first column has three unique values, 'a,' 'b,' and 'c.' Marginalization, in terms of data, simply corresponds to deleting columns, e.g. to obtain the marginal for column 1, we delete (or ignore) columns 2- n . Conditioning, by contrast, corresponds to deleting rows. Conditioned on column 1 data equaling (say) 'a' amounts to looking only at those rows (all columns 2- n included) at which the column 1 entry is 'a.'

3 Lecture 3

We continue discussing independence as a high-dimensional phenomenon, and introduce concentration.

We started with a quick review of conditional probability from last time, continuing geometric interpretation for $A \times B$, $A|B$ and $B|A$ as “sets” in $\mathcal{X} \times \mathcal{Y}$ (remark 2.7).

3.1 Independence

Recall definition 2.8 which says that a joint probability measure $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ is independent if it decomposes as a product of marginals $\mathbb{P}_{\mathcal{X}} \cdot \mathbb{P}_{\mathcal{Y}}$. This extends to products of arbitrarily many factors.

To recognize independence, either of the following suffice:

1. conditional probability is independent of condition, or (what amounts to the same)
2. conditional probability is equal to the marginal

Ultimately, our purpose in going through this rigmarole is for computation. I would like you to feel natural doing symbolic manipulations e.g. of the form $\int_{\mathcal{X} \times \mathcal{Y}} = \int_{\mathcal{X}} \int_{\mathcal{Y}|\mathcal{X}}$. You might see this in other contexts as the “tower” property. If you are already comfortable with iterated and embedded expectations, then you need not refer to the geometric interpretation. Use whichever viewpoint you are most at home with when doing computations with joint probability spaces.

Example 3.1. We start with $(\mathcal{X} \times \mathcal{Y} = \mathbb{R}^k \times \{0, 1\}, \mathbb{P}_{\mathcal{X} \times \mathcal{Y}})$ independent, and suppose that we have “optimal” model $\tilde{y} : \mathcal{X} \rightarrow \mathcal{Y}$. We revisit through a computation what optimality means. Computing accuracy, which is the measure of event $\{\tilde{y} = y\} = \{(x, y) \in \mathcal{X} \times \mathcal{Y} : \tilde{y}(x) = y\}$, we have:

$$\begin{aligned}
 \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(\tilde{y} = y) &= \mathbb{E}(\mathbb{1}_{\tilde{y}(x)=y}) \\
 &:= \int_{\mathcal{X} \times \mathcal{Y}} \mathbb{1}_{\tilde{y}(x)=y} d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(x, y) \\
 &= \int_{\mathcal{X}} \int_{\mathcal{Y}|\mathcal{X}} \mathbb{1}_{\tilde{y}(x)=y} d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) d\mathbb{P}_{\mathcal{X}}(x) \\
 &= \int_{\mathcal{X}} \int_{\mathcal{Y}} \mathbb{1}_{\tilde{y}(x)=y} d\mathbb{P}_{\mathcal{Y}}(y) d\mathbb{P}_{\mathcal{X}}(x) \\
 &= \int_{\mathcal{X}} \int_{\mathcal{Y}} \mathbb{1}_{\tilde{y}(x)=0}(1-p) + \mathbb{1}_{\tilde{y}(x)=1}p d\mathbb{P}_{\mathcal{X}}(x) \\
 &= \int_{\mathcal{X}} \max\{p, 1-p\} d\mathbb{P}_{\mathcal{X}}(x) \\
 &= \max\{p, 1-p\} \int_{\mathcal{X}} d\mathbb{P}_{\mathcal{X}}(x) \\
 &= \max\{p, 1-p\}.
 \end{aligned} \tag{16}$$

In the first equation, we recall that probability may be written as expectation of an indicator. This is useful!, it allows us to do this computation. The second line is the definition of expectation (definition 2.2). The third line is our implicit definition of conditional probability eq. (10), the geometry of which, reminder, you should think of as iterated integration in multivariable calculus. The fourth line is independence (condition #2 above). The fifth is expectation w.r.t. $\mathbb{P}_{\mathcal{Y}}$ of $\mathbb{1}_{\tilde{y}(x)=y}$. The sixth line is optimality of \tilde{y} ; we are trying to optimize accuracy, and we’ve written it such that we may optimize *pointwise* (in x). In the seventh line, we pull out $\max\{p, 1-p\}$ since p is independent of x and in the final line we recall that $\mathbb{P}_{\mathcal{X}}(\mathcal{X}) = 1$.

Next we interpret independence for data.

3.2 Data

Referring to the standard diagram (1), we hope and expect that most instances where supervised machine learning comes in to play, the measure $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ on $\mathcal{X} \times \mathcal{Y}$ is not independent. Sometimes, as in the 0th programming assignment, you'll run into an evil data set, but such are not the norm. Instead, independence is a phenomenon we'd like to associate with *data*.

Definition 3.1. We say that a sequence of (labeled) points $\mathcal{S} = ((x_1, y_1), \dots, (x_m, y_m)) \sim_{\text{iid}} \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ is (sampled) *independent and identically distributed* if $\mathcal{S} \in (\mathcal{X} \times \mathcal{Y})^m$ is a point in joint probability space $(\mathcal{X} \times \mathcal{Y})^m$ with independent measure $\mathbb{P}_{(\mathcal{X} \times \mathcal{Y})^m} = (\mathbb{P}_{\mathcal{X} \times \mathcal{Y}})^m$.

The following picture is for the intuition in the high-dimensional space.




Figure 9

Remark 3.1. While seemingly pedantic, the nuance is important: a data set in ML is properly speaking a sequence. While we might not care about order, we absolutely do care about repetition.

Talk of sampling points suggests talk of randomness. You are allowed to select any point $x \in \mathcal{X}$ that you fancy. How you sample should in some sense be related to your measure $\mathbb{P}_{\mathcal{X}}$. One way of handwaiving this relation is to say that given any set (event) $A \subset \mathcal{X}$, the probability that the point $x \in \mathcal{X}$ you picked happens to (also) be in A is $\mathbb{P}_{\mathcal{X}}(A)$. In other words, we are saying in English that $\mathbb{E}(\mathbb{1}_{x \in A}) = \mathbb{P}_{\mathcal{X}}(A)$. Since we generally do not evaluate $\mathbb{P}_{\mathcal{X}}$ directly, we need a more data-centric way to talk about probability, one which ties with our understanding of “randomness,” which up to this point we’ve barely mentioned! The connection arises (primarily) from the Law of Large Numbers. First a quick review of concentration bounds.

3.3 Concentration Inequalities

Concentration inequalities are sort of the heartbeat (for guarantees) in ML. They are omnipresent and make many important results work. We enumerate a few here and will return to them again later in the semester when we discuss *probably approximately correct* (PAC) learnability.

Proposition 3.1 (Markov). Suppose that $(\mathcal{X} \subset \mathbb{R}^{\geq 0}, \mathbb{P}_{\mathcal{X}})$ a non-negative probability space. Then

$$\mathbb{P}_{\mathcal{X}}(x \geq t) \leq \mathbb{E}(x)/t.$$

Proof. Compute: $\mathbb{E}(x) := \int_{\mathcal{X}} x d\mathbb{P}_{\mathcal{X}}(x) = \int_{[0,t)} x d\mathbb{P}_{\mathcal{X}}(x) + \int_{[t,\infty)} x d\mathbb{P}_{\mathcal{X}}(x)$, the equality by linearity of integration. Since $x \geq 0$, the second term bounds

$$\int_{[t,\infty)} x d\mathbb{P}_{\mathcal{X}}(x) \geq \int_{[t,\infty)} t d\mathbb{P}_{\mathcal{X}}(x),$$

which follows from the fact that $x \geq t$ on $x \in [t, \infty)$. The first integral $\int_{[0,t)} x d\mathbb{P}_{\mathcal{X}}(x) \geq 0$, so we

have $\mathbb{E}(x) \geq \int_{[t,\infty)} t d\mathbb{P}_{\mathcal{X}}(x) = t \cdot \int_{[t,\infty)} d\mathbb{P}_{\mathcal{X}}(x)$. Interpreting the integral as probability and solving proves the result. \square

Proposition 3.2 (Chebyshev). Let $(\mathcal{X} \subset \mathbb{R}, \mathbb{P}_{\mathcal{X}})$ be a probability space with mean $\mu_{\mathcal{X}}$ and finite variance $\sigma_{\mathcal{X}}^2$. Then for $\varepsilon > 0$,

$$\mathbb{P}_{\mathcal{X}}(|x - \mu_{\mathcal{X}}| > \varepsilon) \leq \sigma_{\mathcal{X}}^2 / \varepsilon^2.$$

Proof. Apply proposition 3.1 to $\{|x - \mu_{\mathcal{X}}| > t\} = \{(x - \mu_{\mathcal{X}})^2 > t^2\}$ □

The next two are more advanced; we introduce them here so that when we see them later in the semester it will be for a second time.

Proposition 3.3 (Hoeffding). Let $(\mathcal{X} \subset [0, 1], \mathbb{P}_{\mathcal{X}})$ be a probability space with mean $\mathbb{E}(x) = \mu_{\mathcal{X}}$. Then

$$\mathbb{P}_{\mathcal{X}^m} \left(\left| \frac{1}{m} \sum_{j=1}^m x_j - \mu_{\mathcal{X}} \right| > \epsilon \right) \leq 2e^{-2m\epsilon^2}. \quad (17)$$

The Law of Large Numbers (LLN), which we recall in section 3.3.1, provides a guarantee that sample mean is close to expectation, in the limit. There are various ways of cashing this out (Strong Law, Weak Law, etc.), but Hoeffding is not an asymptotic result: eq. (17) holds for *all* $m \in \mathbb{N}$. Of course, for cases where $\mathcal{X} \subset [0, 1]$ (or more generally, is bounded) Hoeffding readily proves LLN by taking the limit $m \rightarrow \infty$.

Finally, we present Glivenko-Cantelli, a result which guarantees that an empirical cumulative distribution function (ECDF) is close to the CDF.

Theorem 3.1 (Glivenko-Cantelli). Let $(\mathcal{X} = \mathbb{R}, \mathbb{P}_{\mathcal{X}})$ be a probability space, for $t \in \mathbb{R}$, define $F(t) := \mathbb{P}_{\mathcal{X}}(x \leq t)$ the cdf and $F_m(t) : \mathcal{X}^m \rightarrow \mathbb{R}$ by

$$(x_1, \dots, x_m) \mapsto \frac{1}{m} \sum_{j=1}^m \mathbb{1}_{x_j \leq t}$$

the *empirical cdf*. Then

$$\mathbb{P}_{\mathcal{X}^m} \left(\sup_{t \in \mathbb{R}} |F_m(t) - F(t)| > \epsilon \right) \leq 8(m+1)e^{-\frac{m \cdot \epsilon^2}{32}}. \quad (18)$$

The statement of this theorem is phenomenal: no matter what $\mathbb{P}_{\mathcal{X}}$ is, you can articulate precise conditions for satisfaction of a precision specification (ϵ) with arbitrarily high confidence $(1 - \delta)$ provided m is “sufficiently large.” And again... entirely independent of $\mathbb{P}_{\mathcal{X}}$. Sometimes, one must pause to marvel at the beauty of mathematics.

There is a stronger form of this result which replaces the right hand side of (18) with the right hand side of (17).⁶

3.3.1 Law of Large Numbers (LLN)

The version of LLN we provide is the *Weak Law*, which is sufficient for our purposes. The Strong Law denotes a different mode of convergence (‘almost surely’ as opposed to the Weak Law’s ‘in probability’).

Theorem 3.2. Let $(\mathcal{X}, \mathbb{P}_{\mathcal{X}})$ be a probability space and for $m \in \mathbb{N}$, $(\mathcal{X}^m, \mathbb{P}_{\mathcal{X}^m})$ independent (meaning: $\mathbb{P}_{\mathcal{X}^m} = \mathbb{P}_{\mathcal{X}}^m$), and suppose that both mean $\mu_{\mathcal{X}} := \mathbb{E}(x) = \int_{\mathcal{X}} x d\mathbb{P}_{\mathcal{X}}(x)$ and variance $\sigma_{\mathcal{X}}^2 := \mathbb{E}((x - \mu_{\mathcal{X}})^2)$ are finite. Define empirical mean $s_m : \mathcal{X}^m \rightarrow \mathbb{R}$ as a random variable by $(x_1, \dots, x_m) \mapsto \frac{1}{m}(x_1 + \dots + x_m)$. Then for any $\varepsilon > 0$,

$$\lim_{m \rightarrow \infty} \mathbb{P}_{\mathcal{X}^m}(|s_m - \mu_{\mathcal{X}}| > \varepsilon) = 0.$$

⁶See the DKW(M) inequality https://en.wikipedia.org/wiki/Dvoretzky–Kiefer–Wolfowitz_inequality. Massart (the ‘M’) was responsible for the tight coefficient of 2.

Proof. Without loss of generality suppose that $\mu_{\mathcal{X}} = 0$.⁷ By Chebyshev's inequality (proposition 3.2),

$$\begin{aligned} \mathbb{P}_{\mathcal{X}^m}(|s_m - \mu_{\mathcal{X}}| > \varepsilon) &= \mathbb{P}_{\mathcal{X}^m}((x_1 + \dots + x_m)^2 > m^2 \varepsilon^2) \\ &= \mathbb{E} \left[\left(\sum_{i=1}^m x_i \right) \left(\sum_{j=1}^m x_j \right) \right] \\ &\leq \frac{\mathbb{E} \left[\left(\sum_{i=1}^m x_i \right) \left(\sum_{j=1}^m x_j \right) \right]}{m^2 \varepsilon^2} \\ &= \frac{\sum_{i=1}^m \mathbb{E}(x_i^2) + \sum_{i \neq j} \mathbb{E}(x_i) \mathbb{E}(x_j)}{m^2 \varepsilon^2} \\ &= \frac{m \sigma_{\mathcal{X}}^2}{m^2 \varepsilon^2} \xrightarrow{m \rightarrow \infty} 0, \end{aligned}$$

where we use independence and mean-zero to eliminate $\mathbb{E}(x_i x_j) = 0$ for $i \neq j$. \square

Remark 3.2. The statement of LLN measures the set of point $(x_1, \dots, x_m) \in \mathcal{X}^m$ that are at least ε -far from sample mean $s_m^{-1}(\mu_{\mathcal{X}})$.

With LLN under our belt, we may finally interpret randomness into probability: to say that $(x_1, \dots, x_n) \sim_{\text{iid}} \mathbb{P}_{\mathcal{X}}$ (at the moment, we're not talking about labeled data) means, among other things, that we can expect the Law of Large Numbers to hold for probabilities. That is, for any event $A \subset \mathcal{X}$, consider random variable $\mathbb{1}_{x \in A} := \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{else.} \end{cases}$ Then

$$\frac{1}{m} \sum_{j=1}^m \mathbb{1}_{x_j \in A} \xrightarrow{m \rightarrow \infty} \mathbb{E}(\mathbb{1}_{x \in A}) = \mathbb{P}_{\mathcal{X}}(A).$$

As the convergence in this expression is "in probability," we first fix $\varepsilon > 0$; then for any $\delta > 0$, there is $M_\delta > 0$ so that

$$\mathbb{P}_{\mathcal{X}^m} \left(\left| \frac{1}{m} \sum_{j=1}^m \mathbb{1}_{x_j \in A} - \mathbb{P}_{\mathcal{X}}(A) \right| > \varepsilon \right) < \delta \text{ whenever } m > M_\delta. \quad (19)$$

There is a procedure to mechanize this result. Suppose you've set $\varepsilon, \delta > 0$ and (happen to) know M_δ satisfying (19). Sample $x_1, \dots, x_m \sim_{\text{iid}} \mathbb{P}_{\mathcal{X}}$ with $m > M_\delta$ and evaluate the random variable $\mathbb{1}_{|s_m(x_1, \dots, x_m) - \mathbb{P}_{\mathcal{X}}(A)| > \varepsilon}$. The answer will be zero or one. Most of the time it should be zero. If you do this many times,⁸ and take the empirical average of *these* results, the answer should be less than δ .

3.4 Monkeys on a Keyboard: Intuition in High Dimension

Probability does weird things in high dimension. I made a tenuous sounding claim that a high dimensional gaussian has density concentrated at the origin (tracks our low-dimensional intuition) but probability concentrated *away* from it (what?!).

Let's say this a bit more formally. Suppose that $(\mathcal{X} = \mathbb{R}, \mathbb{P}_{\mathcal{X}})$ is a random variable with $\mathbb{P}_{\mathcal{X}}$ normal, zero mean, unit variance, and $(\mathcal{X}^m, \mathbb{P}_{\mathcal{X}^m})$ independent. For $\varepsilon > 0$,

$$\lim_{m \rightarrow \infty} \mathbb{P}_{\mathcal{X}^m} \left(\left| \|x\|^2 - m \right| > \varepsilon \right) = \lim_{m \rightarrow \infty} \mathbb{P}_{\mathcal{X}^m} \left(\left\{ x \in \mathcal{X}^m : \left| \|x\|^2 - m \right| > \varepsilon \right\} \right) = 0.$$

This says that in high dimension, a gaussian concentrates around the sphere of radius \sqrt{m} . In particular, the *solid* sphere of radius (strictly) less than \sqrt{m} is practically empty (despite density being greatest inside)!

We run a computation to more concretely illustrate the point.

⁷Think about why applying the transformation $x \mapsto x - \mu_{\mathcal{X}}$, inducing mean zero random variable, does not affect the LLN result.

⁸As $\mathbb{1}_{|s_m(x_1, \dots, x_m) - \mathbb{P}_{\mathcal{X}}(A)| > \varepsilon}$ is a $[0, 1]$ bounded random variable, proposition 3.3 applies and you can in fact identify what 'many' needs to be to satisfy the subsequent 'should be.'

Example 3.2. Consider random variable $(\mathcal{X} = [0, 1], \mathbb{P}_{\mathcal{X}})$ with uniform measure $\mathbb{P}_{\mathcal{X}}([a, b]) = (b - a)\mathbb{1}_{0 \leq a \leq b \leq 1}$. Suppose that $A \subset \mathcal{X}$ with $1 > \mathbb{P}_{\mathcal{X}}(A) \geq 1 - \varepsilon$ has high probability. The “hypercube” $A^m \subset \mathcal{X}^m$ in high dimension has measure

$$\mathbb{P}_{\mathcal{X}^m}(A^m) = (\mathbb{P}_{\mathcal{X}}(A))^m = (1 - \varepsilon)^m,$$

where the first equality follows by independence. As long as $1 > \varepsilon > 0$, $\lim_{m \rightarrow \infty} (1 - \varepsilon)^m = 0$.

As a practical example, imagine tasking some monkey to button mash on a keyboard in perpetuity. Assuming independence and identical distribution of key strokes, the probability of hitting any character (or punctuation) is roughly $1/30$, and therefore the probability of matching a length ℓ sequence of characters is $1/30^\ell$ (independence!). The complete works of Shakespeare have, let’s estimate, roughly 4,000,000 characters. Thus, the probability that a monkey fails to type out the complete works of Shakespeare, in one iteration of 4,000,000 key strokes, is about $1 - (1/30)^{4000000}$ (fairly close to 1). In one go, it will likely not type out the complete works. But if it keeps trying, eventually it will.

4 Lecture 4

4.1 Setting: Detection / Binary Classification

We are ready to discuss a first instance of machine learning problem: binary classification, where $\mathcal{Y} = \{0, 1\}$. Recalling (1), the task is to construct *model* $\tilde{y} : \mathcal{X} \rightarrow \mathcal{Y}$ so that $\tilde{y}(x) \approx y$ for $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ -most $(x, y) \in \mathcal{X} \times \mathcal{Y}$. Alternatively, we describe this goal in functional form as: $\tilde{y} \circ \pi_{\mathcal{X}} \approx \pi_{\mathcal{Y}}$ in expectation. Pretty much all supervised learning takes this form, and all that we've done at this point is impose a constraint on codomain \mathcal{Y} , namely that it is a discrete set consisting of two elements.

The supervised learning problem for binary classification is typically solved in two steps:

1. construct *score* function (which here we'll denote by) $\tilde{y} : \mathcal{X} \rightarrow \mathbb{R}$ (or to closed interval $[0, 1]$), and
2. *threshold* score to obtain discrete output $\hat{y}_t : \mathcal{X} \rightarrow \mathcal{Y}$ defined by $\hat{y}_t(x) := \mathbb{1}_{\tilde{y}(x) \geq t}$.

We'll return more extensively to step 1. soon enough. For now, suppose that we have a score function \tilde{y} and correspondingly, on labeled data set $S = \{(x_1, y_1), \dots, (x_m, y_m)\}$, a set of scores

$$\{(\tilde{y}(x_1), y_1), \dots, (\tilde{y}(x_m), y_m)\}.$$

4.2 Interpreting the Score

The first point to make is that we would like the scores to separate classes. Namely, if we group scores by class $S_j := \{\tilde{y}(x_k) : y_k = j\}$ for $j = 0, 1$, you should expect a histogram of each to correspondingly separate:




Figure 10: Conditional Histograms

Indeed, the score should serve as some sort of indicator of class! (Here we recall the asymmetry between sampling from “input” space \mathcal{X} and from \mathcal{Y} : the score is evaluable as a function only of input.) The more these two histograms separate, in principle, the better the score. Of course, the histogram represents data we already have, and what you'd like to know is that a histogram of scores on new data looks similar. In probability speak, the normalized histograms approximate *conditional densities* f_j , for $j = 0, 1$, according to which

$$\int_{[a,b]} f_j(s) ds = \mathbb{P}_{\mathcal{X}|\mathcal{Y}}([a,b] | y = j).$$

In a perfect world, both 1. the densities would well-approximate the histograms and 2. the densities would be strongly separated. A measure of separation is given e.g. by the *Kolmogorov-Smirnov (KS) score*, namely:

$$ks := \sup_{t \in \mathbb{R}} |\mathbb{P}_{\mathcal{X}|\mathcal{Y}}(s \leq t | y = 0) - \mathbb{P}_{\mathcal{X}|\mathcal{Y}}(s \leq t | y = 1)|.$$

Understand geometrically why $ks \approx 1$ corresponds to (near) total separation while $ks \approx 0$ corresponds to (near) total overlap.

Example Consider a typical use-case where a bank must decide whether to extend a loan to an applicant based on information they provide at the time of application. With $\mathcal{Y} = \{0, 1\}$, let $y = 0$ correspond to ‘application denied’ and $y = 1$ to ‘loan approved.’ Let $\mathcal{X} = \mathbb{R}^k$ and suppose that each component denotes some measurable attribute, e.g. x^0 debt to income ratio, x^1 income, x^2 age, x^3 years of employment and so on.⁹¹⁰ You can imagine that the projection $\pi_0 : \mathcal{X} \rightarrow \mathbb{R}$ returning debt-to-income (dti) would provide a fairly decent predictor of whether a loan will (should) be extended. In this case, technically our label denotes something more like an answer to the question ‘will they repay their loan?’

Consider a significantly harder example, where input data in $\mathcal{X} = \mathbb{R}^{3 \times k \times k}$ are images and $y = 1$ corresponds to ‘Waldo is in the image.’ In *this* case, π_0 returning the red value of the first pixel will unlikely be a particularly good score.

4.2.1 Optimal Threshold

Now, the score function $\tilde{y} : \mathcal{X} \rightarrow \mathbb{R}$ induces a measure $\mathbb{P}_{\mathbb{R} \times \mathcal{Y}}$ on $\mathbb{R} \times \{0, 1\}$ by $\mathbb{P}_{\mathbb{R} \times \mathcal{Y}}([a, b] \times \{j\}) := \mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(\tilde{y}^{-1}([a, b]) \times \{j\})$. We’ve seen induces measures, though it’s worth recalling them in this context, which is slightly more motivated to a problem at hand. While the measure in simpler space $\mathbb{R} \times \mathcal{Y}$ is *defined* in terms of the one in more complex space $\mathcal{X} \times \mathcal{Y}$, we will compute in the simpler space, in the hopes that class separation is easier to see there. If this sounds counterintuitive keep in mind that this is half the whole point of using random variables (as maps to \mathbb{R}).

Given such a distribution of scores, there remains the question how to predict?, i.e. how to associate output label y to given input point x with score $\tilde{y}(x)$. We feel that scores on “one side” should belong to one label, and indeed, thresholding the score provides half an answer: let label predictor $\hat{y}_t := \mathbb{1}_{\tilde{y}(x) \geq t}$ segment the space: to the left of t belongs one label, and to the right the other. Still, what should t be? To answer this question we must first identify what our objective is. One reasonable objective is to maximize accuracy (or minimize error) $\hat{y}_t(x) = y$ over data $(x, y) \in \mathcal{X} \times \mathcal{Y}$. It isn’t quite right to say this should hold *for all* (x, y) —in particular, because there may be pairs $(x, y), (x, y')$ with $y \neq y'$ —on which account we merely ask for equality to hold for *most* (x, y) .¹¹ Measuring ‘most’ requires a measure ($\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$), and so we ask for approximate equality for ‘most in probability’ or what amounts to the same, in expectation for suitably defined random variable ($\mathbb{E}(\mathbb{1}_{\hat{y}_t(x)=y})$). Maximizing expected accuracy is equivalent to minimizing expected error, and for the sake of uniformity with future convention, we’ll stick to minimizing error.

We compute: for fixed $t \in \mathbb{R}$,

$$\begin{aligned} \mathbb{E}(\mathbb{1}_{\hat{y}_t(x) \neq y}) &= \int_{\mathcal{X} \times \mathcal{Y}} \mathbb{1}_{\hat{y}_t(x) \neq y} d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(x, y) \\ &= \int_{\mathcal{Y}} \int_{\mathcal{X}|\mathcal{Y}} \mathbb{1}_{\hat{y}_t(x) \neq y} d\mathbb{P}_{\mathcal{X}|\mathcal{Y}}(x|y) d\mathbb{P}_{\mathcal{Y}}(y) \\ &= \sum_{y \in \mathcal{Y}} \int_{\mathcal{X}|\mathcal{Y}} \mathbb{1}_{\hat{y}_t(x) \neq y} d\mathbb{P}_{\mathcal{X}|\mathcal{Y}}(x|y) \mathbb{P}_{\mathcal{Y}}(y) \\ &= \mathbb{P}_{\mathcal{Y}}(y = 0) \int_t^\infty f_0(s) ds + \mathbb{P}_{\mathcal{Y}}(y = 1) \int_{-\infty}^t f_1(s) ds. \end{aligned} \tag{20}$$

In the last line, we recall that $\mathbb{1}_{\hat{y}_t(x) \neq y} = 1$ means that $\hat{y}_t(x) \neq y$, which happens when either $\tilde{y}(x) \geq t$ and $y = 0$ or $\tilde{y}(x) \leq t$ and $y = 1$. It’s at this step we sub in the measure $\mathbb{P}_{\mathbb{R} \times \mathcal{Y}}$, whose conditional w.r.t. y is given by integrating the corresponding density f_j ; again this is one reason for working with scores: you can integrate in \mathbb{R} perhaps a lot easier than you can in some wonky space \mathcal{X} .

Let’s interpret limiting cases. When $t \rightarrow -\infty$, $\int_{\mathbb{R}} f_0 = 1$ and the error becomes $\mathbb{P}_{\mathcal{Y}}(y = 0)$. Similarly, when $t \rightarrow \infty$, we have $\int_{\mathbb{R}} f_1 = 1$ and get $\mathbb{P}_{\mathcal{Y}}(y = 1)$. (What does this mean in terms of what

⁹Superscript denotes component, not an exponent, which we use instead of subscript because that typically indexes the data sample.

¹⁰Note that while available data *may* in principle be usable for decisioning, there may be countervailing considerations prohibiting its use, regulation around [consumer finance](#) providing an excellent case in point.

¹¹Think of concrete examples. Two applicants for a loan may have the same debt-to-income ratio and yet different outcomes of repayment. In other words, the input-output relation between input data (dti) and output label (repayment) is not functional. That might be a problem if we didn’t have probability!

our model predicts?)

To find optimal t^* minimizing (20), take the derivative (w.r.t. t) and set to zero. By the Fundamental Theorem of Calculus, we obtain

$$t^* = \arg_{t \in \mathbb{R}} -\mathbb{P}_{\mathcal{Y}}(y=0)f_0(t) + \mathbb{P}_{\mathcal{Y}}(y=1)f_1(t) = 0.$$

When $\mathbb{P}(y=0) = \mathbb{P}(y=1)$ (both labels are equally likely), a candidate for the optimal point occurs when both conditional densities are equal. Considering fig. 10, does this make sense?

4.3 Constructing the Score

The procedure we outlined for finding t^* to threshold a score (and obtain hard predictions) is a microcosm of the same procedure we employ for *constructing* score $\tilde{y} : \mathcal{X} \rightarrow \mathbb{R}$. In that case, we looked for the minimum $\min_{t \in \mathbb{R}} \mathbb{E}(\mathbb{1}_{\tilde{y}_t \neq y})$ of an expectation for a (one-parameter) parameterized random variable. In this case, we similarly execute optimization $\min_{\tilde{y} : \mathcal{X} \rightarrow \mathbb{R}} \mathbb{E}(\ell_{\tilde{y}})$ for a random variable $\ell_{\tilde{y}} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}$, called a *loss* function, parameterized (this time) by model \tilde{y} . Two things to note: first, we have yet to specify what the loss function is (there will be many options). Secondly, our search is no longer over \mathbb{R} but instead a function space, namely the collection of maps $\mathcal{X} \rightarrow \mathbb{R}$. This space is much larger, and therefore the problem signif harder!

Still, let's take a stab at solving it. Suppose that you want score to reflect (something like) likelihood, so that higher scores (near 1) correspond to $y = 1$ and lower scores (near 0) correspond to $y = 0$. Then a perfectly fine loss function may be $\ell_{\tilde{y}}(x, y) := (\tilde{y}(x) - y)^2$.¹² To find optimal (call it) $y^* : \mathcal{X} \rightarrow \mathbb{R}$, we compute

$$\min_{\tilde{y} : \mathcal{X} \rightarrow \mathbb{R}} \mathbb{E}(\ell_{\tilde{y}}) = \int_{\mathcal{X} \times \mathcal{Y}} \ell_{\tilde{y}}(x, y) d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(x, y)$$

and this time condition the other way to get

$$\int_{\mathcal{X} \times \mathcal{Y}} (\tilde{y}(x) - y)^2 d\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}(x, y) = \int_{\mathcal{X}} \int_{\mathcal{Y}|\mathcal{X}} (\tilde{y}(x) - y)^2 d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) d\mathbb{P}_{\mathcal{X}}(x).$$

Minimizing the inner integral $\int_{\mathcal{Y}|\mathcal{X}} (\tilde{y}(x) - y)^2 d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x)$ *pointwise in x* will minimize the entire expectation. Therefore, we fix x and consider the optimization problem:

$$\min_{\tilde{y}(x) \in \mathbb{R}} \int_{\mathcal{Y}|\mathcal{X}} (\tilde{y}(x) - y)^2 \mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x).$$

The advantage of *this* expression is that we've once again reduced the problem to an optimization in \mathbb{R} (and swept under the rug whether a map $\mathcal{X} \rightarrow \mathbb{R}$ defined by $y^*(x) = \arg \min_{v \in \mathbb{R}} \int_{\mathcal{Y}|\mathcal{X}} (v - y)^2 d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x)$ is legit, i.e. measurable). Computing the conditional expectation by expanding the integrand, we obtain

$$\int_{\mathcal{Y}|\mathcal{X}} \tilde{y}(x)^2 - 2\tilde{y}(x)y + y^2 d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) = \tilde{y}(x)^2 \int_{\mathcal{Y}|\mathcal{X}} d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) - 2\tilde{y}(x) \int_{\mathcal{Y}|\mathcal{X}} y d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x) + \int_{\mathcal{Y}|\mathcal{X}} y^2 d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x).$$

Recall that the conditional integral integrates w.r.t. the "free variable" y , not conditioned one x ; that allows e.g. pulling $\tilde{y}(x)$ outside the integral. We rinse and repeat steps from thresholding, take the derivative of the right hand side with respect to $\tilde{y}(x)$ ¹³ and set to zero to obtain

$$2\tilde{y}(x) \underbrace{\int_{\mathcal{Y}|\mathcal{X}} d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x)}_1 - 2 \underbrace{\int_{\mathcal{Y}|\mathcal{X}} y d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x)}_{\mathbb{E}(y|x)} = 0,$$

which implies that $y^*(x) = \mathbb{E}(y|x)$. For this chosen loss function, the task is totally solved, and if such were the case we'd basically be done with ML, we could go home and move on to bigger and harder problems. There's one catch: we typically do not know the measure \mathbb{P} (either $\mathbb{P}_{\mathcal{X} \times \mathcal{Y}}$ or $\mathbb{P}_{\mathcal{Y}|\mathcal{X}}$)!

¹²Notice when $\tilde{y}(x) \in \{0, 1\}$ this loss function is identical to the one we used for thresholding!

¹³With x fixed, we're looking for the value $v \in \mathbb{R}$ to which we'll assign $\tilde{y}(x)$. So if you prefer, expand $\int_{\mathcal{Y}|\mathcal{X}} (v - y)^2 d\mathbb{P}_{\mathcal{Y}|\mathcal{X}}(y|x)$, take derivative w.r.t. v , and set to zero.

4.4 Metrics

(TODO: complete this section) When we have score-induced measure $\mathbb{P}_{\mathbb{R} \times \mathcal{Y}}$ we may write

$$\mathbb{P}_{\mathbb{R} \times \mathcal{Y}}([a, b] \times \{j\}) = \alpha_j \int_a^b f_j(s) ds.$$

Apparently $\alpha_j = \mathbb{P}_{\mathcal{Y}}(y = j)$ and $\int_a^b f_j(s) ds = \mathbb{P}_{\mathcal{X}|\mathcal{Y}}(x|y)$.