

The GAPic Package

Lukas Schnelle

GAPDays Spring 2024

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

We call (\prec, X_0, X_1, X_2) *triangular complex* if

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

We call (\prec, X_0, X_1, X_2) *triangular complex* if

- (i) The relation \prec is transitive, i.e. $\forall v \in X_0, e \in X_1, f \in X_2$:

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f$$

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

We call (\prec, X_0, X_1, X_2) *triangular complex* if

- (i) The relation \prec is transitive, i.e. $\forall v \in X_0, e \in X_1, f \in X_2$:

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f$$

- (ii) $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$ such that $v_1, v_2 \prec e$

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

We call (\prec, X_0, X_1, X_2) *triangular complex* if

- (i) The relation \prec is transitive, i.e. $\forall v \in X_0, e \in X_1, f \in X_2$:

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f$$

- (ii) $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$ such that $v_1, v_2 \prec e$
(iii) $\forall e \in X_1 \exists f \in X_2$ such that $e \prec f$

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

We call (\prec, X_0, X_1, X_2) *triangular complex* if

- (i) The relation \prec is transitive, i.e. $\forall v \in X_0, e \in X_1, f \in X_2$:

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f$$

- (ii) $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$ such that $v_1, v_2 \prec e$
(iii) $\forall e \in X_1 \exists f \in X_2$ such that $e \prec f$
(iv) $\forall f \in X_2 \exists! v_1 \neq v_2 \neq v_3 \in X_0$ such that $v_1, v_2, v_3 \prec f$

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

We call (\prec, X_0, X_1, X_2) *triangular complex* if

- (i) The relation \prec is transitive, i.e. $\forall v \in X_0, e \in X_1, f \in X_2$:

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f$$

- (ii) $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$ such that $v_1, v_2 \prec e$
(iii) $\forall e \in X_1 \exists f \in X_2$ such that $e \prec f$
(iv) $\forall f \in X_2 \exists! v_1 \neq v_2 \neq v_3 \in X_0$ such that $v_1, v_2, v_3 \prec f$
(v) $\forall f \in X_2 \exists! e_1 \neq e_2 \neq e_3 \in X_1$ such that $e_1, e_2, e_3 \prec f$

Definition

Let $X_0, X_1, X_2 \subset \mathbb{N}_0$ be finite subsets and \prec a symmetric relation between the sets called *incidence*.

We call (\prec, X_0, X_1, X_2) *triangular complex* if

- (i) The relation \prec is transitive, i.e. $\forall v \in X_0, e \in X_1, f \in X_2$:

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f$$

- (ii) $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$ such that $v_1, v_2 \prec e$
(iii) $\forall e \in X_1 \exists f \in X_2$ such that $e \prec f$
(iv) $\forall f \in X_2 \exists! v_1 \neq v_2 \neq v_3 \in X_0$ such that $v_1, v_2, v_3 \prec f$
(v) $\forall f \in X_2 \exists! e_1 \neq e_2 \neq e_3 \in X_1$ such that $e_1, e_2, e_3 \prec f$

The elements in X_0 are called *vertices*, the elements in X_1 are called *edges* and the elements in X_2 are called *faces*.

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.
Then we call (\prec, X_0, X_1, X_2) *simplicial surface* if

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.

Then we call (\prec, X_0, X_1, X_2) *simplicial surface* if

$$(i) \quad \forall e \in X_1 : |\{f \in X_2 \mid e \prec f\}| \leq 2$$

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.

Then we call (\prec, X_0, X_1, X_2) *simplicial surface* if

- (i) $\forall e \in X_1 : |\{f \in X_2 \mid e \prec f\}| \leq 2$
- (ii) $\forall v \in X_0 : |\{f \in X_2 \mid v \prec f\}| < \infty$

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.

Then we call (\prec, X_0, X_1, X_2) *simplicial surface* if

- (i) $\forall e \in X_1 : |\{f \in X_2 \mid e \prec f\}| \leq 2$
- (ii) $\forall v \in X_0 : |\{f \in X_2 \mid v \prec f\}| < \infty$
- (iii) $\forall v \in X_0$: there is an ordering of the $e_i, f_j \prec v$ such that

$$e_1 \prec f_1 \prec e_2 \prec f_2 \prec \cdots \prec f_{n-1} \prec e_n \prec f_n \prec e_1$$

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.


Then we call (\prec, X_0, X_1, X_2) *simplicial surface* if

- (i) $\forall e \in X_1 : |\{f \in X_2 \mid e \prec f\}| \leq 2$
- (ii) $\forall v \in X_0 : |\{f \in X_2 \mid v \prec f\}| < \infty$
- (iii) $\forall v \in X_0 : \text{there is an ordering of the } e_i, f_j \prec v \text{ such that}$


$$e_1 \prec f_1 \prec e_2 \prec f_2 \prec \cdots \prec f_{n-1} \prec e_n \prec f_n \prec e_1$$

the last condition is called the *umbrella condition*.

Example





Example




Simplicial Surface

Example




Simplicial Surface

Example



Simplicial Surface



Triangular complex

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.

Then we define an *embedding* of (\prec, X_0, X_1, X_2) as a map

$$c : X_0 \rightarrow \mathbb{R}^3$$

Definition

Let (\prec, X_0, X_1, X_2) be a triangular complex.

Then we define an *embedding* of (\prec, X_0, X_1, X_2) as a map

$$c : X_0 \rightarrow \mathbb{R}^3$$

The image of $v \in X_0$ is called *coordinate of v* .

Definition


Let (\prec, X_0, X_1, X_2) be a triangular complex.

Then we define an *embedding* of (\prec, X_0, X_1, X_2) as a map

$$c : X_0 \rightarrow \mathbb{R}^3$$

The image of $v \in X_0$ is called *coordinate of v* .

Example



Embedded
triangular complex

Definition


Let (\prec, X_0, X_1, X_2) be a triangular complex.

Then we define an *embedding* of (\prec, X_0, X_1, X_2) as a map


$$c : X_0 \rightarrow \mathbb{R}^3$$

The image of $v \in X_0$ is called *coordinate of v* .

Example



Embedded
triangular complex



Embedded simplicial
surface



Simplicial Surfaces Package

- Has functionality for displaying simplicial surfaces
 - Generates a .html file
 - Uses three.js

Simplicial Surfaces Package

- Has functionality for displaying simplicial surfaces
 - Generates a .html file
 - Uses three.js

Example (Number 2.1 and 2.2 from [1])



Fachpraktikum

Was a project with the goal to improve the visualizations by adding shading/local lighting.

Fachpraktikum

Was a project with the goal to improve the visualizations by adding shading/local lighting.

After some work it turns out: central class used in implementation is deprecated.

THREE.Geometry will be removed from core with r125

Discussion

geometry

**Mugen87** ✓

3 Jan '21

The upcoming release r125 will contain a major, potentially breaking change. The class `THREE.Geometry` will be no longer part of the core but moved to `jsm/deprecated/Geometry.js`. It will only be available as an ES6 module and not as a global script.

Fachpraktikum

Was a project with the goal to improve the visualizations by adding shading/local lighting.

After some work it turns out: central class used in implementation is deprecated.

THREE.Geometry will be removed from core with r125

Discussion

geometry

**Mugen87** ✓

3 Jan '21

The upcoming release r125 will contain a major, potentially breaking change. The class `THREE.Geometry` will be no longer part of the core but moved to `jsm/deprecated/Geometry.js`. It will only be available as an ES6 module and not as a global script.

→Decided to rewrite whole functionality

Advancements after rewrite

- New security requirements of JavaScript and modern browsers: need to load the code from some server → way smaller file sizes (for small examples 9kB vs. 539kB)

Advancements after rewrite

- New security requirements of JavaScript and modern browsers: need to load the code from some server → way smaller file sizes (for small examples 9kB vs. 539kB)
- More efficient Animations, faster loading, less memory (Demo in Browser)

Advancements after rewrite

- New security requirements of JavaScript and modern browsers: need to load the code from some server → way smaller file sizes (for small examples 9kB vs. 539kB)
- More efficient Animations, faster loading, less memory (Demo in Browser)
- Also works for triangular complexes
→ Does not depend on incidence structure for visualization (Demo in Browser)

Afterwards decided to roll this feature into new package:

GAP image creator

Afterwards decided to roll this feature into new package:

GAP image creator

Goal is to divide up working with triangular complexes/simplicial surfaces in `SimplicialSurfaces` and to visualize them in `GAPic`.

New features since creating GAPic

- Most settings available via GUI in real time

New features since creating GAPic

- Most settings available via GUI in real time
- `normalsMaterial` for visualizing angles of faces
 - maps colors of a face smoothly depending on the normal

New features since creating GAPic

- Most settings available via GUI in real time
- `normalsMaterial` for visualizing angles of faces
→ maps colors of a face smoothly depending on the normal
- Intersection planes
→ allows seeing inside complexes

New features since creating GAPic

- Most settings available via GUI in real time
- `normalsMaterial` for visualizing angles of faces
 - maps colors of a face smoothly depending on the normal
- Intersection planes
 - allows seeing inside complexes
- Parameterized coordinates
 - allows coordinates to be defined as any equation JavaScript can evaluate

Future plans

- Currently in beta
 - need to finish manual

Future plans

- Currently in beta
 - need to finish manual
- Partial rewrite with new data structure
 - will reduce storage especially for big animations
- Add library of often used coordinates
 - e.g. icosahedra of length 1[1]

Future plans

- Currently in beta
 - need to finish manual
- Partial rewrite with new data structure
 - will reduce storage especially for big animations
- Add library of often used coordinates
 - e.g. icosahedra of length 1[1]
- Make compatible with objects from other packages
 - e.g. simpcomp package or

Future plans

- Currently in beta
 - need to finish manual
- Partial rewrite with new data structure
 - will reduce storage especially for big animations
- Add library of often used coordinates
 - e.g. icosahedra of length 1[1]
- Make compatible with objects from other packages
 - e.g. simpcomp package or **your package?**

Thank you for your attention

Want to get involved? → github.com/GAP-ART-RWTH/GAPic

References:

- [1] Karl-Heinz Brakhage et al. *The icosahedra of edge length 1*. 2019. DOI: [10.48550/ARXIV.1903.08278](https://doi.org/10.48550/ARXIV.1903.08278). URL: <https://arxiv.org/abs/1903.08278>.