

# The GAPic Package

Lukas Schnelle

GAPDays Spring 2024

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

We call  $(\prec, X_0, X_1, X_2)$  *triangular complex* if

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

We call  $(\prec, X_0, X_1, X_2)$  *triangular complex* if

- (i) The relation  $\prec$  is transitive, i.e.  $\forall v \in X_0, e \in X_1, f \in X_2$ :

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f.$$

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

We call  $(\prec, X_0, X_1, X_2)$  *triangular complex* if

- (i) The relation  $\prec$  is transitive, i.e.  $\forall v \in X_0, e \in X_1, f \in X_2$ :

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f.$$

- (ii)  $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$  such that  $v_1 \prec e, v_2 \prec e$ .

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

We call  $(\prec, X_0, X_1, X_2)$  *triangular complex* if

- (i) The relation  $\prec$  is transitive, i.e.  $\forall v \in X_0, e \in X_1, f \in X_2$ :

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f.$$

- (ii)  $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$  such that  $v_1 \prec e, v_2 \prec e$ .  
(iii)  $\forall e \in X_1 \exists f \in X_2$  such that  $e \prec f$ .

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

We call  $(\prec, X_0, X_1, X_2)$  *triangular complex* if

- (i) The relation  $\prec$  is transitive, i.e.  $\forall v \in X_0, e \in X_1, f \in X_2$ :

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f.$$

- (ii)  $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$  such that  $v_1 \prec e, v_2 \prec e$ .

- (iii)  $\forall e \in X_1 \exists f \in X_2$  such that  $e \prec f$ .

- (iv)  $\forall f \in X_2 \exists! v_1 \neq v_2 \neq v_3 \in X_0$  such that  $v_1, v_2, v_3 \prec f$ .



## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

We call  $(\prec, X_0, X_1, X_2)$  *triangular complex* if

- (i) The relation  $\prec$  is transitive, i.e.  $\forall v \in X_0, e \in X_1, f \in X_2$ :

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f.$$

- (ii)  $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$  such that  $v_1 \prec e, v_2 \prec e$ .  
(iii)  $\forall e \in X_1 \exists f \in X_2$  such that  $e \prec f$ .  
(iv)  $\forall f \in X_2 \exists! v_1 \neq v_2 \neq v_3 \in X_0$  such that  $v_1, v_2, v_3 \prec f$ .  
(v)  $\forall f \in X_2 \exists! e_1 \neq e_2 \neq e_3 \in X_1$  such that  $e_1, e_2, e_3 \prec f$ .

## Definition

Let  $X_0, X_1, X_2 \subset \mathbb{N}_0$  be finite subsets and  $\prec$  a symmetric relation between the sets called *incidence*.

We call  $(\prec, X_0, X_1, X_2)$  *triangular complex* if

(i) The relation  $\prec$  is transitive, i.e.  $\forall v \in X_0, e \in X_1, f \in X_2$ :

$$(v \prec e) \wedge (e \prec f) \Rightarrow v \prec f.$$

(ii)  $\forall e \in X_1 \exists! v_1 \neq v_2 \in X_0$  such that  $v_1 \prec e, v_2 \prec e$ .

(iii)  $\forall e \in X_1 \exists f \in X_2$  such that  $e \prec f$ .

(iv)  $\forall f \in X_2 \exists! v_1 \neq v_2 \neq v_3 \in X_0$  such that  $v_1, v_2, v_3 \prec f$ .

(v)  $\forall f \in X_2 \exists! e_1 \neq e_2 \neq e_3 \in X_1$  such that  $e_1, e_2, e_3 \prec f$ .

The elements in  $X_0$  are called *vertices*, the elements in  $X_1$  are called *edges* and the elements in  $X_2$  are called *faces*.

## Definition

Let  $(\prec, X_0, X_1, X_2)$  be a triangular complex.


Then we call  $(\prec, X_0, X_1, X_2)$  *simplicial surface* if

- (i)  $\forall e \in X_1 : |\{f \in X_2 \mid e \prec f\}| \leq 2$
- (ii)  $\forall v \in X_0 : |\{f \in X_2 \mid v \prec f\}| < \infty$
- (iii)  $\forall v \in X_0$  : there is an ordering of the  $e_i, f_j \prec v$  such that


$$e_1 \prec f_1 \prec e_2 \prec f_2 \prec \cdots \prec f_{n-1} \prec e_n \prec f_n \prec e_1$$

the last condition is called the *umbrella condition*.

## Example





## Example




Simplicial Surface

## Example




Simplicial Surface

## Example



Simplicial Surface



Triangular complex

## Definition

Let  $(\prec, X_0, X_1, X_2)$  be a triangular complex.



## Definition

Let  $(\prec, X_0, X_1, X_2)$  be a triangular complex.

Then we define an *embedding* of  $(\prec, X_0, X_1, X_2)$  as a map

$$c : X_0 \rightarrow \mathbb{R}^3.$$

The image of  $v \in X_0$  is called *coordinate of  $v$* .

## Definition


Let  $(\prec, X_0, X_1, X_2)$  be a triangular complex.

Then we define an *embedding* of  $(\prec, X_0, X_1, X_2)$  as a map

$$c : X_0 \rightarrow \mathbb{R}^3.$$

The image of  $v \in X_0$  is called *coordinate of  $v$* .

## Example



Is an embedded triangular complex.



# Simplicial Surfaces Package

- Has functionality for displaying surfaces
  - Generates a .html file
  - Uses three.js

# Simplicial Surfaces Package

- Has functionality for displaying surfaces
  - Generates a .html file
  - Uses three.js

Example (Number 2.1 and 2.2 from [1])



## Fachpraktikum

Was a project with the goal to improve the visualizations by adding shading/local lighting

## Fachpraktikum

Was a project with the goal to improve the visualizations by adding shading/local lighting After some work it turns out: central class used in implementation is deprecated.

### THREE.Geometry will be removed from core with r125

**Discussion**

geometry

**Mugen87** ✓

3 Jan '21

The upcoming release r125 will contain a major, potentially breaking change. The class `THREE.Geometry` will be no longer part of the core but moved to `jsm/deprecated/Geometry.js`. It will only be available as an ES6 module and not as a global script.

# Goal

## Goal

Implement shading in the visualizations of the Simplicial Surface package.

# Goal

## Goal

Implement shading in the visualizations of the Simplicial Surface package.

First Approach: Implement directly

→ Learn how the output is generated

Uses a class called `THREE.Geometry`



# Workflow

**But:** In newer revisions of three.js shading is already implemented.  
→ After some promising tests: Decided to rewrite the entire function.

# Demo

We need to switch to the browser for this.  
For one example we use [1]

# Results

## ToDo

- Some of the functions are not yet implemented again

# Results

## ToDo

- Some of the functions are not yet implemented again

## Advantages

- New security requirements of javascript and modern browsers: need to load the code from server → way smaller file sizes (for small examples 9kB vs. 539kB)

# Results

## ToDo

- Some of the functions are not yet implemented again

## Advantages

- New security requirements of javascript and modern browsers: need to load the code from server → way smaller file sizes (for small examples 9kB vs. 539kB)
- More efficient Animations, faster loading, fewer memory (Demo in Browser)

# Results

## ToDo

- Some of the functions are not yet implemented again

## Advantages

- New security requirements of javascript and modern browsers: need to load the code from server → way smaller file sizes (for small examples 9kB vs. 539kB)
- More efficient Animations, faster loading, fewer memory (Demo in Browser)
- Also works for triangular complexes  
→ Does not depend on incidence structure for visualization (Demo in Browser)

## Future

- More functions in the GUI, e.g.
  - Turning the vertices on and off
  - Changing location of a vertex on the fly
- More options materials  
e.g. Color dependent on the normal of the polygon  
(Demo in browser)
- Intersection planes  
(Demo in browser)

# Thank You for your attention

Are there Questions?

- [1] Karl-Heinz Brakhage et al. *The icosahedra of edge length 1*. 2019. DOI: 10.48550/ARXIV.1903.08278. URL: <https://arxiv.org/abs/1903.08278>.