

Mechanical Comparison of Arrangement Strategies for Topological Interlocking Assemblies

Lukas Schnelle

Dec 2023

Topological Interlocking

Definition (Topological Interlocking)

A topological interlocking assembly can be defined as an arrangement of blocks that are in contact with each other together with a frame such that, if the frame is fixed, any non-empty finite subset of blocks of the arrangement is prevented from moving.

Topological Interlocking

Definition (Topological Interlocking)

A topological interlocking assembly can be defined as an arrangement of blocks that are in contact with each other together with a frame such that, if the frame is fixed, any non-empty finite subset of blocks of the arrangement is prevented from moving.

Here:

- planar topological interlocking assemblies, i.e. between two parallel planes in 3D-space,

Topological Interlocking

Definition (Topological Interlocking)

A topological interlocking assembly can be defined as an arrangement of blocks that are in contact with each other together with a frame such that, if the frame is fixed, any non-empty finite subset of blocks of the arrangement is prevented from moving.

Here:

- planar topological interlocking assemblies, i.e. between two parallel planes in 3D-space,
- use perimeter as the frame,

Topological Interlocking

Definition (Topological Interlocking)

A topological interlocking assembly can be defined as an arrangement of blocks that are in contact with each other together with a frame such that, if the frame is fixed, any non-empty finite subset of blocks of the arrangement is prevented from moving.

Here:

- planar topological interlocking assemblies, i.e. between two parallel planes in 3D-space,
- use perimeter as the frame,
- only copies of the same block differently arranged.

The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by

The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by vertices

$$B_0 := \{v_1, \dots, v_9\},$$

The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by vertices

$$B_0 := \{v_1, \dots, v_9\},$$

edges

$$\begin{aligned} B_1 := & \{ \{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_1, v_5\}, \{v_1, v_9\}, \{v_2, v_3\}, \{v_2, v_5\}, \\ & \{v_2, v_6\}, \{v_2, v_7\}, \{v_3, v_4\}, \{v_3, v_7\}, \{v_4, v_7\}, \{v_4, v_8\}, \{v_4, v_9\}, \\ & \{v_5, v_6\}, \{v_5, v_7\}, \{v_5, v_9\}, \{v_6, v_7\}, \{v_7, v_8\}, \{v_7, v_9\}, \{v_8, v_9\} \}, \end{aligned}$$

The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by vertices

$$B_0 := \{v_1, \dots, v_9\},$$

edges

$$B_1 := \{\{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_1, v_5\}, \{v_1, v_9\}, \{v_2, v_3\}, \{v_2, v_5\}, \\ \{v_2, v_6\}, \{v_2, v_7\}, \{v_3, v_4\}, \{v_3, v_7\}, \{v_4, v_7\}, \{v_4, v_8\}, \{v_4, v_9\}, \\ \{v_5, v_6\}, \{v_5, v_7\}, \{v_5, v_9\}, \{v_6, v_7\}, \{v_7, v_8\}, \{v_7, v_9\}, \{v_8, v_9\}\},$$

and triangular faces

$$B_2 := \{\{v_1, v_2, v_3\}, \{v_1, v_2, v_5\}, \{v_1, v_3, v_4\}, \{v_1, v_4, v_9\}, \{v_1, v_5, v_9\}, \\ \{v_2, v_3, v_7\}, \{v_2, v_5, v_6\}, \{v_2, v_6, v_7\}, \{v_3, v_4, v_7\}, \{v_4, v_7, v_8\}, \\ \{v_4, v_8, v_9\}, \{v_5, v_6, v_7\}, \{v_5, v_7, v_9\}, \{v_7, v_8, v_9\}\}.$$

The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by B_0, B_1, B_2 together with coordinates

$$v_1 = (0, 0, 0), v_2 = (1, 1, 0), v_3 = (2, 0, 0),$$

$$v_4 = (1, -1, 0), v_5 = (0, 1, 1), v_6 = (1, 1, 1),$$

$$v_7 = (1, 0, 1), v_8 = (1, -1, 1), v_9 = (0, -1, 1).$$


The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by B_0, B_1, B_2 together with coordinates

$$v_1 = (0, 0, 0), v_2 = (1, 1, 0), v_3 = (2, 0, 0),$$

$$v_4 = (1, -1, 0), v_5 = (0, 1, 1), v_6 = (1, 1, 1),$$

$$v_7 = (1, 0, 1), v_8 = (1, -1, 1), v_9 = (0, -1, 1).$$




The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by B_0, B_1, B_2 together with coordinates

$$v_1 = (0, 0, 0), v_2 = (1, 1, 0), v_3 = (2, 0, 0),$$

$$v_4 = (1, -1, 0), v_5 = (0, 1, 1), v_6 = (1, 1, 1),$$

$$v_7 = (1, 0, 1), v_8 = (1, -1, 1), v_9 = (0, -1, 1).$$






The Versatile Block

The Versatile Block is a polyhedron embedded in \mathbb{R}^3 , given by B_0, B_1, B_2 together with coordinates

$$v_1 = (0, 0, 0), v_2 = (1, 1, 0), v_3 = (2, 0, 0),$$

$$v_4 = (1, -1, 0), v_5 = (0, 1, 1), v_6 = (1, 1, 1),$$

$$v_7 = (1, 0, 1), v_8 = (1, -1, 1), v_9 = (0, -1, 1).$$



Isometries

Definition

Let $n \in \mathbb{N}$, V an euclidean K vectorspace (with metric $d : V \times V \rightarrow K$). Then $\varphi : V \rightarrow V$ is called an *isometry* if

Isometries

Definition

Let $n \in \mathbb{N}$, V an euclidean K vectorspace (with metric $d : V \times V \rightarrow K$). Then $\varphi : V \rightarrow V$ is called an *isometry* if

$$\forall x, y \in V : d(x, y) = d(\varphi(x), \varphi(y)).$$

Isometries

Definition

Let $n \in \mathbb{N}$, V an euclidean K vectorspace (with metric $d : V \times V \rightarrow K$). Then $\varphi : V \rightarrow V$ is called an *isometry* if

$$\forall x, y \in V : d(x, y) = d(\varphi(x), \varphi(y)).$$

Note: An isometry is always a combination of an affine transformation and an orthogonal matrix. Such an isometry $\varphi = (A, a)$ operates as

$$(A, a)(x) := A \cdot x + a.$$

Isometries

Definition

Let $n \in \mathbb{N}$, V an euclidean K vectorspace (with metric $d : V \times V \rightarrow K$). Then $\varphi : V \rightarrow V$ is called an *isometry* if

$$\forall x, y \in V : d(x, y) = d(\varphi(x), \varphi(y)).$$

Note: An isometry is always a combination of an affine transformation and an orthogonal matrix. Such an isometry $\varphi = (A, a)$ operates as

$$(A, a)(x) := A \cdot x + a.$$

The isometries of a given euclidean vectorspace are a group with the operation

$$((A, a) \circ (B, b))(x) := (A \cdot B, A \cdot a + b)$$

denoted by $E(n)$

Isometries

Definition

Let $n \in \mathbb{N}$, V an euclidean K vectorspace (with metric $d : V \times V \rightarrow K$). Then $\varphi : V \rightarrow V$ is called an *isometry* if

$$\forall x, y \in V : d(x, y) = d(\varphi(x), \varphi(y)).$$

Note: An isometry is always a combination of an affine transformation and an orthogonal matrix. Such an isometry $\varphi = (A, a)$ operates as

$$(A, a)(x) := A \cdot x + a.$$

The isometries of a given euclidean vectorspace are a group with the operation

$$((A, a) \circ (B, b))(x) := (A \cdot B, A \cdot a + b)$$

denoted by $E(n)$ ($\implies E(n) = O(n) \ltimes \mathbb{R}^n$).

Wallpaper Groups

Definition

Crystallographic/Wallpaper Groups Let $\Gamma \leq E(n)$ a subgroup of the group of isometries of dimension n .

Then Γ is called *crystallographic group* if Γ is cocompact and discrete, it is called *wallpaper group* if $n = 2$.

Wallpaper Groups

Definition

Crystallographic/Wallpaper Groups Let $\Gamma \leq E(n)$ a subgroup of the group of isometries of dimension n .

Then Γ is called *crystallographic group* if Γ is cocompact and discrete, it is called *wallpaper group* if $n = 2$.

Definition

Let $\Gamma \leq E(n)$.

Then Γ is cocompact if the space $E(n)/\Gamma$ is compact.

Wallpaper Groups

Definition

Crystallographic/Wallpaper Groups Let $\Gamma \leq E(n)$ a subgroup of the group of isometries of dimension n .

Then Γ is called *crystallographic group* if Γ is cocompact and discrete, it is called *wallpaper group* if $n = 2$.

Definition

Let $\Gamma \leq E(n)$.

Then Γ is cocompact if the space $E(n)/\Gamma$ is compact.

Proposition ([1, Prop. 1.9])

Let $\Gamma \leq E(n)$.

Then $E(n)/\Gamma$ is compact iff the orbit space \mathbb{R}^n/Γ is compact.

Examples of wallpaper groups

$p1 :=$

$$\left\langle \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right), \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) \right\rangle$$

Examples of wallpaper groups

$p1 :=$

$$\left\langle \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right), \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) \right\rangle$$

$pg :=$

$$\left\langle \left(\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \end{pmatrix} \right), \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) \right\rangle$$

Examples of wallpaper groups

$p1 :=$

$$\left\langle \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right), \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) \right\rangle$$

$pg :=$

$$\left\langle \left(\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \end{pmatrix} \right), \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) \right\rangle$$

$p4 :=$

$$\left\langle \left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 2 \end{pmatrix} \right), \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 \\ -2 \end{pmatrix} \right), \left(\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) \right\rangle$$

Planar Assemblies of the Versatile Block

Finite Element Method

Problem formulation

Simulation Setup

Stresses

Combinatorial Method

Combinatorial Results

- [1] A. Szczepanski and W. Scientific. *Geometry of Crystallographic Groups*. Algebra and discrete mathematics. World Scientific, 2012. URL:
<https://books.google.de/books?id=wX26CgAAQBAJ>.