Inward Pointing Vectors on Manifolds with Corners

Hannah Scholz

Mathematical Institute of the University of Bonn

06.11.2026

Model with Corners

We will focus on real manifolds.
Let E be a normed space over $\mathbb{R}$ and H a topological space.

Definition: Model with Corners

A *model with corners* from E to H is a partial equivalence $I : H \to E$ such that

(i) the forward and backward map are continuous,

(ii) then source is the entirety of H, and

(iii) the range is convex and has nonempty interior.

Example picture here (one for boundary one for corners one circle)

Topological Manifold with Corners

Definition: Topological Manifold with Corners

A topological space M is a *topological n-manifold* if

(i) M is Hausdorff.

(ii) M is second-countable, i.e. M has a countable base.

(iii) M is locally H, i.e. for every point p there is a neighbourhood of p that is homeomorphic to an open set in H.

Definition: Chart

Let M be a topological manifold with corners. Let U be an open subset of M and $\varphi\colon U \to H$ a homeomorphism onto an open subset of H. Then we call (U, φ) a chart of M.


Boundary of a Topological Manifold

Definition: Boundary

Let M be a topological manifold with boundary.

(i) A point $p \in M$ is a *boundary point* if it is in the domain of a *boundary chart*, i.e. a chart (U, φ) with $I(\varphi(p)) \in \partial(I(H))$.

(ii) A point $p \in M$ is an *interior point* if it is in the domain of an *interior chart*, i.e. a chart (U, φ) with $I(\varphi(U)) \cap \partial(I(H)) = \varnothing$.



Smooth Manifold with Corners

Definition: Smooth Atlas

Let M be a topological manifold. Let $\mathcal{A}$ be a collection of charts whose domains cover M. $\mathcal{A}$ is a smooth atlas for M if any two charts (φ, U) and (ψ, V) are *smoothly compatible*, i.e. either $U \cap V$ is empty or $I \circ \psi \circ \varphi^{-1} \circ I^{-1} \colon I(\varphi(U \cap V)) \to \psi I((U \cap V))$ is smooth in the ordinary sense.

Definition: Smooth Manifold

A pair $(M, \mathcal{A})$ of a topological manifold M and a smooth altas $\mathcal{A}$ on M, which is maximal with respect to inclusion, is called a *smooth manifold*.

Intuition

pictures of inward pointing and not inward pointing vectors

Definition on Manifold without Corners

Inward pointing [Lee06]

Let M be a smooth manifold with boundary without corners. Let $p \in M$. Then $v \in T_pM \setminus T_p\partial M$ is inward pointing if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma\colon [0, \varepsilon) \to M$ such that $\gamma(0) = p$ and $\gamma'(0) = v$.

first have picture for intuition here

Definition on Manifold without Corners

Inward pointing [Lee06]

Let M be a smooth manifold with boundary without corners. Let $p \in M$. Then $v \in T_pM \setminus T_p\partial M$ is inward pointing if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma\colon [0, \varepsilon) \to M$ such that $\gamma(0) = p$ and $\gamma'(0) = v$.

then picture of square here

Options for definition

Definition: (weakly) inward pointing/realisable

We call a vector that is inward pointing or tangent to the boundary *realisable* or *weakly inward pointing*.

We have the following two options for the definition of inward-pointing vectors:

(A) intrinsically exclude tangent vectors, or

(B) define realisable vectors and then exclude tangent vectors

Option A: attempt 1

Attempt 1: our idea inspired by [Lee06]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is inward pointing if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma\colon [0, \varepsilon) \to M$ such that $\gamma(0) = p$, $\gamma'(0) = v$ and $\gamma((0, \varepsilon)) \subseteq \mathrm{int}(M)$.

Option A: attempt 1

Attempt 1: our idea inspired by [Lee06]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is inward pointing if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma\colon [0, \varepsilon) \to M$ such that $\gamma(0) = p$, $\gamma'(0) = v$ and $\gamma((0, \varepsilon)) \subseteq \mathrm{int}(M)$.

picture of why this doesn't work

Option A: attempt 2

Attempt 2: adapted from [Mel96]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is inward pointing if and only if $D_v f(p) \geq 0$ for every non-negative $f \in C^\infty(M)$ that is zero on ∂M.

This definition is clearly designed to mean weakly inward-pointing.

Option A: attempt 3

Attempt 3: our idea inspired by [Mel96]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is inward pointing if and only if there is a non-negative $f \in C^\infty(M)$ that is zero on ∂M such that $\mathrm{D}_v f(p) > 0$.

Option A: attempt 3

Attempt 3: our idea inspired by [Mel96]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is inward pointing if and only if there is a non-negative $f \in C^\infty(M)$ that is zero on ∂M such that $\mathrm{D}_v f(p) > 0$.

picture of why this doesn't work

Option A: attempt 4

Attempt 4: e.g. [Joy09]

Let M be a smooth manifold with corners over the model with corners $\mathbb{R}^n_k = [0,\infty)^k \times \mathbb{R}^{n-k}$. Let $p \in M$ be a corner point of codimension k, i.e. let there be a chart (U, φ) with $U \subseteq \mathbb{R}^n_k$ such that $0 \in U$ and $\varphi(U) = p$. Then a vector $v \in T_pM$ is inward pointing if and only if it is in $\mathrm{d}\varphi_0((0,\infty)^k \times \mathbb{R}^{n-k})$ where $\mathrm{d}\varphi_0 \colon \mathbb{R}^n = T_0\mathbb{R}^n_k \to T_pM$.

Option A: attempt 4

Attempt 4: e.g. [Joy09]

Let M be a smooth manifold with corners over the model with corners $\mathbb{R}^n_k = [0,\infty)^k \times \mathbb{R}^{n-k}$. Let $p \in M$ be a corner point of codimension k, i.e. let there be a chart (U, φ) with $U \subseteq \mathbb{R}^n_k$ such that $0 \in U$ and $\varphi(U) = p$. Then a vector $v \in T_pM$ is inward pointing if and only if it is in $\mathrm{d}\varphi_0((0,\infty)^k \times \mathbb{R}^{n-k})$ where $\mathrm{d}\varphi_0 \colon \mathbb{R}^n = T_0\mathbb{R}^n_k \to T_pM$.

Defining inward-pointing from realisability

Definition: inward-pointing, [Amr21]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is inward pointing if and only if it is in the interior of the set of realisable vectors.

Conditions for a definition inward-pointing

A sensible definition of realisability needs to fulfil these two conditions:

(i) The definition needs to be preserved by local diffeomorphisms (in particular it should be preserved by smooth charts).

(ii) The definition should reduce to a sensible notion on the model space that is easily verifiable in the cases of the typical examples (e.g. upper half space in $\mathbb{R}^n$).

Conditions for a definition inward-pointing

A sensible definition of realisability needs to fulfil these two conditions:

(i) The definition needs to be preserved by local diffeomorphisms (in particular it should be preserved by smooth charts).

(ii) The definition should reduce to a sensible notion on the model space that is easily verifiable in the cases of the typical examples (e.g. upper half space in $\mathbb{R}^n$).

Sensible notion in normed spaces: tangent cones

Definition: contingent Tangent Cone, [AF90]

Let E be a normed vector space, $K \subset E$ and $x \in \overline{E}$. Then the *contingent tangent cone* to K at x is the set of all $v \in E$ such that there is a sequence of scalars $c\colon \mathbb{N} \to \mathbb{R}_{\geq}$ and a sequence of vectors $d\colon \mathbb{N} \to E$ such that

(i) $\lim_{n\to\infty} d_n = 0$,

(ii) $x + d_n \in K$ for all $n \in \mathbb{N}$, and

(iii) $\lim_{n\to\infty} c_n \cdot d_n = v$.

We denote this tangent cone by $T_K(x)$.

This is in Mathlib as `posTangentConeAt`. For convex sets this notion agrees with other types of tangent cones such as the Clarke tangent cone.
Pictures for definition and examples of tangent cones

Option B: attempt 1

Definition: realisability, [Lee06]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is *realisable* if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma\colon [0, \varepsilon) \to M$ such that $\gamma(0) = p$ and $\gamma'(0) = v$.

Option B: attempt 1

Definition: realisability, [Lee06]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is *realisable* if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma\colon [0, \varepsilon) \to M$ such that $\gamma(0) = p$ and $\gamma'(0) = v$.

picture of why this is really hard

Option B: attempt 1

Definition: realisability, [Lee06]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is *realisable* if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma\colon [0, \varepsilon) \to M$ such that $\gamma(0) = p$ and $\gamma'(0) = v$.

Requiring γ to be smooth instead of continuous would work better.

Option B: final pick

Definition: Realisability, [RW98]

Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_pM$ is *realisable* if and only if there is a function $\gamma\colon [0, \infty) \to M$ that is differentiable at 0 and fulfils $\gamma(0) = p$ and $\gamma'(0) = v$.

```
def IsRealizable {M H E : Type*} [TopologicalSpace H]
    [TopologicalSpace M] [ChartedSpace H M]
    [NormedAddCommGroup E] [NormedSpace ℝ E]
    {I : ModelWithCorners ℝ E H} {p : M}
    (v : TangentSpace I p) : Prop :=
  ∃ (γ : ℝ → M) (_ : MDiffAt[Ici 0] γ 0), γ 0 = p ∧
    mfderiv[Ici 0] γ (0 : ℝ) 1 = v
```

Properties of definition

Lemma: invariance under charts

Let $f : M \to M'$ be a local diffeomorphism between manifolds with corners. Let $p \in M$ and $v \in T_pM$. Then v is realisable if and only if $\mathrm{D}_v f(p)$ is realisable.

References I

[AF90] Jean-Pierre Aubin and Hélène Frankowska. *Set-valued analysis*. Systems & control 2. Boston: Birkhaeuser, 1990.

[Amr21] Amr (https://math.stackexchange.com/users/29267/amr). *Definition of an inward-pointing tangent vector for smooth manifolds with corners*. Mathematics Stack Exchange. 2021.

[Joy09] Dominic Joyce. "On manifolds with corners". In: *Advances in Geometric Analysis* (Oct. 2009).

References II

[Lee06] John M. Lee. *Introduction to smooth manifolds*. Graduate texts in mathematics 218. New York: Springer, 2006.

[Mel96] Richard Melrose. *Differential analysis on manifolds with corners*. Sept. 1996.

[RW98] Ralph Tyrrell Rockafellar and Roger J.-B. Wets. *Variational analysis*. Grundlehren der mathematischen Wissenschaften 317. Springer, 1998.