

????

Hannah Scholz

Born 21st July, 2003 in Bonn, Germany

22nd August, 2024

Master's Thesis Mathematics

Advisor: Dr. Michael Rothgang

Second Advisor: Prof. Dr. Floris van Doorn

MATHEMATISCHES INSTITUT

MATHEMATISCH-NATURWISSENSCHAFTLICHE FAKULTÄT DER
RHEINISCHEN FRIEDRICH-WILHELMS-UNIVERSITÄT BONN

Contents

1	The mathematics of collar neighborhoods	5
1.1	Prerequisites	5
1.2	Basics about collar neighbourhoods	5
2	The proof of the collar neighbourhood theorem for smooth manifolds	9
2.1	Prerequisites	9
2.1.1	Vector fields	9
2.1.2	Flows and Flowouts	10
3	Inward Pointing Vectors in Manifolds with Corners	11
	Bibliography	13

1 The mathematics of collar neighborhoods

1.1 Prerequisites

Lemma 1.1.1 (Smooth Urysohn's lemma, in mathlib I think). *Let M be a $\mathcal{C}^k$ -manifold and A and B two disjoint closed sets in M . Then there is a $\mathcal{C}^k$ -function $f: M \rightarrow [0, 1]$ such that f is 0 on A and 1 on B .*

Remark 1.1.2. Again, this requires a definition of manifold that makes it metrizable.

1.2 Basics about collar neighbourhoods

Definition 1.2.1. A *topological manifold* is a second-countable Hausdorff space that is locally Euclidean.

Remark 1.2.2. Apparently, there are some subtleties with collars of non-metrizable manifolds but I assume we don't care. (At least with the above definition our manifolds are always metrizable.)

Definition 1.2.3. Let M and N be two $\mathcal{C}^k$ -manifolds. Let $f: M \rightarrow N$.

- (i) f is a $\mathcal{C}^k$ -diffeomorphism if it is bijective and both f and its inverse functions are k -times continuously differentiable.
- (ii) f is a $\mathcal{C}^k$ -embedding if it is a $\mathcal{C}^k$ -diffeomorphism onto its image.

Definition 1.2.4 (Due to [Con71]). Let M be a $\mathcal{C}^k$ -manifold (with boundary) and let $B \subseteq M$ be closed.

- (i) B is *locally collared* if it is covered by sets $U \subseteq B$ which are open in B such that there are $\mathcal{C}^k$ -embeddings $\varphi_U: \overline{U} \times [0, 1] \rightarrow M$ with
 - a) $\varphi_U^{-1}(B) = \overline{U} \times \{0\}$.
 - b) For all $x \in \overline{U}$, $\varphi_U(x, 0) = x$.
 - c) $\varphi_U(\overline{U} \times [0, 1])$ is closed.
 - d) $\varphi_U(U \times [0, 1])$ is open.
- (ii) B is *collared* if B already fulfils the conditions given above. In this case we call the image of $B \times [0, 1)$ under the corresponding map $\varphi_B: B \times [0, 1] \rightarrow M$ a *collar neighbourhood* of B .

Remark 1.2.5. This definition is a lot more restrictive than other definition which don't consider the closure or the closed interval. I think if this one doesn't work, I can just take the other one.

Remark 1.2.6. For $k = 0$ we can even extend this definition to topological spaces instead of just defining it for topological manifolds.

Theorem 1.2.7 (Due to [Con71]). *Let X be a Hausdorff space and $B \subseteq X$ be a closed subset. If B is locally collared in X , then it is collared.*

Proof. Let $U_1, \dots, U_n$ be a cover of B that witnesses that B is locally collared. Let $\varphi_i: \overline{U_i} \times [0, 1] \rightarrow X$ be the corresponding embeddings. B is normal because it is compact and Hausdorff. By the shrinking lemma (already in mathlib) there is a new open cover $V_1, \dots, V_n$ of B such that $\overline{V_i} \subseteq U_i$ for all $1 \leq i \leq n$. Define

$$X^+ := (X \amalg B \times [-1, 0]) / x \sim (x, 0).$$

Clearly, X^+ is collared. So it suffices to show that there is a homeomorphism $G: X \rightarrow X^+$ s.t. $G(x) = (x, -1)$ for all $x \in B$. Let $\Phi_i: \overline{U_i} \times [-1, 1] \rightarrow X^+$ be defined by

$$\Phi_i(x) = \begin{cases} \phi_i(x) & \text{if } x \in \overline{U_i} \times [0, 1] \\ x & \text{if } x \in \overline{U_i} \times [-1, 0]. \end{cases}$$

We now inductively define maps $f_i: B \rightarrow [0, 1]$ and embeddings $g_i: X \rightarrow X^+$ for all $0 \leq i \leq n$ such that

- (i) $f_i(x) = -1$ if $x \in \bigcup_{j \leq i} \overline{V_j}$
- (ii) $g_i(x) = (x, f_i(x))$ if $x \in B$
- (iii) $g_i(X) = X \cup \{(x, t) \mid t \geq f_i(x)\}$

Then g_n will be the desired homeomorphism G from above. Set f_0 to be constant 0. Then we can choose g_0 to be the inclusion and trivially $g_0(X) = X = X \cup X \times \{0\} = X \cup \{(x, t) \mid t \geq 0\}$.

Assume that g_{i-1} has been defined. Using Urysohn's Lemma, choose a continuous function $\lambda_i: \overline{U_i} \rightarrow [0, 1]$ such that it is 0 on $\overline{U_i} \setminus U_i$ and 1 on $\overline{V_i}$. Given $x \in B$ set $a(x) := f_{i-1}(x)$ and $b(x) := (1 - \lambda_i(x))f_{i-1}(x) + \lambda_i(x)(-1)$. Now let

$$s(x, _): [a(x), 1] \rightarrow [b(x), 1], \quad t \mapsto \frac{b(x) - 1}{a(x) - 1}(t - 1) + 1.$$

Now set

$$\psi_i: \Phi_i^{-1}(g_{i-1}(X)) \rightarrow \overline{U_i} \times [-1, 1], \quad (x, t) \mapsto (x, s(x, t)).$$

This allows us to define

$$\Psi_i: g_{i-1}(X) \rightarrow X^+, \quad x \mapsto \begin{cases} \Phi_i(\psi_i(\Phi_i^{-1}(x))) & \text{if } x \in g_i(X) \cap \Phi_i(\overline{U_i} \times [-1, 1]) \\ x & \text{otherwise} \end{cases}$$

Claim 1: Ψ_i is continuous.

Proof. It suffices to show that ψ_i is continuous. We need to show that $s: B \times [a(x), 1] \rightarrow [b(x), 1]$ is continuous.

Ahhh this is ugly. I somehow would prefer to just get going with this in Lean. $\square$

□

I want to try to separate out some parts here:

Definition 1.2.8. Let X be a $(n+1)$ -dimensional $\mathcal{C}^k$ -manifold and $B \subseteq X$ a closed subset with some nice properties (n -dimensional). We define the topological space

$$X_B^+ := (X \amalg B \times [-1, 0]) / x \sim (x, 0).$$

Lemma 1.2.9. Let X be an n -dimensional $\mathcal{C}^k$ -manifold and $B \subseteq X$ a closed subset. Then X_B^+ is an n -dimensional $\mathcal{C}^k$ -manifold.

Proof. We first need to show that X_B^+ is a Hausdorff space. I am not sure if this is best done manually or if there is some cool lemma like: a pushout along closed embeddings preserves Hausdorff spaces. (Is this true?)

Then we need to show that X_B^+ is second countable. Again, there should be some general lemma about closed embeddings and pushouts.

Now we need to pick an atlas for X_B^+ . Let $\mathcal{A}_X$ be a $\mathcal{C}^k$ -atlas for X , $\mathcal{A}_B$ be a $\mathcal{C}^k$ -atlas for B and $\mathcal{A}_{[-1,0]}$ be a $\mathcal{C}^k$ -atlas for $[-1, 0]$. We define

$$\begin{aligned} \mathcal{A} := & \{(U, \varphi) \in \mathcal{A}_X \mid U \cap B = \emptyset\} \\ & \cup \{(U \times I, \varphi \times \psi) \mid (U, \varphi) \in \mathcal{A}_B, (I, \psi) \in \mathcal{A}_{[-1,0]}, 0 \notin I\} \\ & \cup \{(U \cup ((U \cap B) \times I), \varphi_I^+) \mid (U, \varphi) \in \mathcal{A}, (I, \psi) \in \mathcal{A}_{[-1,0]}, 0 \in I\} \end{aligned}$$

where we define φ_I^+ as follows. Let (U, φ) and (I, ψ) as above. Set

$$\varphi_I^+ : U \cup ((U \cap B) \times I) \rightarrow \mathbb{R}^n, x \mapsto \begin{cases} \varphi(x) & x \in U \\ (\varphi(y), z) & x = (y, z) \in (U \cap B) \times I \end{cases}$$

For $x \in B$ we have $\varphi(x)_I^+ = \varphi(x) = (\varphi(x), 0) = \varphi_I^+(x, 0)$, so φ_I^+ is continuous.

It seems to me that this might not be $\mathcal{C}^k$ at all. I don't know if you can even glue this without basically already having the vector field available and then one might as well do the other proof. Maybe a local vector field is enough? Not sure :(

Now you would need to verify that this is a $\mathcal{C}^k$ atlas. Possibly using a version of the manifold chart lemma from Lee. I will only try to show that the transition maps are $\mathcal{C}^k$. The rest should be easier. The only interesting intersections are those of two charts of the form $(U \cup ((U \cap B) \times I), \varphi_I^+)$ and $(V \cup ((V \cap B) \times J), \psi_J^+)$. We need to check that the transition map is $\mathcal{C}^k$ on $U \cap V \cap B$. This is clear for all combinations of partial derivatives that do not include the n -th one. □

Lemma 1.2.10. Let X be an n -dimensional $\mathcal{C}^k$ -manifold and $B \subseteq X$ a closed subset. Then X_B^+ is the pushout in the category of $\mathcal{C}^k$ -manifolds.

Remark 1.2.11. It seems that this is not at all trivial and in fact much of the issue with gluing manually along boundaries. I am not sure if we can even guarantee the above lemma. The above lemma would hopefully mean that one never has to look at the explicit construction again.

Definition 1.2.12. Let X be a Hausdorff space and $B \subseteq X$ a closed subset. Let $U \subseteq B$ be an open subset together with a

We would like the following theorem:

Theorem 1.2.13. *Let M be a C^k -manifold (with boundary) and $B \subseteq X$ a closed subset. If B is locally collared in M , then it is collared.*

Then we will need the following lemma:

Lemma 1.2.14. *Let M be an n -dimensional C^k -manifold with boundary. Then ∂M is locally collared.*

Proof. Let $p \in \partial M$. Then p is contained in a regular half-coordinate ball U , i.e. there is a coordinate half-ball $U' \supseteq \bar{U}$ and a coordinate map $\psi: U' \rightarrow \mathbb{H}^n$ such that for some $r' > r > 0$ we have $\psi(U) = B_r(0) \cap \mathbb{H}^n$, $\psi(\bar{U}) = \overline{B_r(0)} \cap \mathbb{H}^n$ and $\psi(U') = B_{r'}(0) \cap \mathbb{H}^n$.

Now pick an open neighbourhood V of $0 \in \partial \mathbb{H}^n$ and $l > 0$ s.t. $V \times [0, l] \subseteq B_r(0)$.

Considering $\bar{V}$ and V as submanifolds of $\mathbb{H}^n$ and $\psi^{-1}(\bar{V}) = \overline{\psi^{-1}(V)}$ and $\psi^{-1}(V)$ as submanifolds. Then

$$\psi|_{\overline{\psi^{-1}(V)}}: \overline{\psi^{-1}(V)} \rightarrow \bar{V}$$

and

$$\psi|_{\psi^{-1}(V)}: \psi^{-1}(V) \rightarrow V$$

are C^k -diffeomorphisms. So it suffices to show that $\psi^{-1}|_{\bar{V} \times [0, l]}: \bar{V} \times [0, l] \rightarrow M$ fulfils the conditions given in Definition 1.2.4.

I think this should work... I think it would be nicer to do this directly in Lean and then later write up the maths proof. $\square$

which will then allow us to derive

Theorem 1.2.15 (Collar neighbourhood theorem). *Let M be a C^k -manifold with boundary, then ∂M is collared.*

2 The proof of the collar neighbourhood theorem for smooth manifolds

2.1 Prerequisites

2.1.1 Vector fields

Lemma 2.1.1. *Let M be a smooth n -manifold with boundary. Let B be a coordinate half ball with corresponding map $\varphi: B \rightarrow \mathbb{H}^n$. Then $\frac{\partial}{\partial x^n}$ is a smooth vector field on B which is inward-pointing when restricted to $\partial M \cap B$.*

Proof. The vector field is smooth by Example 8.2 in [Lee06]. Let $p \in B$. We need to show that $\frac{\partial}{\partial x^n} \Big|_p$ is inward-pointing. By Proposition 5.42 in [Lee06], it suffices to show that the n -th component of $\frac{\partial}{\partial x^n} \Big|_p$ is positive. Its n -th component is 1 so we are done. $\square$

Lemma 2.1.2. *Let M be a smooth manifold, A a closed and U an open subset of M with $A \subseteq U$. Let $X: A \rightarrow TM$ be a smooth vector field along A . Then there is a smooth vector field $\tilde{X}: M \rightarrow TM$ with support in U which agrees with X on A .*

Proof. We first notice that the proof of Lemma 2.26 (Extension Lemma for Smooth Functions) in [Lee06] also works for the domain TM . Now we need to verify that a smooth map of the form

$$\tilde{X} = \sum_{p \in A} \tau_p(x) \tilde{X}_p(x)$$

is a vector field. Here the $\tau_p x$ are bump functions and the $\tilde{X}_p$ local extensions of the vector field. Now locally this sum is finite. Module over $C^\infty(M)$... blah blah blah $\square$

Lemma 2.1.3 (Problem 8-4 in [Lee06]). *Let M be a smooth n -manifold with boundary. There exists a smooth vector field on M , which is inward-pointing when restricted to the boundary.*

Proof. Cover ∂M with a countable number of regular coordinate half-balls $(B_i)_i$. This is possible by Exercise 1.42 in [Lee06]. Let us call the corresponding larger coordinate half-balls B'_i and the corresponding smooth coordinate map $\varphi_i: B'_i \rightarrow \mathbb{H}^n$. By Lemma 2.1.1, $\frac{\partial}{\partial x^n}$ is a smooth vector field on B'_i which is inward-pointing when restricted to $\partial M \cap B$. In particular, $\frac{\partial}{\partial x^n}$ is a smooth vector field along B_i . Using Lemma 2.1.2, we can extend these to smooth vector fields $X_i: M \rightarrow TM$ with support in B'_i and which agree with $\frac{\partial}{\partial x^n}$ on B_i . This is again inward-pointing which is easy to see using Proposition 5.42 in [Lee06]. Now take a partition of unity subordinate to the cover $\{B_i \mid i\} \cup M \setminus \partial M$. Then do the same thing as in the proof of Lemma 2.1.2 and get a global smooth vector field that is inward-pointing on the boundary. $\square$

General
Lemma?

write
Lemma

2.1.2 Flows and Flowouts

Lemma 2.1.4. *Let U be an open subset of $\mathbb{H}^n$ that has non-zero intersection with the boundary. Let V be a smooth vector field on U . Then there exists a smooth map $\delta_U: U \rightarrow \mathbb{R}^+$ and a smooth open embedding Φ_U :*

Lemma 2.1.5. *Let M be a smooth n -manifold with boundary. Let N be a smooth vector field on M that is inward-pointing when restricted to the boundary. Let (U, φ) be a boundary chart. Then there exists a smooth map $\delta_U: U \rightarrow \mathbb{R}^+$ and a smooth open embedding $\Phi_U: P_U \rightarrow M$ where $P_U := \{(t, p) \in \mathbb{R} \times U \mid 0 \leq t < \delta_U(p)\}$...*

Proof. Consider the pushforward φ_*X of X by φ on $\varphi(U) \subseteq \mathbb{H}^n$. Extend φ_*X to a vector field V on an open subset U' of $\mathbb{R}^n$ with $U' \cap \mathbb{H}^n = U$ such that still $V^n(p) > 0$ for all $p \in \partial\mathbb{H}^n$. Now let $\theta_V: \mathcal{D} \rightarrow U'$ be the flow of V as given by the Fundamental Theorem of Flows (Theorem 9.12 in [Lee06]) where $\mathcal{D} \subseteq \mathbb{R} \times M$ is a flow domain. Since $\varphi(U) \cap \partial\mathbb{H}^n$ is an embedded $n - 1$ -dimensional submanifold of U' and V is clearly nowhere tangent to it, we can apply the Flowout Theorem (Theorem 9.20 in [Lee06]) to get a smooth function $\delta'_U: \varphi(U) \cap \partial\mathbb{H}^n \rightarrow \mathbb{R}^+$ such that $\theta_V|_{\mathcal{O}}: \mathcal{O} \rightarrow U$ is a smooth embedding where $\mathcal{O} := \{(t, p) \in (\mathbb{R} \times (\varphi(U) \cap \partial\mathbb{H}^n) \cap \mathcal{D}) \mid 0 \leq t < \delta'_U(p)\}$.

Now define

□

Lemma 2.1.6 (Boundary Flowout Theorem, Theorem 9.24 and Problem 9-11 in [Lee06]). *Let M be a smooth manifold with non-empty boundary. Let N be a smooth vector field on M that is inward-pointing when restricted to the boundary. Then there exists a smooth map $\delta: \partial M \rightarrow \mathbb{R}^+$ and a smooth embedding $\Phi_\delta: P_\delta \rightarrow M$ where $P_\delta := \{(t, p) \in \mathbb{R} \times \partial M \mid 0 \leq t < \delta(p)\}$ such that $\Phi_\delta(P_\delta)$ is a neighbourhood of ∂M and for every $p \in \partial M$, $t \mapsto \Phi_\delta(t, p)$ is an integral curve of N starting at p .*

Proof. Cover ∂M with a countable number of boundary charts (U, φ) .

□

3 Inward Pointing Vectors in Manifolds with Corners

Defining the notion of inward pointing vectors in manifolds with corners is substantially harder than in manifolds with boundary without corners. Let us first examine why this is.

Here is a common definition for manifolds with boundary without corners:

Definition 3.0.1 ([Lee06] p. 136). Let M be a smooth manifold with boundary without corners. Let $p \in M$. Then $v \in T_p M \setminus T_p \partial M$ is inward pointing if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma: [0, \varepsilon) \rightarrow M$ such that $\gamma(0) = p$ and $\gamma'(0) = v$.

This definition cannot be used for manifolds with corners as in this case ∂M is not always a manifold and thus $T_p \partial M$ is not defined.

The existence of the $\gamma: [0, \varepsilon) \rightarrow M$ by itself characterises whether a vector is inward pointing or tangent to the boundary, a notion which we will call *realisable*.

So for manifolds with corners we either need a completely new definition that intrinsically excludes the vectors tangent to the boundary or we need a definition of realisability and a way to exclude the tangent vectors.


We first tried the first option and came up with these candidates for definitions which we either came up with ourselves or found in the literature:

Definition 3.0.2 (Our idea inspired by Definition 3.0.1). Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_p M$ is inward pointing if and only if there is an $\varepsilon > 0$ and a smooth curve $\gamma: [0, \varepsilon) \rightarrow M$ such that $\gamma(0) = p$, $\gamma'(0) = v$ and $\gamma((0, \varepsilon)) \subseteq \text{int}(M)$.

Unfortunately, this definition doesn't exclude vectors tangent to the boundary:

Example 3.0.3. Take M to be the upper half plane $\mathcal{H}$ in $\mathbb{R}^2$ let p be $(0, 0)$ and let $\gamma: [0, 1) \rightarrow \mathcal{H}, x \mapsto x^2$. Then $\gamma'(0) = (1, 0)$ and thus by this definition $(1, 0)$ would be inward pointing even though it is clearly tangent to the boundary.

Definition 3.0.4. Let M be a smooth manifold with corners. Let $p \in M$. Then $v \in T_p M$ is inward pointing if and only if $D_v f(p) \geq 0$ for every non-negative $f \in C^\infty(M)$ that is zero on ∂M .



Bibliography

- [Con71] Robert Connelly. “A New Proof of Brown’s Collaring Theorem”. In: *Proceedings of the American Mathematical Society* 27.1 (1971), pp. 180–182. ISSN: 00029939, 10886826. URL: <http://www.jstor.org/stable/2037284> (visited on 04/29/2026).
- [Lee06] John M. Lee. *Introduction to smooth manifolds*. Graduate texts in mathematics 218. New York: Springer, 2006. ISBN: 0387954481. URL: <http://www.gbv.de/dms/goettingen/347921388.pdf>.