

12 Specialist Mathematics Summarised Notes  
(Unofficial)  
Work in Progress

Functions

Composite Functions

Given  $f : x \mapsto f(x)$  and  $g : x \mapsto g(x)$ , the composite function of  $f$  and  $g$  is:

$$(f \circ g)(x) = f(g(x))$$
  
or  
$$f \circ g : x \mapsto f(g(x))$$

In general,  $(f \circ g)(x) \neq (g \circ f)(x)$ .

Inverse Functions

An inverse function returns the original value from the output of a function.  $f(x)$  has an inverse if it is injective (one-to-one), if  $f(a) = f(b)$  only when  $a = b$ ,  $\therefore$  passes the horizontal line test.

For  $f^{-1}(x)$ , the inverse of  $f(x)$ :

- Is a reflection of  $y = f(x)$  over  $y = x$ .
- $(f \circ f^{-1})(x) = (f^{-1} \circ f)(x) = x$
- Domain of  $f^{-1}$  = range of  $f$ .
- Range of  $f^{-1}$  = domain of  $f$ .

Self-Inverse Functions

An invertible function that is symmetrical about  $y = x$ .

$$f^{-1}(x) = f(x)$$

Reciprocal Functions

A function of the form  $f(x) = \frac{k}{x}$ , where  $k \neq 0$  is a constant.

Reciprocal of Other Functions

The reciprocal of a function  $f(x)$  is  $\frac{1}{f(x)}$ . Graphing  $y = \frac{1}{f(x)}$  from  $y = f(x)$ :

- Zero  $f(x) \rightarrow$  vertical asymp  $\frac{1}{f(x)}$
- Vertical asymp  $f(x) \rightarrow$  zero  $\frac{1}{f(x)}$
- Local max  $f(x) \rightarrow$  local min  $\frac{1}{f(x)}$
- Local min  $f(x) \rightarrow$  local max  $\frac{1}{f(x)}$
- When  $f(x) > 0$ ,  $\frac{1}{f(x)} > 0$
- When  $f(x) < 0$ ,  $\frac{1}{f(x)} < 0$
- When  $f(x) \rightarrow 0$ ,  $\frac{1}{f(x)} \rightarrow \pm\infty$
- When  $f(x) \rightarrow \pm\infty$ ,  $\frac{1}{f(x)} \rightarrow 0$

Invariant Points:

Points that do not move under a transformation occurring at  $y = \pm 1$ .

Rational Functions

Results from the division of one polynomial by another. Vertical asymptote occurs when denominator is zero. Horizontal asymptote ascertained from behaviour of graph as  $|x| \rightarrow \infty$ .

- If the degree of denominator > numerator, horizontal asymptote at  $y = 0$ .
- If the degree of denominator < numerator, function has slanted asymptote found through polynomial division.
- If the degree of denominator = numerator horizontal asymptote at  $y = \frac{a}{b}$  where  $a$  and  $b$  are the leading coefficients.

Absolute Value Functions

The absolute value or modulus  $|x|$  of a real number  $x$  is its distance from 0 on the number line.

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Alternatively,

$$|x| = \sqrt{x^2}$$

Properties:

- $|x| \geq 0$
- $|x|^2 = x^2$
- $\left|\frac{x}{y}\right| = \frac{|x|}{|y|}$
- $|-x| = |x|$
- $|xy| = |x||y|$
- $|x - y| = |y - x|$

If  $|x| = a$  where  $a > 0$ , then  $x = \pm a$ . If  $|x| = |b|$  then  $x = \pm b$ .

Graphs Involving the Absolute Value Function

Graphing  $y = f(|x|)$  from  $y = f(x)$ :

- Discard the graph for  $x < 0$
- Reflect the graph for  $x \geq 0$  in the  $y$ -axis
- Points on the  $y$ -axis are invariant

Graphing  $y = |f(x)|$  from  $y = f(x)$ :

- Keep the graph for  $f(x) \geq 0$
- Reflect the graph for  $f(x) < 0$  in the  $x$ -axis
- Points on the  $x$ -axis are invariant

Trigonometric Identities

Angle Relationships

$$\begin{aligned} \sin(-\theta) &= -\sin \theta & \cos(-\theta) &= \cos \theta \\ \sin(\pi - \theta) &= \sin \theta & \cos(\pi - \theta) &= -\cos \theta \\ \sin\left(\frac{\pi}{2} - \theta\right) &= \cos \theta & \cos\left(\frac{\pi}{2} - \theta\right) &= \sin \theta \end{aligned}$$

Pythagorean Theorem

$$\begin{aligned} \sin^2 \theta + \cos^2 \theta &= 1 \\ \tan^2 \theta + 1 &= \sec^2 \theta \\ \cot^2 \theta + 1 &= \csc^2 \theta \end{aligned}$$

Double Angle Identities

$$\begin{aligned} \sin 2\theta &= 2 \sin \theta \cos \theta \\ \cos 2\theta &= \cos^2 \theta - \sin^2 \theta \\ &= 1 - 2 \sin^2 \theta \\ &= 2 \cos^2 \theta - 1 \\ \tan 2\theta &= \frac{2 \tan \theta}{1 - \tan^2 \theta} \end{aligned}$$

Angle Sum and Difference

$$\begin{aligned} \sin(A \pm B) &= \sin A \cos B \pm \cos A \sin B \\ \cos(A \pm B) &= \cos A \cos B \mp \sin A \sin B \\ \tan(A \pm B) &= \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \end{aligned}$$

Sum to Product

$$\begin{aligned} \sin A \pm \sin B &= 2 \sin\left(\frac{A \pm B}{2}\right) \cos\left(\frac{A \mp B}{2}\right) \\ \cos A + \cos B &= 2 \cos\left(\frac{A + B}{2}\right) \cos\left(\frac{A - B}{2}\right) \\ \cos A - \cos B &= -2 \sin\left(\frac{A + B}{2}\right) \sin\left(\frac{A - B}{2}\right) \end{aligned}$$

Product to Sum

$$\begin{aligned} 2 \sin A \cos B &= \sin(A + B) + \sin(A - B) \\ 2 \sin A \sin B &= \cos(A - B) - \cos(A + B) \\ 2 \cos A \cos B &= \cos(A + B) + \cos(A - B) \end{aligned}$$

Mathematical Induction

The Principle of Mathematical Induction

Suppose  $P_n$  is a proposition which is defined for every integer  $n \geq a$ ,  $a \in \mathbb{Z}$ . If  $P_a$  is true, and if  $P_{k+1}$  is true whenever  $P_k$  is true, then  $P_n$  is true for all  $n \geq a$ .

Complex Numbers

Imaginary Numbers

A number which cannot be placed on a real number line in the form  $ai$  where  $a \in \mathbb{R}$  and  $i = \sqrt{-1}$ .

Complex Numbers

Any number in the form  $a + bi$  where  $a, b \in \mathbb{R}$  and  $i = \sqrt{-1}$ .

If  $z = a + bi$   
 $\Re(z) = a \quad \Im(z) = b$

The Complex Plane

Complex numbers can be plotted on the complex plane or Argand plane as a vector where the  $x$ -axis is the real axis and the  $y$ -axis is the imaginary axis.

$\overrightarrow{OP} = \begin{pmatrix} x \\ y \end{pmatrix}$  represents  $x + yi$

Complex Conjugates

The complex conjugate of

$z = a + bi$  is  $z^* = a - bi$

In the complex plane,  $z^*$  is the reflection of  $z$  in the real axis.

Modulus and Argument

The modulus of the complex number  $z = a + bi$  is the length of the vector  $\begin{pmatrix} a \\ b \end{pmatrix}$ , which is the real number:

$|z| = \sqrt{a^2 + b^2}$

The argument of  $z$ ,  $\arg(z)$  is the angle  $\theta$  between the positive real axis and  $\begin{pmatrix} a \\ b \end{pmatrix}$ . Real numbers have an argument of 0 or  $\pi$ . Purely imaginary numbers have argument of  $\frac{\pi}{2}$  or  $-\frac{\pi}{2}$ .

Properties of Modulus:

- $|z^*| = |z|$

- $|z^*|^2 = zz^*$
- $|z_1 z_2| = |z_1| |z_2|$
- $\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}, z_2 \neq 0$
- $|z_1 z_2 z_3 \dots z_n| = |z_1| |z_2| |z_3| \dots |z_n|$
- $|z^n| = |z|^n, n \in \mathbb{Z}^+$

Polar Form

$\text{cis } \theta = \cos \theta + i \sin \theta$

A complex number  $z$  has the polar form

$z = |z| \text{cis } \theta$

where  $\theta = \arg(z)$ .  
The conjugate of  $z$  is:

$z^* = |z| \text{cis } (-\theta)$

Properties of cis  $\theta$ :

- $\text{cis } \theta \times \text{cis } \phi = \text{cis } (\theta + \phi)$
- $\frac{\text{cis } \theta}{\text{cis } \phi} = \text{cis } (\theta - \phi)$
- $\text{cis } (\theta - 2k\pi) = \text{cis } \theta, k \in \mathbb{Z}$

De Moivre's Theorem

$(|z| \text{cis } \theta)^n = |z|^n \text{cis } n\theta, \text{ for all } n \in \mathbb{Q}$

Roots of Complex Numbers

The  $n^{\text{th}}$  roots of the complex number  $c$  are the solutions of  $z^n = c$ .

The  $n^{\text{th}}$  Roots of Unity

The  $n^{\text{th}}$  roots of unity are the solutions of  $z^n = 1$ .

Distances in the Complex Plane

If  $z_1 \equiv \overrightarrow{OP_1}$  and  $z_2 \equiv \overrightarrow{OP_2}$  then  $|z_1 - z_2|$  is the distance between points  $P_1$  and  $P_2$ .

Real Polynomials

Zeros and Roots

A zero of a polynomial is a value of the variable which makes the polynomial equal to zero.  
 $\alpha$  is a zero of polynomial

$P(x) \iff P(\alpha) = 0$

The roots of a polynomial equation are the solutions to the equation.  
 $\alpha$  is a root (or solution) of

$P(x) \iff P(\alpha) = 0$

The roots of  $P(x) = 0$  are the zeros of  $P(x)$  and the  $x$ -intercepts of the graph  $y = P(x)$

Factors

$(x - \alpha)$  is a factor of the polynomial  $P(x) \iff$  there exists a polynomial  $Q(x)$  such that  $P(x) = (x - \alpha)Q(x)$ .

Polynomial Equality

Two polynomials are equal if and only if they have the same degree (order) and corresponding terms have equal coefficients.

Polynomial Division by Linears

If  $P(x)$  is divided by  $D(x) = ax + b$  until a quotient  $Q(x)$  and constant remainder  $R$  is obtained, then

$\frac{P(x)}{ax + b} = Q(x) + \frac{R}{ax + b}$

Notice that  $P(x) = Q(x) \times (ax + b) + R$ .

Polynomial Division by Quadratics

If  $P(x)$  is divided by  $D(x) = ax^2 + bx + c$ , then

$\frac{P(x)}{ax^2 + bx + c} = Q(x) + \frac{ex + f}{ax^2 + bx + c}$

where  $ex + f$  is the remainder.

The Remainder Theorem

When a polynomial  $P(x)$  is divided by  $x - k$  until a constant remainder  $R$  is obtained, then  $R = P(k)$ .

The Factor Theorem

For any polynomial  $P(x)$ ,  $k$  is a zero of  $P(x) \iff (x - k)$  is a factor of  $P(x)$ .

The Fundamental Theorem of Algebra

If  $P(x)$  is a polynomial of degree  $n$ , then  $P(x)$  has  $n$  zeros, each in the form  $a + bi$  where  $a, b \in \mathbb{R}$ , some of which may be repeated.

Vectors

Vectors in Space

Any point  $P$  in space can be specified  $(x, y, z)$  corresponding to steps in the  $X$ ,  $Y$  and  $Z$  direction from the origin  $O$ .  
The position vector of  $P$  is

OP→ = (x y z) = x i + y j + z k

where i = (1 0 0), j = (0 1 0), and k = (0 0 1), the base unit vectors.

The Magnitude of a Vector

The magnitude or length of the vector a = (a1 a2 a3) is

|a| = sqrt(a1^2 + a2^2 + a3^2)

Operations with Vectors

If a = (a1 a2 a3) and b = (b1 b2 b3) then:

-a = (-a1 -a2 -a3) a+b = (a1+b1 a2+b2 a3+b3) a-b = (a1-b1 a2-b2 a3-b3) ka = (ka1 ka2 ka3)

Vector Algebra

- a + b = b + a
- (a + b) + c = a + (b + c)
- a + 0 = 0 + a = a
- a + (-a) = (-a) + a = 0
- k(a + b) = ka + kb
- |ka| = |k||a|
- If a + b = c, then a = c - b
- If ka = b, k ≠ 0, then a = 1/k b

Vector Between Two Points

If A(a1, a2, a3) and B(b1, b2, b3) then the position vector of B relative to A is

AB→ = OB→ - OA→ = (b1-a1 b2-a2 b3-a3)

The distance from A to B is |AB→| = sqrt((b1-a1)^2 + (b2-a2)^2 + (b3-a3)^2)

Parallelism

a = kb ⇔ a and b are non-zero parallel vectors.

Collinear Points

A, B and C are collinear if AB→ = kBC→.

Unit Vectors

The unit vector v-hat, a vector of length 1 in the direction of v is

v-hat = 1/|v| v

Dot Product (Scalar Product)

If a = (a1 a2 a3) and b = (b1 b2 b3), then the scalar dot product is defined as

a · b = a1b1 + a2b2 + a3b3

Properties

- a · b = b · a
- a · a = |a|^2
- a · (b + c) = a · b + a · c
- (a+b) · (c+d) = a · c + a · d + b · c + b · d
- k(a · b) = (ka) · b = a · (kb), k ∈ R

The Angle Between Two Vectors

The angle θ between two vectors a and b can be found using

cos θ = (a · b) / (|a||b|)

Scalar Product Geometric Properties

- For non-zero vectors a and b: a · b = 0 ⇔ a and b are perpendicular.
- |a · b| = |a||b| ⇔ a and b are non-zero parallel vectors.
- Given a · b = |a||b| cos θ If θ is acute, cos θ > 0 and so a · b > 0 If θ is obtuse, cos θ < 0 and so a · b < 0

Cross Product (Vector Product)

The vector cross product of a = (a1 a2 a3) and b = (b1 b2 b3) is

a × b = (a2b3 - a3b2 a3b1 - a1b3 a1b2 - a2b1)

Alternatively,

a × b = (i j k a1 a2 a3 b1 b2 b3) = (a2 a3 b2 b3) i - (a1 a3 b1 b3) j + (a1 a2 b1 b2) k

Properties

- a × b is a vector perpendicular to both a and b.
- a × a = 0 for all a.
- a × b = -b × a, ∴ a × b and b × a have equal length but opposite direction.
- a · (b × c) is called the scalar triple product.
- a × (b × c) = a × b + a × c
- (a + b) × (c + d) = a × c + a × d + b × c + b × d

Direction of a × b

|a × b| = |a||b| sin θ

where θ is the angle between a and b.

Area

If a triangle has defining vectors a and b then its area is 1/2 |a × b| units^2. If a parallelogram has defining vectors a and b then its area is |a × b| units^2.

Lines in 2 and 3 Dimensions

r = a + λb, λ ∈ R is the vector equation of the line.

The Shortest Distance From a Point to a Line

A point P is closest to a point R on a line in direction b when PR→ is perpendicular to b.

PR→ · b = 0

## Relationships Between Lines

### Line Classification in 2 Dimensions

- Intersecting: Unique solution
- Parallel: No solutions
- Coincident: Infinite Solutions

### Line Classification in 3 Dimensions

- Lines are coplanar if they lie in the same plane. In this case, they may be intersecting, parallel or coincident.
- Otherwise, they are skew.

### Shortest Distance Between Skew Lines

For two skew lines with vector equations  $\mathbf{r}_1 = \mathbf{a}_1 + \lambda \mathbf{b}_1$  and  $\mathbf{r}_2 = \mathbf{a}_2 + \mu \mathbf{b}_2$ , the shortest distance  $d$  between them is

$$d = \frac{|(\mathbf{a}_1 - \mathbf{a}_2) \cdot (\mathbf{b}_1 \times \mathbf{b}_2)|}{|\mathbf{b}_1 \times \mathbf{b}_2|}$$

.....

## Planes

A plane is a flat surface that extends forever and has zero thickness.

### The Vector Equation of a Plane

$$\mathbf{r} = \mathbf{a} + s\mathbf{v} + t\mathbf{w}$$

- $\mathbf{r}$  is the position vector of any point on the plane.
- $\mathbf{a}$  is the position vector of a known point on the plane.
- $\mathbf{v}$  and  $\mathbf{w}$  are any two non-parallel vectors that are parallel to the plane.
- $s, t \in \mathbb{R}$  are two independent parameters.

### The Normal of a Plane

A vector is normal to a plane if it is perpendicular to all vectors which are parallel to the plane. If  $\mathbf{n}$  is a normal to a plane such as  $\mathbf{n} = \mathbf{v} \times \mathbf{w}$ , an equivalent vector equation of a plane is

$$\mathbf{n} \cdot (\mathbf{r} - \mathbf{a}) \quad \text{or} \quad \mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$$

### The Cartesian Equation of a Plane

If a plane has normal vector  $\mathbf{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$  and passes through  $P(X, Y, Z)$  then it has equation

$$ax + by + cz = aX = bY = cZ = d$$

where  $d$  is a constant.

### Distance Between a Point and a Plane

The distance between a point  $P(x_1, y_1, z_1)$  and the plane  $Ax + By + Cz + D = 0$  is

$$d = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

## Angles in Space

### The Angle Between a Line and a Plane

The acute angle  $\phi$  between a line with direction vector  $\mathbf{a}$  and a plane with normal vector  $\mathbf{n}$  is

$$\phi = \sin^{-1} \left( \frac{|\mathbf{n} \cdot \mathbf{b}|}{|\mathbf{n}| |\mathbf{b}|} \right)$$

### The Angle Between Two Planes

If two planes have normal vectors  $\mathbf{n}_1$  and  $\mathbf{n}_2$  and  $\theta$  is the acute angle between them then

$$\theta = \cos^{-1} \left( \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{|\mathbf{n}_1| |\mathbf{n}_2|} \right)$$

.....

## Row Reduction

Linear systems of equations can be solved using augmented matrices. A general  $3 \times 3$  system has the form:

$$\begin{cases} a_1x + b_1y + c_1z = d_1 \\ a_2x + b_2y + c_2z = d_2 \\ a_3x + b_3y + c_3z = d_3 \end{cases}$$

In augmented matrix form, the system is:

$$\left[ \begin{array}{ccc|c} a_1 & b_1 & c_1 & d_1 \\ a_2 & b_2 & c_2 & d_2 \\ a_3 & b_3 & c_3 & d_3 \end{array} \right]$$

Using row operations it is reduced to the echelon form.

$$\left[ \begin{array}{ccc|c} a & b & c & d \\ 0 & e & f & g \\ 0 & 0 & h & i \end{array} \right]$$

$\therefore hz = i \Rightarrow z = \frac{i}{h}$ ,  $ey + fz = g$  and  $ax + by + cz = d$

.....

## Intersecting Planes

Two planes in space could have the following arrangements:

- Intersecting
- Parallel
- Coincident

Three planes in space could have the following arrangements:

- All coincident
  - Two coincident and one intersecting
  - Two coincident and one parallel
  - Two parallel and one intersecting
  - All parallel
  - All meet at one point
  - All meet in a common line
  - The line of intersection of any two planes is parallel to the third plane.
-

Integration

Indefinite Integrals

$$\int k \, dx = kx + c$$
$$\int x^n \, dx = \frac{1}{n+1}x^{n+1} + c, \, n \neq -1$$
$$\int e^x \, dx = e^x + c$$
$$\int \frac{1}{x} \, dx = \ln|x| + c$$
$$\int \cos x \, dx = \sin x + c$$
$$\int \sin x \, dx = -\cos x + c$$

Integrating  $f(ax + b)$

$$\int (ax + b)^n \, dx = \frac{(ax + b)^{n+1}}{a(n+1)} + c,$$
$$n \neq -1$$
$$\int e^{ax+b} \, dx = \frac{1}{a}e^{ax+b} + c$$
$$\int \frac{1}{ax+b} \, dx = \frac{1}{a} \ln|ax+b| + c$$
$$\int \cos(ax+b) \, dx = \frac{1}{a} \sin(ax+b) + c$$
$$\int \sin(ax+b) \, dx = -\frac{1}{a} \cos(ax+b) + c$$

Inverse Trigonometric Functions

$$\int \frac{1}{\sqrt{1-x^2}} \, dx = \arcsin x + c$$
$$\int -\frac{1}{\sqrt{1-x^2}} \, dx = \arccos x + c$$
$$\int \frac{1}{1+x^2} \, dx = \arctan x + c$$
$$\int \frac{1}{\sqrt{a^2-x^2}} \, dx = \arcsin\left(\frac{x}{a}\right) + c$$
$$\int -\frac{1}{\sqrt{a^2-x^2}} \, dx = \arccos\left(\frac{x}{a}\right) + c$$
$$\int \frac{a}{a^2+x^2} \, dx = \arctan\left(\frac{x}{a}\right) + c$$

Integrating  $\sin^2 x$  and  $\cos^2 x$

Use double-angle identities when integrating.

$$\sin^2 x = \frac{1}{2} - \frac{1}{2} \cos 2x$$
$$\cos^2 x = \frac{1}{2} + \frac{1}{2} \cos 2x$$

Definite Integrals

If  $F(x)$  is the antiderivative of  $f(x)$  where  $f(x)$  is continuous over  $a \leq x \leq b$ , the definite integral is:

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Properties of Definite Integrals

$$\int_a^a f(x) \, dx = 0$$
$$\int_a^b k \, dx = k(b-a)$$
$$\int_a^b f(x) \, dx = -\int_b^a f(x) \, dx$$
$$\int_a^b kf(x) \, dx = k \int_a^b f(x) \, dx$$
$$\int_a^b f(x) \, dx + \int_b^c f(x) \, dx = \int_a^c f(x) \, dx$$
$$\int_a^b [f(x) \pm g(x)] \, dx = \int_a^b f(x) \, dx \pm \int_a^b g(x) \, dx$$

Integration by Substitution

$$\int f(u) \frac{du}{dx} \, dx = \int f(u) \, du$$

When solving definite integrals with bounds  $a$  and  $b$ , adjust as  $u(a)$  and  $u(b)$ .


Integration by Parts

$$\int uv' \, dx = uv - \int u'v \, dx$$

Area Under a Curve

If  $f(x)$  is positive and continuous for  $a \leq x \leq b$ , the area bound by  $y = f(x)$ , the  $x$ -axis,  $x = a$  and  $x = b$  is:

$$A = \int_a^b f(x) \, dx \quad \text{or} \quad A = \int_a^b y \, dx$$




Area Between Two Curves

Upper function:  $f(x)$  or  $y_U$   
Lower function:  $g(x)$  or  $y_L$

$$A = \int_a^b [f(x) - g(x)] \, dx$$

or

$$A = \int_a^b [y_U - y_L] \, dx$$



Solids of Revolution

When the region enclosed by  $y = f(x)$ , the  $x$ -axis,  $x = a$  and  $x = b$  is revolved through  $2\pi$  about the  $x$ -axis, the volume is:

$$V = \pi \int_a^b y^2 \, dx$$

When the region enclosed by  $y = f(x)$ , the  $y$ -axis,  $y = f(a) = c$  and  $y = f(b) = d$  is revolved through  $2\pi$  about the  $y$ -axis, the volume is:

$$V = \pi \int_c^d x^2 \, dy$$

Volumes for Two Defining Functions

Upper function:  $f(x)$  or  $y_U$   
Lower function:  $g(x)$  or  $y_L$

$$V = \int_a^b ([f(x)]^2 - [g(x)]^2) \, dx$$

or

$$V = \int_a^b (y_U^2 - y_L^2) \, dx$$

# Rates of Change and Differential Equations

## Implicit Differentiation

To find  $\frac{dy}{dx}$  for an implicit relationship between  $x$  and  $y$ . A useful property is

$$\frac{d}{dx} [x^n] = nx^{n-1}$$

*and*

$$\frac{d}{dx} [y^n] = nx^{n-1} \frac{dy}{dx}$$

.....

## Related Rates

.....

## Differential Equations

A differential equation is an equation that describes the relationship between a function and its derivatives.

.....

## Differential Equations of the Form $\frac{dy}{dx} = f(x)$

---

## Vector Calculus

### Parametric Equations

.....

### Pairs of Uniformly Varying Quantities

An object initially at  $(x_0, y_0)$  and moving with velocity vector  $\mathbf{v} = \begin{pmatrix} a \\ b \end{pmatrix}$  has parametric equations

$$x(t) = x_0 + at, \quad y(t) = y_0 + bt, \quad t \geq 0$$

The speed of the object is

$$|\mathbf{v}| = \sqrt{a^2 + b^2}$$