Game Theory
Nash Theorem[1]

Christoph Schottmüller

University of Copenhagen

[1] License: CC Attribution Share Alike 4.0

Outline

1. Existence of Nash equilibria
2. Maths: Fixed point theorems
3. Nash theorem

Existence of Nash equilibrium I

- not all games have a Nash equilibrium

Example

- 2 players ($N = \{P1, P2\}$)
- each player says a number ($A_i = \mathbb{R}$)
- player that says the higher number wins
  (e.g. winner has payoff 1 while loser has payoff 0, if both say the same number each has payoff 1/2)

Existence of Nash equilibrium II

Why are we interested in existence?

- worthwhile to search for an equilibrium
- steady state interpretation of equilibrium, existence of NE says that the process might become stable
- sometimes possible to do comparative statics without computing the NE (only sensible if NE exists)

Fixed point theory I

Definition (fixed point)

Let f be a real function, i.e. $f : A \to \mathbb{R}$ where $A \subseteq \mathbb{R}$. A point $a \in A$ is a *fixed point* of f if $f(a) = a$.

- Does $f(x) = 1$ have a fixed point? What about $f(x) = x + 1$? What about $f(x) = x^2$?
- fixed point theory gives general conditions under which functions have fixed points
- fixed points for functions from $f : A \to \mathbb{R}^n$ where $A \subseteq \mathbb{R}^n$ are defined analogously

Fixed point theory II

Theorem (Brouwer's fixed point theorem, 1-dimensional)

Let $f : [0, 1] \to [0, 1]$ be a continuous function. Then f has a fixed point.

Fixed point theory III

Example (2 player Cournot)

- two firms each choose a quantity q_i
- both firms have costs $c(q_i) = cq_i$ for some $c > 0$
- inverse demand is $P(q_1 + q_2)$ where we assume that P is two times continuously differentiable with $P' < 0$ and $P'' \leq 0$
- assume that $P(1) < c$
  $\Rightarrow$ a firm will never offer a quantity greater than 1
- firm 1 chooses a quantity from $[0, 1]$ to maximize profits

$$\max_{q_1} (P(q_1 + q_2) - c) q_1$$

we get the first order condition

$$P'(q_1 + q_2) q_1 + P(q_1 + q_2) - c = 0$$

Fixed point theory IV

Example (2 player Cournot continued)

- the second order condition holds by assumption

$$P''(q_1 + q_2)q_1 + 2P'(q_1 + q_2) < 0$$

- the first order condition defines a best response function $q_1(q_2)$
- the best response function is continuous because P and P' are continuous by assumption
- Brouwer: best response function has a fixed point!
- game is symmetric $\Rightarrow$ fixed point is an equilibrium $\Rightarrow$ equilibrium exists
- We showed this without being able to actually calculate the equilibrium!!!

Fixed point theory V

Theorem (Brouwer fixed point theorem)

Let S be a convex and compact set in $\mathbb{R}^n$ and let $f : S \to S$ be a continuous function. Then there exists an $s^* \in S$ such that $f(s^*) = s^*$.

Nash theorem

Theorem (Nash theorem)

A strategic game $\langle N, (A_i), (u_i) \rangle$ with

- *a finite number of players*
- *a finite number of actions for each player*

has a Nash equilibrium in mixed strategies.

- proof uses Brouwer's fixed point theorem
- to simplify notation, we prove the theorem for 2×2 games

Proof of Nash theorem for 2×2 games I

- 2×2 game:

	L	R
U	a,b	c,d
D	e,f	g,h

- mixed strategy of P1: probability $\alpha \in [0, 1]$ of playing U
- mixed strategy of P2: probability $\beta \in [0, 1]$ of playing L
- idea of proof:
  - define a function $f(\alpha, \beta)$ such that
    - f is continuous
    - if $f(\alpha^*, \beta^*) = (\alpha^*, \beta^*)$, then (α^*, β^*) is a Nash equilibrium of the game
  - use Brouwer's theorem to establish that f has a fixed point

Proof of Nash theorem for 2×2 games II

- $u_1(U, \beta) = \beta a + (1 - \beta)c$ is the expected utility of P1 when playing U and P2 uses the mixed strategy β
- $u_1(U, \beta)$ is linear and therefore continuous in β
- define

$$g(\alpha, \beta) = max\left\{0, \frac{\alpha + u_1(U, \beta) - u_1(D, \beta)}{1 + |u_1(U, \beta) - u_1(D, \beta)|}\right\}$$

  - $g(\alpha, \beta)$ is higher than α if U is the best response to β and lower than α if D is best response
  - g is continuous because u_1 is continuous in β
- define

$$h(\alpha, \beta) = max\left\{0, \frac{\beta + u_2(L, \alpha) - u_2(R, \alpha)}{1 + |u_2(L, \alpha) - u_2(R, \alpha)|}\right\}$$

Proof of Nash theorem for 2×2 games III

- $h(\alpha, \beta)$ is higher than β if L is best response to α and lower than β if R is best response
- h is continuous because u_2 is continuous in α

- let f be

$$f(\alpha, \beta) = (g(\alpha, \beta), h(\alpha, \beta))$$

- if $f(\alpha^*, \beta^*) = (\alpha^*, \beta^*)$ then
  - $g(\alpha^*, \beta^*) = \alpha^* \Rightarrow \alpha^*$ is best response to β^*
  - $h(\alpha^*, \beta^*) = \beta^* \Rightarrow \beta^*$ is best response to α^*
  - (α^*, β^*) is Nash equilibrium

  every fixed point of f is Nash equilibrium
- f is continuous because g and h are continuous
- Brouwer: f has a fixed point

Existence revisited I

Example (revisited)

- 2 players ($N = \{P1, P2\}$)
- each player says a number ($A_i = \mathbb{R}$)
- player that says the higher number wins
  (e.g. winner has payoff 1 while loser has payoff 0, if both say the same number each has payoff 1/2)

- why does the Nash theorem not apply?

Existence revisited II

Note that the propositions give sufficient conditions for NE, i.e. NE can exist even if the conditions are not satisfied:

Example (war of attrition 1, OR ex. 18.4)

- 2 players dispute an object
- time is a continuous variable starting at 0 and running indefinitely
- actions: chose a time a_i when to concede object
- player i attaches value v_i to object and loses one payoff unit for every time unit (until one of the players concedes)

action space is not finite

Review Questions

- Why are we interested in the question "Does a Nash equilibrium exist?"?
- What is a "fixed point"?
- Explain Brouwers fixed point theorem (1-dimensional case)!
- What does Nash's theorem state?

reading: MSZ 5.3, (supplementary *reading: OR 2.4)

Exercises I

1. We look at a 2 player game in which $A_1 = A_2 = [0, 1]$. Assume that P1 has a best response function that is strictly increasing and continuous. For all subquestions draw the best response functions in an a_1, a_2 diagram.
   1. First, assume that P2 also has a best response function that is continuous. Does this guarantee the existence of a Nash equilibrium?
   2. Second, assume that P2 also has a best response function that is increasing (though not necessarily continuous). Does this guarantee the existence of a Nash equilibrium?
   3. Third, assume that P2 has a best response function that is discontinuous at one point only. Does this guarantee the existence of a Nash equilibrium?
2. *Use the Brouwer fixed point theorem to prove that a pure strategy Nash equilibrium exists in the following strategic game:
   - $N = \{P1, P2\}$

Exercises II

- $A_1 = A_2 = [0, 1]$
- $u_1(a_1, a_2)$ is differentiable and strictly concave in a_1. The partial derivative $\frac{\partial u_1}{\partial a_1}(a_1, a_2)$ is continuous in both variables.
- $u_2(a_1, a_2)$ is differentiable and strictly concave in a_2. The partial derivative $\frac{\partial u_2}{\partial a_2}(a_1, a_2)$ is continuous in both variables.

Hint: This exercise is an extension of the Cournot example we did in the lecture. The argument is similar.