Calculus Problem Book:
Form V and Form VI

S. Chu

Contents

Preface 7

Chapter 1. Functions and Other Miscellaneous Content 9
1. Set Notation and Numbers 9
2. Working with Inequalities 10
3. Polynomials & Rational Functions 10
4. Trigonometric Functions 11
5. Miscellaneous 11

Chapter 2. Limits and Continuity 13
1. Intuitive Notion 13
2. Epsilon and Delta 13
3. Continuity 13
4. Intermediate Value Theorem 13

Chapter 3. The Derivative 15
1. Limit Definition and Properties of the Derivative 15
2. Tangent Line Problem 15
3. Higher Order Derivatives 15
4. Power Rule and others 15
5. Implicit Differentiation 15

Chapter 4. Applications of the Derivative 17
1. Related Rates 17
2. Mean Value Theorem 17
3. Maximums and Minimums 17
4. Optimization and Graphing 17

Chapter 5. The Integral 19
1. Series 19
2. Riemann Sums 19
3. Riemann Integrable 19
4. Properties of the Integral 19

Chapter 6. The Fundamental Theorem of Calculus 21
1. Anti-Derivatives 21
2. FTC part I 21
3. FTC part 2 21
4. Functions Defined by an Integral 21
5. Integral as Accumulator 21
6. U Substitution 21

7. Probability Distributions 21

Chapter 7. Logarithms, Exponentials, and Inverses 23
1. Logarithms 23
2. Inverse Functions 23
3. Exponentials 23
4. Inverse Trigonometric Functions 24
5. Hyperbolic Trigonometric Functions 24

Chapter 8. Methods of Integration 25
1. Integration by Parts 25
2. Method of Partial Fractions 25
3. Trigonometric Substitutions 25
4. Improper Integrals 25

Chapter 9. Differential Equations 27
1. Slope Fields and Euler's Method 27
2. Separation of Variables 27
3. Modeling 27
4. Logistic Growth 27
5. Systems of Differential Equations 27
6. Reduce 2nd order equations to two 1st order equations 27

Chapter 10. More Applications of Integration 29

Chapter 11. Infinite Series 31

Chapter 12. Conics, Parametric Equations, Polar Coordinates 33

Chapter 13. Vectors and the Geometry of $\mathbb{R}^3$ 35

Chapter 14. Vector Valued Functions 37

Chapter 15. The Partial Derivative and Function of Several Variables 39

Chapter 16. The Gradient and the Method of Lagrange Multipliers 41

Chapter 17. The Derivative of maps from $\mathbb{R}^n \to \mathbb{R}^m$ 43

Chapter 18. Longer Problem Sets 45
Working with Parameters and the Derivative 46
An Algorithm to Find Roots 47
Instantaneous Rate of Turning 49
Efficient Foraging 50
When is Venus Bright in the Sky? 54
Rainbows 57
Constructing the Demand Curve 60
The Shape of Bee Hive Cells 62
The Art Gallery Problem 64
Getting to π using Integration by Parts 66
Predicting Peak Oil 67
SIR: A Model for the Spread of Infectious Disease 69

Modeling Air Resistance 71
A Model for Combat 73
A Model for Relationships 74
Unbounded Integrals and $\sqrt{\pi}$ 75
A Derivation of the Least Squares Regression Line 76

Chapter 19. Answers 77

Bibliography 79

Preface

The following short booklet contains a collection of problems with solutions that roughly follow the topics typically covered in the calculus sequence for Form V and Form VI mathematics classes at Ethical Culture Fieldston School. The hope is to provide a book of problems of varying difficulty that will provide non-routine exercises for students learning Calculus, as well as to record general themes and approaches to tackling these types of problems. Ideally these problems can be useful to all courses that cover these topics. Problems are roughly categorized by difficulty level and notated via *, **, or *** in increasing difficulty.

At the end of text we have included several longer problems that allow students to tackle more complex problems that often synthesize material from several areas as well as giving students a look at some applications of Calculus to other fields. We have adopted problems from memory, and from years of working with the material. The following references were consulted for problems as well as other considerations such as narrative structure in introducing various topics and thematic development of the material. [FGI96], [Pat04], Spivak [Spi94], [more later].

CHAPTER 1

Functions and Other Miscellaneous Content

1. Set Notation and Numbers

Exercise 1 Sets, Unions and Intersections
Set $A \subset B$ iff every element of A in also in B. Set $A \supset B$ iff every element of B is also in A and two sets are equal iff both $A \supset B$ and $A \subset B$.

(1) If $A \subset B$ then what is the set $A \cap B$?

(2) If $A \subset B$ then what is the set $A \cup B$?

(3) If $A \cup B = B$, give an argument that explains why $A \subset B$.

Exercise 2 Complements
Given a set A, and a set B we can define the complement of that set, A^c or sometimes: $\overline{A}$, A', or $B - A$, as the elements in B but not in A. Notice A^c, A' and $\overline{A}$ do not reference the set B and are often used when no confusion can arise or there is some all encompassing universal set U that contains A.

(1) Given a universal set U and A, B contained in U. Show that $(A \cap B)^c = A^c \cup B^c$ by first showing that $(A \cap B)^c \subset A^c \cup B^c$. Then show that $(A \cap B)^c \supset A^c \cup B^c$. This property is often called DeMorgan's Law.

(2) Describe the elements that are contained in the set $(A - B) \cup (B - A)$. This set is called the symmetric difference of A and B, or $A \Delta B$.

** *Exercise 3 Working with Sets*

In the previous problem we defined the symmetric difference. The following exercises will work through some properties of this operation on sets. Recall that to show two sets A, B are equal you must show that $A \subset B$ ie. every element is A is also in B, as well as that $B \subset A$, ie. every element of B is also in A.

(1) Show that $(A - B) \cup (B - A) = (A \cup B) - (A \cap B)$ as sets. These are two different ways to think about the set that is the symmetric difference of two sets $A \Delta B$. Let A, B be arbitrary sets.

(2) Show that $A \Delta B = B \Delta A$ by showing that every element in the set on the left is in the set on the right and vice versa.

(3) Show that $A \Delta \varnothing = A$. Where $\varnothing$ is the empty set.

(4) Show that $A \Delta A = \varnothing$.

2. Working with Inequalities

3. Polynomials & Rational Functions

Exercise 4 Polynomials I
Let $f(x) = x^2 - 3$ and $g(x) = x^3 - 2x$. Both f and g are polynomials. You can form new polynomials by adding, subtracting, multiplying, and composing.

(1) Find $(f + g)(x)$

(2) Find $(f - 2g)(x)$

(3) Find $(f \cdot g)(x)$

(4) Find $f \circ g)(x)$

(5) Explain why $(f/g)(x)$ is not a polynomial.

solution 4

* *Exercise 5 Defining Polynomials*
A polynomial f has the general form $f(x) = a_0 + a_1 x + a_2 x^2 + \ldots + a_{n-1} x^{n-1} + a_n x^n$, where $n \in \mathbb{Z}$ and $a_i \in \mathbb{R}$. Show that adding, subtracting, or multiplying two polynomials always results in a polynomial. Explain why this is not the case for division.

solution 5

Exercise 6 Polynomials II
A function f is even if and only if $f(x) = f(-x)$ for all x in its domain. A function is odd if and only if $f(-x) = -f(x)$ for all x in its domain.

(1) Give an example of an even polynomial.

(2) Give an example of an odd polynomial.

(3) Show that if f and g are even functions, then $h(x) = f(x) + g(x)$ is also even.

(4) If f and g are odd functions, what can we say about $h(x) = f(x) + g(x)$? What about $k(x) = f(x) \cdot g(x)$?

* *Exercise 7 Infinity is Not a Number*
Let $F(x) = \dfrac{x}{2 - x}$.

(1) We can decompose any function into its even part and its odd part with the following definitions: $f_e(x) = \dfrac{f(x) + f(-x)}{2}$ and $f_o(x) = \dfrac{f(x) - f(-x)}{2}$. Find and simplify the even and odd parts of $F(x)$.

(2) Graph $F_e(x)$ and $F_o(x)$.

(3) Simplify $F_e(x) + F_o(x)$. How does this relate to $F(x)$?

(4) What is $F_e(-2)$? What is $F_o(-2)$? How doe they relate to $F(-2)$?

(5) Explain why the previous result should make you wary of infinity, or at least wary of this method of decomposing functions into their odd and even parts.

4. Trigonometric Functions

5. Miscellaneous

CHAPTER 2

Limits and Continuity

1. Intuitive Notion

2. Epsilon and Delta

* *Exercise 8 Dirichlet Function*

The Dirichlet function D is define as $D(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & \text{otherwise} \end{cases}$. Two sets of numbers are *not seperable* if it is impossible to find an open interval around an element from one set that contains only elements from that set and not the other. Rational numbers and irrational numbers are not separable. Use this to argue that $\lim_{x \to a} D(x)$ does not exist for any value of a. *solution 8*

*** *Exercise 9 A Strange Function*

Let $f(x) = \begin{cases} 0 & x \in \text{ irrational} \cap [0,1] \\ \frac{1}{q} & x = \frac{p}{q} \text{ in lowest terms}, 0 < x < 1 \end{cases}$

For any $a \in [0,1]$, find the $\lim_{x \to a} f(x)$. Show the limit you calculated is in fact the limit by giving an appropriate delta neighborhood for a given $\epsilon > 0$. Spivak [Spi94] *solution 9*

3. Continuity

* *Exercise 10 A Bounded Function around Zero*

Let $f : \mathbb{R} \to \mathbb{R}$ be a function such that $|f(x)| \leq |x|$ for all x in the domain of f. Prove that f is continuous at $x = 0$.

solution 10

* *Exercise 11 Some Points Don't Move That Much*

Let $f : [0,1] \to [0,1]$ be a function such that $|f(x)| \leq |x|$ for all x in the domain of f. Prove that f is continuous at $x = 0$.

solution 11

4. Intermediate Value Theorem

* *Exercise 12 A Polynomial with Real Roots*

Let $f(x) = (x-a)(x-b) + (x-b)(x-c) + (x-a)(x-c)$ where a, b, c are distinct real numbers. Prove that f must have 2 real distinct zeroes. *solution 12*

* *Exercise 13 Fixed points*

Let f be a continous function from the unit interval to itself. $f : [0, 1] \to [0, 1]$. Show that f must have at least one fixed point x_0. In other words, $f(x_0) = x_0$.

** *Exercise 14 Antipodes*

Two points on a sphere are antipodal if the line connecting them passes through the center of the sphere. They are 'directly opposite' one another, or maximally distant from one another while still being on the sphere.

(1) If you assume that the earth is a sphere and that temperature varies continuously from point to point on the earth, show that there must be a pair of antipodal points that are the same temperature at a given moment in time.

(2) If we also assume that air pressure varies continuously across the earth, show that there must be a pair of antipodal points on the surface of the earth that have the same temperature and air pressure at a given moment in time.

solution 14

CHAPTER 3

The Derivative

1. Limit Definition and Properties of the Derivative

Exercise 15 *Introducing a Discontinuity*

Let $f(x) = |x|$. Use the limit definition of the derivative to show that $f'(x) = \dfrac{|x|}{x}$. Show using the definition of continuity how the derivative


2. Tangent Line Problem

3. Higher Order Derivatives

4. Power Rule and others

* *Exercise 16* *Unit Circle and Sine*

Let $P = (\cos\theta, \sin\theta)$. Use similar triangles to calculate an approximation for $\dfrac{\Delta y}{\Delta \theta}$ and explain how this shows what the derivative of $f(x) = \sin(x)$ is. Do the same for $g(x) = \cos(x)$ by looking at $\dfrac{\Delta x}{\Delta \theta}$.



solution 16

5. Implicit Differentiation

CHAPTER 4

Applications of the Derivative

1. Related Rates

2. Mean Value Theorem

3. Maximums and Minimums

4. Optimization and Graphing

CHAPTER 5

The Integral

1. Series

2. Riemann Sums

3. Riemann Integrable

4. Properties of the Integral

CHAPTER 6

The Fundamental Theorem of Calculus

1. Anti-Derivatives

2. FTC part I

3. FTC part 2

4. Functions Defined by an Integral

5. Integral as Accumulator

6. U Substitution

7. Probability Distributions

CHAPTER 7

Logarithms, Exponentials, and Inverses

1. Logarithms

Exercise 17 *A Better Antiderivative*
We defined the function $L(x) := \int_1^x \frac{1}{t} dt$.

(1) This definition combined with FTC part 2 gives us what relationship between $L(x)$ and the function $f(x) = \frac{1}{x}$.

(2) Calculate the derivative of $g(x) = \ln|x|$.

(3) Explain why part a and part b do not contradict the theorem that tells us if a function f has antiderivatives F and G that F and G must differ by a constant.

solution 17

Exercise 18

** *Exercise 19* *A Nice Bound for Log*
Show that for any $0 < x < y$, we have the inequality:

$$\frac{1}{y} < \frac{\ln(y) - \ln(x)}{y - x} < \frac{1}{x}.$$

solution 19

2. Inverse Functions

3. Exponentials

** *Exercise 20* x^y *vs* y^x
Let $f(x) = \frac{\ln(x)}{x}$

(1) Determine on what intervals f is strictly increasing versus strictly decreasing.

(2) Use part (a) to show that if $e \leq A < B$ then $A^B > B^A$.

(3) Use part (b) to show that $e^\pi > \pi^e$.

solution 20

4. Inverse Trigonometric Functions

5. Hyperbolic Trigonometric Functions

CHAPTER 8

Methods of Integration

1. **Integration by Parts**
2. **Method of Partial Fractions**
3. **Trigonometric Substitutions**
4. **Improper Integrals**

CHAPTER 9

Differential Equations

1. Slope Fields and Euler's Method

2. Separation of Variables

3. Modeling

4. Logistic Growth

5. Systems of Differential Equations

6. Reduce 2nd order equations to two 1st order equations

CHAPTER 10

More Applications of Integration

CHAPTER 11

Infinite Series

CHAPTER 12

Conics, Parametric Equations, Polar Coordinates

CHAPTER 13

Vectors and the Geometry of $\mathbb{R}^3$

CHAPTER 14

Vector Valued Functions

CHAPTER 15

The Partial Derivative and Function of Several Variables

CHAPTER 16

The Gradient and the Method of Lagrange Multipliers

CHAPTER 17

The Derivative of maps from $\mathbb{R}^n \to \mathbb{R}^m$

CHAPTER 18

Longer Problem Sets

Working with Parameters and the Derivative

In this exploration we will work with derivative to find tangent lines to a curve. We will also introduce working with parameters and see how this gives us a set of tools to solve more complex problems. We will end by using our skill with parameters to explore a little bit of the early history of Calculus by taking a look at Descartes' Method of Normals and Fermat's Method of Adequality.

First Let's Look at Tangent Lines to a Parabola.

(1) Let $f(x) = x^2 + 1$ and point $C(0, -2)$. Find the equation of the two lines that pass through C and are tangent to f.

(2) Let C vary its position along the y-axis. Let $C(0, k)$. Find the equations of the two lines that pass through C and are tangent to f. Notice your solutions will be in terms of the parameter k.

(3) Lastly, let's generalize and let C be any point in the plane $C(h, k)$. What are the equations of the two lines that pass through C and are tangent to f, in terms of the parameters h and k?

(4) Notice this gives you the equations for the lines through any point in the plane that are tangent to the given parabola. Demonstrate the ease with which you can find these lines that are tangent to the parabola, by finding them for $P(5, 3)$ and $Q(3, -5)$.

(5) What do these equations tell you about the family of lines tangent to the graph of f? What points on the plane have no lines through them that are tangent to the curve? How can you see this from the equations? Explain how this is different than just looking at $f'(x)$?

Descartes' Method Of Normals versus Fermat's Method. In this next section we will take a look at two early methods of finding slopes of lines tangent to a a curve.

An Algorithm to Find Roots

We will learn how to find the zeroes of a function $f(x)$ using Newton's Method, which involves using the derivative to find lines tangent to the curve and using these to successively better approximate the zeros of a function.

(1) First recognize that finding the zero of a line is straightforward. We will reduce finding the zeroes of an arbitrary differentiable function to finding the zeroes of a bunch of lines. Let $y - a = m(x - b)$ be a line through (b, a) with slope m. Find its x-intercept in terms of a, b and m.

(2) We can get a sense for how the method works by looking at a quadratic $f(x) = -x^2 + 5x$.
   (a) First let's solve for the zeroes by factoring or using the quadratic formula. What are the zeroes of f?

   (b) Let's use the point $(4, 4)$ on f, and call our initial x value $x_0 = 4$. Find the line tangent to f at $(4, 4)$. Let's call this line l_1.

   (c) What is the x-intercept of l_1? Call the x-coordinate of the zero of l_1, x_1.

   (d) Notice x_1 is not equal to a zero of f, but it is closer than x_0. Let's repeat this process but using $(x_1, f(x_1))$.

   (e) What is the line tangent to f at $(x_1, f(x_1))$? Find its x-intercept and call it x_2. It should be close to one of the zeroes at this point.

   (f) Give a sketch of each tangent line that you found in this problem and its x-intercept. Detail how this process works to approximate the zeroes of a function.

   (g) If your initial guess was $x_0 = 3$ Would the process converge to $x = 5$ in more or fewer steps. Explain without carrying out the calcuations.

   (h) What initial guess would cause the process to fail? Explain why.

(3) The power of Newton's Method is that the same process can be applied to any differentiable function to find its zeroes. One of its drawbacks is that it can involve a lot of calculation. Let's offload some of this to a spreadsheet, but first let's develop an expression to generate the sequence of x-coordinates, $\{x_0, x_1, x_2, \dots\}$ that should converge to the a zero of the given function.

(a) Let x_0 be your first guess at a zero. Then as we saw before we are looking for the tangent line to $f(x)$ that passes through $x_0, f(x_0)$. x_1 is the x-intercept of that tangent line. Find an expression for x_1 in terms of $x_0, f(x_0), and f'(x_0)$. Generalize this to an expression that gives x_n.

(b) Look at the spreadsheet linked here.

(c) Recall how difficult it is to find where the sine curve intersects with various lines. We will find when $y = 2\sin(x)$ meets $y = x$ by letting $g(x) = 2\sin(x) - x$ and solving for its zeroes.

(d) Use the IVT to get a guess for where $g(x)$ has zeroes. Then use the spreadsheet to find these zeroes with a reasonabe first guess determined by your use of the IVT. Take a screen shot of your spreadsheet that shows the calculations that give each zero.

Instantaneous Rate of Turning

Efficient Foraging

An Application of Optimization[1]

Foraging animals, whose food is arranged in clumps must make a decision about how long to stay in a given clump of food before moving on in search of another. We will try to arrive at an optimal strategy to foraging given a few simplifying assumptions. Let us first assume that patches of food are distributed uniformly (equally spaced) and that they are equally abundant, meaning an animal's optimal time spend in one patch would be the same for any other patch.

(1) Let us first try to model the rate at which resources can be gathered when arriving at a fresh patch.
   (a) What would a sketch of the amount of food gathered versus time look like assuming at time zero that no food had been gathered and the patch has a total of L units of food?
   (b) Explain why your sketch has the shape it has. What physical idea about the patch of food are you trying to capture? What criteria do you think would be common to all proposed models? Explain.
   (c) Does the graph $F(t) = L(1 - e^{-t/2})$ meet the criteria you gave in the previous question? Explain. Do you have a different function in mind? If so compare it to $F(t)$. What are its comparative strengths and weaknesses.


(2) Another factor to take into consideration is how long it takes to get from one patch of food to the next.
   (a) Let's look to extend our model from 1c by assuming it takes A seconds to travel from each patch to a new patch, and that the closest patch to home is also A seconds away. What does the graph of F versus time look like for an animal that leaves home and forages in one patch? Give a labeled sketch.
   (b) Give a piecewise defined function that will model the above situation. Use $f(t) = L(1 - e^{-t/2})$ as the food gathering profile for one patch and A as the time it takes to get to the patch.
   (c) Sketch and give a piecewise defined function that will model F versus time for an animal that leaves home and forages in two patches of food, where every patch is A seconds away and the animal spends B seconds in each patch. It may be easier to define $F(t)$ by using translates of $f(t)$.

(3) Let's take a moment to look at the various characteristics of our model so far.
   (a) What are some of the assumptions we are making in modeling this situation?
   (b) The model gives energy gathered(F) in terms of time spent foraging (t). What are the three other factors or parameters that affect this model?

[1] adapted from Graves at NCSSM


(c) Which of these parameters does the foraging animal have influence on?


(d) What is the same and what is different in the two following foraging strategies?




(4) Our main goal then is to model an animal's foraging behavior in choosing _____ given a certain density of food patches (time between patches), and richness of food patches (ie.energy gathering profile of a patch) so that it will maximize the food it collects in a given time.

(a) Here we have varied the density of the food patches while keeping each patch's food gathering profile the same. For which situation will the animal stay longer in one food patch? Why?






(b) Here we have varied the food gathering profile of each patch, while keeping the density the same. For which situation will the animal stay onlger in one food patch? Why?



(5) Let's get more familiar with these ideas with some concrete models of bees foraging for pollen.

(a) First, let's look at things graphically. Given the travel distance of 3 seconds and the food gathering profile given in the figure below, how much pollen is gathered over 12 seconds, if the bee stays on each flower 1 second versus 3 seconds.



(b) Assuming the food gathering profile is given by $f(t) = \sqrt{t}$, and a travel time of 3 seconds, let's look at the effect of various foraging times (time spent on a flower) on total collected pollen as well as average rate of pollen collected. Fill out the following table.

total time	foraging time	total pollen	avg rate of pollen collection
8	1		
10	2		
12	3		
12	1		
14	4		
15	2		

(c) Which foraging time seems optimal? Why?

(6) It may then be reasonable to assume that animals will choose the foraging time that maximizes the average rate of food gathering. Let $F(t)$ be one cycle of foraging that includes travel time A with a food gathering profile $f(t)$.
So $F(t) = \begin{cases} 0 & , \quad 0 \leq x < A \\ f(t-A) & , \quad A \leq x \end{cases}$

(a) The average rate of food gathering is given by $G(t) = \dfrac{F(t)}{t}$. Assuming $t > 0$, what equation must critical points of $G(t)$ satisfy? What does this look like on a graph of $F(t)$?

(b) Charnov in establishing the marginal value theorem says, "The predator should leave the patch it is presently in when the marginal capture rate in the patch drops to the average capture rate for the habitat." Explain how this is a restatement of what you found in the part a.

(7) Determine the optimal foraging time B for each of the following food gathering profiles and travel times.

(a) $F(t) = \begin{cases} 0 & , \quad 0 \leq t < 3 \\ \sqrt{t-3} & , \quad t \geq 3 \end{cases}$

(b) $F(t) = \begin{cases} 0 & , \quad 0 \leq t < A \\ K\sqrt{t-A} & , \quad t \geq A \end{cases}$

(c) $F(t) = \begin{cases} 0 & , \quad 0 \leq t < 3 \\ \sqrt[3]{t-3} & , \quad t \geq 3 \end{cases}$


(d) $F(t) = \begin{cases} 0 & , \quad 0 \leq t < A \\ \sqrt[n]{t-A} & , \quad t \geq A \end{cases}$

(8) In the previous set of questions what do n, A, and K represent in our foraging model, and how did changing these parameters affect the optimal foraging time? Explain.


When is Venus Bright in the Sky?

We will be looking at when Venus is brightest in the sky. We will make a lot of simplifying assumptions such as circular, coplanar orbits, constant intensity of solar rays and constant albedo of Venus.

(1) First we can look at a general picture of the situation. In the figure below both venus and the earth are orbiting the sun in a counterclockwise direction.



(2) Second, it will be helpful to come up with the fraction of Venus that is illuminated as seen from earth for different angles θ. Call that $P(\theta)$. We can do this by looking at the picture below and asking what portion of the diameter $\overline{FG}$ appears lit in terms of θ.



(3) Explain why we can use the proportion of the diameter that appears lit to represent the proportion of the sphere that appears lit from the perspective of earth. It may help to visualize where *theta* is on the figure above and to the right.

(4) Then make use of the relation that the apparent brightness of an object is inversely proportional to the square of the distance from the source and directly proportional to $P(\theta)$. Let's call apparent brightness B and use k as our constant of proportionality. Write the equation that expresses B in terms of $P(\theta)$ and the distance from the earth to Venus.

(5) Take your expression for B and attempt to simplify the expression so that it is just in terms of the distance between the Earth and Venus.

(6) Lastly find the maximum of your brightness function. What is the Sun-Venus-Earth angle, or distance between Earth and Venus that maximizes the brightness of Venus?


(7) At maximal brightness, what proportion of the surface of Venus appears lit from the perspective of Earth?

(8) Find the maximum value of the Sun-Earth-Venus angle? Use this to explain why Venus is often called the morning star or the evening star.

(9) Some additional resources.
   (a) A simple animation in desmos of the situation with no angles.
   (b) A link to a geogebra animation of the Sun, Venus, Earth System with angles. You need to click the play button by the t parameter to start the animation.
   (c) An interesting view of the path of Venus if we took a geocentric view, which shows the planet moving in retrograde.

Rainbows

First to describe light moving from air into a raindrop and back out we need Snell's Law of refraction, which explains why a straight straw looks bent when put into a glass of water. It is based upon the idea that light takes the path of *least* time to go from point A to point B, and that light travels at different speeds through different media.



The diagram above shows a light ray passing from point A to point B. Its path could potentially meet the air-water boundary at any point on the boundary, but it turns out that the light will actually go through the point P that leads to the path of least time.

(1) Give an expression for t_1 that describes how long it takes to go from point A to point P if light travels at v_1 through air.

(2) Give an expression for t_2 that describes how long it takes to go from point P to point B if light travels at v_2 through air.

(3) Let $T = t_1 + t_2$ and now find $\dfrac{dT}{dx}$.


(4) If there is a local minimum of T at $x = x_0$ then what must be true about x_0?

(5) Does this give you an equation relating $\sin(\theta_1)$ and $\sin(\theta_2)$ to v_1 and v_2?

(6) If light moves faster in air than it does in water, draw a figure that shows a ray of light passing from air to water. Draw the line normal to the boundary and label θ_1 and θ_2.


(7) Draw a figure that shows a ray of light passing from water to air. Again draw the line normal to the boundary and label the angles.

We also need to know how light reflects.



(1) Use the fact that light will 'pick' the point P that minimizes the time it takes to go from A to B. Mimic the previous steps and show that the path of minimal time implies a relation between the angle of incidence (θ_1) and the angle of reflection (θ_2).

Now we look at a raindrop in cross section.



(1) Find the the total angle τ that the light ray turns as it enters the raindrop, is reflected at point B and then exits the light ray. Your expression for τ should be in terms of α and β.

(2) Re-write your expression for τ by using Snell's law that $\dfrac{\sin\alpha}{\sin\beta} = k$ where $k = \dfrac{v_\alpha}{v_\beta}$, so that it is only in terms of the angle β.

(3) Find the angle β that minimizes the total angle deflection τ for $k = 0.75$.

(4) Take a look at your function for τ with $k = .75$ in Desmos. Mark the minimum value of τ and the β that corresponds to it. What do you notice

about the value of τ for all incident angles β around the minimum? If we disregard color for a moment and think of rainbows as just brighter parts of the sky, what does the water droplet do to incoming light rays, and what does this have to do with the minimum of the τ vs. β graph?

(5) $\pi - \tau$ is sometimes called the rainbow angle. If you imagine all the water droplets in the sky acting according to our model of a droplet, can you predict what shape a rainbow should have?

(6) To explain the colors we need to know that the higher the frequency of light, the more it slows down more in water. The ordering of visible light from highest to lowest frequency is violet-blue-green-yellow-orange-red. Draw a diagram of how the colors should separate as a light ray enters a water droplet. What happens to color order after the internal reflection? What about as it exits the droplet?

Constructing the Demand Curve

(1) Here is a quick introduction. The demand curve relates the price of a commodity with the quantity of the commodity that would be demanded at that price. The demand curve could represent one person's quantity demanded versus price, or the entire market (the set of all consumers) for that commodity. The supply curve is a similar relationship between price and the quantity of the commodity that suppliers are willing to sell at that price. The intersection of the supply and demand curved is the market equilibrium price for that commodity.
   (a) Let the demand curve of some good be given by $p = 10 - 2q_d$ and supply be given by $p = 1 + q_s$. What is the market equilibrium price for this good?
   (b) Explain why in general a demand curve will be decreasing, whereas a supply curve is typically increasing?

(2) We will construct the demand curve from the more basic idea of utility. Economic utility can be thought of as a way to measure or compare levels of happiness so that we can talk about consumer preferences by saying one good is preferred because it is worth more utils than another. Let's say we have only two choices of good x, or y, and that our utility function is given by $U(x, y) = \sqrt{xy}$.
   (a) Give a sketch of the curve $1 = \sqrt{xy}$. In the context of the given utility function, what do the points on the curve represent. Explain why this curve is called an indifference curve.
   (b) Sketch the curves $2 = \sqrt{xy}$ and $4 = \sqrt{xy}$. What does this tell you about the difference in the mind of a consumer between the following bundles of goods (x, y): $(1, 1)$; $(1, 4)$; $(1, 16)$. How does this connect with the idea that 'more is always better'?
   (c) Is it possible for two distinct indifference curves to intersect? (hint: remember to apply what you said in part a)

(3) From the utility curve we can construct the marginal utilities for good x by taking $\dfrac{dU}{dx}$ while treating y as constant. The marginal utility curve for good y is calculated in a similar fashion by differentiating with respect to y while holding x constant. Let $U(x, y) = \sqrt{xy}$.
   (a) In words what does the marginal utility function for good x represent?
   (b) Calculate $\dfrac{dU}{dx}$ and $\dfrac{d^2U}{dx^2}$. What does this tell you about U as x increases? What does this tell you about marginal utility as x increases?

(4) To construct a demand curve we will want to know given a limited amount of income, what is the optimal bundle of goods (x, y) where x is the good of interest and y is every other possible good lumped together. So, we need to introduce the budget constraint which relates income I, with purchasing x at p_x and y at p_y.

(a) Say we have income $I = 100$, and $p_x = 2$, $p_y = 5$. Sketch $I = p_x(x) + p_y(y)$. What do the points on this curve represent? Why is this curve typically called the budget curve?

(b) What does $\frac{dy}{dx}$ mean in this context?

(c) If we treat I, p_x, and p_y as constants, find $\frac{dy}{dx}$ in terms of those constants.

(5) Let us now calculate the optimal bundle of goods (x, y) to maximize the utility of the consumer given a budget constraint.

(a) Draw a family of utility curves $U(x, y) = \sqrt{xy}$ by setting $U = \{1, 2, 3, 4\}$ and say your budget curve is given by $5 = x + y$. What is the optimal bundle of goods that you can afford? Explain.

(b) To mimic the above, we will want to find the point on the budget curve that makes the budget curve tangent to an indifference curve. Find the optimal bundle (x^*, y^*) in terms of I and p_x, p_y.

(6) Finally we are able to construct the demand curve. Look at your optimal bundle. Notice it gives a relationship between x^* and p_x. Give a sketch of this for the utility curve $U(x, y) = \sqrt{xy}$. Does the shape of this curve match what you expect from question 1?


(7) Thinking about how we constructed this demand curve, what are the main assumptions that underly its construction?

(8) What questions came up for you? What would you want to further explore?


The Shape of Bee Hive Cells


(1) First let's look at the 2D problem. We will explore why hexagons, but only by restricting to tessellations of the plane by one regular polygon.
   (a) First what are the possible regular polygons that tessellate the plane. Can you prove this?
   (b) Considering your answers from part a, fix the perimeter of each polygon to 1, what is the area of each polygon?
   (c) Fixing the area to 1, what are the respective perimeters of each polygon from part a?
   (d) What can you say about the hexagon compared to the other polygons?

(2) Let's solve a different but somewhat related problem. There are four cities that are located at the four vertices of a square of sidelength 1. How can one connect the four cities with a minimal length of road so that is a path from each city to every other city? The paths can overlap. The two boundary cases are pictured below. The minimum is some intermediary figure. After finding the minimum, how does this relate to the hexagon tessellation?



(3) Moving to 3D, let us first think of the honeycomb cells as two layers of hexagonal prisms of equal heights, and the cell openings are on the top and the bottom. The wall in the middle of the two layers is not planar, and in fact it will take a shape that maintains the volume but minimizes the surface area, thus minimizing the wax required to make the hive. (This is mostly true.)





(a) Let's walk through how to go from the left diagram to the right. First choose a distance, y, for $\overline{JN}$. Notice this defines a pyramid $JKIN$, rotate this pyramid 180° around the line $\overline{IK}$. This gives the rhombus $PKNI$ on the right. Explain why replacing the top base hexagon with 3 congruent rhombi preserves the volume of the honeycomb cell. Let's fix the volume of a cell by choosing s constant as the sidelength of the hexagon and h constant as the height of the hexagonal prism.

(b) Instead of y as the parameter we vary, let's use the parameter θ to vary the shape of the honeycomb. It will be the angle between the vertical line through P and one of the congruent rhombi. Express the area of one rhombus in terms of θ. It will help to picture the pyramid $JKIN$ and that we started with regular hexagons as the base.

(c) Express the area of one of the trapezoidal lateral faces in terms of θ.

(d) Give the expression for the surface area of the cell, and takes its derivative with respect to θ.

(e) What is the value of θ that minimizes the surface area? How do you know it is a minimum?

(4) A possible extension is to look at the tetrahedral molecule Methane (CH_4). Show the angles of the bonds are related to the angles of the honeycomb.

(5) What other observations can you make? Feel free to research online at this point. What else might you be curious about?

The Art Gallery Problem

Understanding the Problem. We are given a gallery in the shape of an arbitrary n-sided polygon. The problem is to find the minimum number of guards needed to guarantee that every part of the gallery is in the line-of-sight of a guard without knowing the exact shape of the n-gon ahead of time. Let's define line-of-sight as points that can be connected to the guard with a line segment that stays within the polygon.

(1) A good way to start this problem is to look at a few galleries and position guards, so that all of the gallery is within sight of a guard.

(2) Notice for some polygons this problem is trivial, for instance how many guards do you need to watch a convex regular hexagon?

(3) However, other polygons can be a little more difficult to visualize. Try to find the minimum number of guards for the following 15-gon.



(4) Do you have a guess as to the minimum number of guards required to handle any n-gon.

A Possible Outline.

(1) Prove any n-gon can be triangulated (cut into non-overlapping triangles) by connecting vertices. (be wary of concave polygons)

(2) Prove that once you have triangulated your polygon, you can 3-color the vertices, meaning assign 1 of 3 colors to each vertex such that no two connected vertices share a color.

(3) Show that your 3-coloring of the polygon allows you to assign the minimal number of guards that will work for any n-gon.

A More Detailed Outline. It will help to first show that any polygon can be triangulated. A polygon is triangulated when it is divided into triangles by drawing diagonals which are formed by connecting pairs of vertices. We will prove this by induction on the number of sides of the polygon.

(1) Why is the base case $n = 3$ true?

(2) The induction step here is a little tricky so we will break it into smaller steps.
   (a) First thinking about our $k + 1$-gon, why must there be at least one convex ($< 180°$) vertex?
   (b) Label that convex vertex Q and the adjacent vertices P and R. If $\overline{PR}$ is a diagonal we are done. Explain.
   (c) If $\overline{PR}$ is not a diagonal that means it must slip outside of the polygon, but then you can argue that there is a diagonal that has Q as one of its endpoints. Explain. Use a diagram.

(3) Your proof that any polygon can be triangulated should be done. Look it over to make sure it makes sense and reads well.

Now we will prove that for any n-gon, $\lfloor n/3 \rfloor$ guards will always be sufficient. First we will prove that any triangulation of a polygon can be 3-colored. For a triangulation to be 3-colored we mean the vertices of each triangle can be assigned a color 1, 2, or 3, so that no connected vertices are the same color.

(1) Prove by induction that any triangulation of an n-gon can be 3-colored.

(2) Suppose a guard is placed at the color that occurs least frequently. First explain why that gives $\lfloor n/3 \rfloor$ guards as the maximum number of guards. Then explain why this ensures all of the gallery is in view of a guard.

(3) You're done! Write out your proof into one coherent argument. Read it one last time to make sure it reads well and makes sense. Be careful not to leave out key ideas.

Getting to π using Integration by Parts

In the following set of exercises you will derive an expression for π that looks very appealing in its separation of odd and even integers.

(1) Let $I_n = \int_0^{\pi} \sin^n(x)dx$. Use integration by parts to find a relationship between I_n and I_{n-2}

(2) Calculuate I_0 and I_1. Use this and the relationship you found inpart part a to find I_k for $k = 1, 2, 3, 4, 5, 6, 7$. Don't simplify by multiplying. What patterns do you see?

(3) Make a guess for the formulas for I_{2n} and I_{2n+1}. Make sure you get the indexing correct here, by checking against part b.

(4) Use mathematical induction to prove the equations for I_{2n} and I_{2n+1} that you guessed in part c.

(5) Explain why $\sin^{2n+1} x \leq \sin^{2n} x \leq \sin^{2n-1} x$ for $x \in [0, \pi]$

(6) Explain why this gives the inequality $I_{2n+1} \leq I_{2n} \leq I_{2n-1}$

(7) Show that $\lim_{n\to\infty} \dfrac{I_{2n}}{I_{2n+1}} = 1$

(8) Show this gives a way to calculate $\dfrac{\pi}{2}$ as an infinite product.

(9) In a spreadsheet calculate the partial products of this infinite product as rows in a table. For each partial product of n terms, calculate its percent error with the actual value of π (at least to 10 decimlas). Then using linearization determine the relationship between the number of terms n in your partial product and the absolute value of the percent error of your partial product with n terms. Does this infinite product converge to π quickly or slowly? Explain.

Predicting Peak Oil

(1) First let's look at our model and make sense of it using what we already know about the logistic model.
   (a) Let $P(t)$ be the total amount of oil the U.S. has extracted at time t in years. Then let us assume the function can be modeled by the differential equation: $\dfrac{dP}{dt} = kP\left(1 - \dfrac{P}{L}\right)$. Give a sketch with two possible solutions curves on a P vs. t axes, clearly labeling any asymptotes.

   (b) Give a sketch of $\dfrac{dP}{dt}$ vs P. Where are its zeroes, and what does that correspond to for the differential equation? Using this graph, make an argument as to the presence and value of any points of inflections for possible solution curves.

   (c) Assuming the solution curve has y-int between 0 and $L/2$, what is the maximal rate of increase of solution curve $P(t)$, in terms of k and L?

   (d) Give a sketch of $\dfrac{1}{P}\dfrac{dP}{dt}$ vs. P. Label any attributes of the curve you can determine in terms of k or L.

   (e) If we now are able to collect values for P at various years and we assume P fits a logistic model, what then can we conclude about the plot of $\dfrac{1}{P}\dfrac{dP}{dt}$ vs P? How does this data help to find a specific solution curve for $P(t)$?

(2) Now onto some actual data.
   (a) On the worksheet labeled US oil, use excel formulas to transform the Oil production from an annual number in thousands of barrels to a cumulative number in billions of barrels.
   (b) In the next column give an approximate for $\frac{dP}{dt}$. It's fine to just use secant approximation.
   (c) Transfer the data from 1931 to present into the US oil worksheet tab. From there use the ideas we developed in question 1, and the linear least squares regression, restricted to 1931-1950) to find values for the parameters of the logistic model. (use time as your x-axis instead of year.)
   (d) Peak oil production (P^*) is the point in time where $\frac{dP}{dt}$ is a maximum. What is the value of P^* that corresponds to peak oil?
   (e) What is the rate of oil production that correspond to the model's peak oil prediction?
   (f) What year does the model predict for peak oil production in the US? (first, make use of the the fact that peak oil P^* can be expressed in terms of L and keep in mind the need to convert from time to year).
   (g) Give the solution curve that we have modeled from the data that describes $P(t)$ with t as time.

(3) Lastly we can now compare the model with actual data.
   (a) Generate the values of P predicted by your solution curve from question 2, for the years 1931 to 2020.
   (b) Plot both your predicted P and the actual values of P on the same axes as scatterplots. What do you see of interest?
   (c) Plot both your predicted $\frac{dP}{dt}$ vs P with the actual $\frac{dP}{dt}$. What do you see of interest?
   (d) This analysis was done by M. King Hubbert in 1956. He predicted peak oil some time between 1965-1970. What is somewhat remarkable is he only used this analysis to solve for k, instead preferring to rely on geologists' best guesses for L. Where does the actual data differ dramatically from the pattern established by the model? What events may have had an influence on oil production in the US to account for these deviations?

(4) Share an observation about the activity. This could be a summary in your own words of what we did, it could be a flaw in the model, it could be something you noticed that you feel is essential to understand the process.

SIR: A Model for the Spread of Infectious Disease

(1) First let's look at our model and make sense of the symbols we will use in the SIR model. Let S be the number of people in the population that are a susceptible to a given disease. I will be the number that are currently infected, and R will be the number who have recovered. We will assume that all infected people are infectious, and that the total population does not change (ie $\dot{S} + \dot{I} + \dot{R} = 0$).
   (a) We will assume the only change to the susceptible population is due to infections, and that the rate at which this happens is proportional to the product of the susceptible and infectious populations. Write the differential equation for $\dot{S}$ with α as the constant of proportionality.
   (b) We will assume that the recovered population increases at a rate proportional to the number of infectious people. Write the differential equation for $\dot{R}$ with β as the constant of proportionality.
   (c) We will assume that the infectious population increases at the same rate the susceptible population decreases and that it decreases at the rate at which the recovered population is increasing. Write the differential equation for $\dot{I}$.

(2) Find a trajectory in phase space
   (a) By using the chain rule for $\frac{dI}{dt}$ where I is a function of S which is a function of t, write out an expression for $\frac{dI}{dS}$.
   (b) Solve the differential equation. Include a picture of the slope field and a solution curve in the first quadrant of an I vs S coordinate system.
   (c) Solve for the constant of integration in terms of S_0 and I_0, and rewrite your solution curve $I(S)$.
   (d) Give a sketch indicating the direction along the solution curve that corresponds to increasing time? How do you know this?
   (e) What is the maximum value of I in terms of S_0, I_0, α, and β? Analytically, how do you know it is a maximum?

(3) The argument for vaccination and the idea of herd immunity.
   (a) First let's scale our slope field by choosing $\beta = 0.1$ and $\alpha = .4$. Now S and I are expressed as percentages of the total population. You can then choose $I_0 = 0.02$ and $S_0 = 0.98$ to represent $2/100$ people be infectious, and the rest of the population being susceptible. Adjust the parameters of the slope field in desmos and take a screenshot of a potential solution curve $I(S)$.
   (b) An outbreak occurs when $\frac{dI}{dt} > 0$. Looking back at our original system of first order differential equations, at what values of S is $\dot{I} > 0$? Let S_c be the value of S where $\frac{dI}{dt}$ changes from positive to negative.

(c) What do you notice about all solution curves $I(S)$ when $S < S_c$?

(d) If $S_0 < S_c$ how would the disease spread among the population?

(e) Vaccination is one way to lower S_0. If 60% of the population is vaccinated, then $S_0 = 0.40$. Using the already given values of α and β what percentage of the population do we need to vaccinate to ensure no outbreak? This is typically called herd immunity.

(f) Assuming the vaccine is only 95% effective, what percentage of the population needs to be vaccinated to achieve herd immunity.

(g) The ratio of $\dfrac{\alpha}{\beta} = R_0$ is called the basic reproductive ratio. It represents the average number of secondary cases that arise from an average primary case in a completely susceptible population. What is R_0 given our parameter values?

(h) The measles have an $R_0 = 15$. What percent vaccination is required to achieve herd immunity given the Measles vaccine has 97

(4) Building $S(t)$, $I(t)$ and $R(t)$ using a spreadsheet and Euler's method.

(a) Use Euler's methods to calculate the next values for S, I, and R in columns E, F, G starting in row 8. Remember it is helpful to fix cell references using F4 so that you can drag formulas down.

(b) Recall in part one you found an expression for the maximum value of I, call it I_m. Calculate this maximum value of I using the parameters in the spreadsheet: $\alpha = 0.4$, $\beta = 0.1$, $S_0 = 0.999$, and $I_0 = 0.001$. Is this the same as the I_m found using Euler's method?

(c) How can you make Euler's method give a better approximation of the curve $I(t)$? Does this decrease the difference in I_m that you found in part b? Explain, and take any relevant screenshots of your spreadsheet.

(d) β is often thought of as the reciprocal of the infectious period of the disease. So if a certain disease has an infectious period of 5 days, then $\beta = 0.2$. This parameter is mostly determined by the particular disease. However, α is parameter we can influence. It is in part a measure of infectiousness of the disease, but it is also a measure of how 'well mixed' a population is. Make a conjecture as to the effect of masking and quarantining on the parameter α. How does this change the curve of infected over time?

(e) In your model decrease α to 0.3. What do you observe?

(5) What do you think is most important that the SIR model does not capture? Explain why you think it is important and how you might incorporate it into the model.

Modeling Air Resistance

> Aristotle says that a hundred pound ball falling from a height of a hundred braccia [arm lengths] hits the ground before a one pound ball has fallen one braccio. I say they arrive at the same time. You find, on making the test, that the larger ball beats the smaller one by two inches. Now, behind those two inches you want to hide Aristotle's ninety-nine braccia and speaking only of my tiny error, remain silent about his enormous mistake.

The above quote is from Galileo after conducting the experiment of dropping two weights from the top of the tower of Pisa. His idea that falling objects accelerate at the same rate was criticized on the basis that the heavier object hit the ground before the lighter one. Galileo did note that this was most likely due to the effect of air resistance. We will take a simple model of air resistance and try to calculate how much sooner the heavier weight hits the ground. We will work in standard metric units and assume the tower is 56 meters high.

(1) First let's derive our models of the velocity of a falling object assuming air resistance is proportional to velocity, or to velocity squared.
   (a) If we assume air resistance and gravity are the only two forces acting on the ball, we can write that $m\dot{v} = mg - kv$. Solve this differential equation using separation of variables. Assume $v_0 = 0$.
   (b) What is the $\lim_{t\to\infty} v(t)$? What is the physical interpretation of this expression?
   (c) Use Desmos to generate a picture of the slope field generated by the above differential equations and a solution curve given $k = 0.08$ and $m = 1$.
   (d) If you ignore air resistance, how long will it take an object to fall 56 meters (approximately the height of the tower)?
   (e) Using the solution curve you generated in part c, how long will it take your 1kg object to fall the 56 meters? (hint: change in position is given by the integral of the velocity curve.) Include a screen shot of your desmos calculation.
   (f) Double the mass of the object. How long does it take now to fall 56 meters? Does this corroborate Galileo's observation? Again include a screenshot of your desmos calculation.
   (g) Just by looking at the differential equation, what would you expect to happen if you reduced the mass to 0.5 kg? How long does it now take to travel 56 meters?

(2) Let us now assume that air resistance is proportional to velocity squared.
   (a) Write the new differential equation, but use a different constant of proportionality.
   (b) Solve the differential equation.
   (c) What is the equlibrium solution for this differential equation? If you wanted both models to have the same equilibrium solution, what should be the relationship between constants?
   (d) What is the specific solution curve given a mass of 1 kg, and a constant that results in the same equilibrium solution from the previous model.
   (e) How long does it take this object to travel 56 meters? Include a screenshot of the desmos calculation.
   (f) As you double and halve the mass what do you observe about the time to travel 56 meters? Include screen shots.
   (g) How do the two models compare? Observations? Conjectures?

(3) not finished . . . [here there will be a data component I just need to actually gather some data]

A Model for Combat

[Still working on it. Based upon Lanchester Laws]

A Model for Relationships

[Systems of Differential Equations. Still developing this one. May require matrices]

Unbounded Integrals and $\sqrt{\pi}$

[The idea here is to show how one way to extend the factorial function to all real numbers involves integrating the logarithm from 0 to 1. Note it gives a value for gamme(1/2) I think as $\sqrt{\pi}$.

A Derivation of the Least Squares Regression Line

Given a set of data in two variables x_i, y_i you might want to see if there

CHAPTER 19

Answers

Exercise 4

blah blah

Exercise 5

The expression $A \subset B$ is read, the set A is contained in the set B. Whereas the expression $A \supset B$ is read the set A contains the set B.

(1) $\mathbb{Z} \supset \mathbb{N}$

(2) $\mathbb{R} \supset \mathbb{Q}$

(3) $\mathbb{N} \subset \mathbb{Q}$

Exercise 6

Exercise 7

Exercise 8

blah blah

Exercise 9

First make a sketch of the function.

Exercise 10

blah blah

Exercise 11

blah blah

Exercise 12

Look at $f(a)$, $f(b)$ and $f(c)$. Since a, b, and c are distinct we can assume they are ordered $a < b < c$, this will then put you in a position to apply IVT.

Exercise 14

Both require looking at the difference in the function evaluated at a pair of antipodes.

Exercise 15

Exercise 16

blah blah

Exercise 17

Exercise 18

Exercise 19

Note that the term bounded by $\frac{1}{y}$ and $\frac{!}{x}$ is the average rate of change of $\ln(x)$. So applying the MVT gives that there is some $c \in [x, y]$ such that $\frac{d}{dx}\ln(x)$ at $x = c$ is equal to the average rate of change of log. Finally use that $y = \frac{1}{x}$ is decreasing.

Exercise 20

After finding f' you can show that f is decreasing on the interval (e, ∞). This implies that since $e < \pi$ we get $f(e) > f(\pi)$. Then multiplying through by $e * \pi$ and taking exponentials base e of both sides gives the result.

Bibliography

[Spi94] Michael Spivak. *Calculus*. Publish or Perish Inc, 1994.

[FGI96] Fomin, Genkin, and Itenberg. *Mathematical Circles (Russian Experience)*. Vol. 7. Mathematical World. American Mathematical Society, 1996.

[Pat04] Patrick. *Introduction to Counting and Probability*. 2nd. AoPS Inc., 2004.