
3D Surface: Site fraction of MOO4-2 on sublattice 2 in LIQUID
Site fraction of MOO4-2 on sublattice 2 in LIQUID
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Site fraction of MOO4-2 on sublattice 2 in LIQUID


Contour: Site fraction of MOO4-2 on sublattice 2 in LIQUID
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Site fraction of MOO4-2 on sublattice 2 in LIQUID
0.90
0.75
0.60
0.45
0.30
0.15
0.00


3D Surface: Site fraction of CS+ on sublattice 1 in LIQUID
Site fraction of CS+ on sublattice 1 in LIQUID
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Site fraction of CS+ on sublattice 1 in LIQUID
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Site fraction of CS+ on sublattice 1 in LIQUID
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
1e−5+9.999e−1
9.50
8.75
8.00
7.25
6.50
5.75
5.00
Site fraction of CS+ on sublattice 1 in LIQUID


3D Surface: Mole fraction of LIQUID
Mole fraction of LIQUID
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Mole fraction of LIQUID
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Mole fraction of LIQUID
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of LIQUID
0.90
0.75
0.60
0.45
0.30
0.15
0.00


3D Surface: Mole fraction of MOO2
Mole fraction of MOO2
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Mole fraction of MOO2
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Mole fraction of MOO2
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of MOO2
0.42
0.36
0.30
0.24
0.18
0.12
0.06
0.00


3D Surface: Mole fraction of CS2MOO4_S2
Mole fraction of CS2MOO4_S2
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
1.0
0.8
0.6
0.4
0.2
0.0
Mole fraction of CS2MOO4_S2


Contour: Mole fraction of CS2MOO4_S2
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of CS2MOO4_S2
0.84
0.72
0.60
0.48
0.36
0.24
0.12
0.00


3D Surface: Mole fraction of CS2MOO4_S1
Mole fraction of CS2MOO4_S1
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
1.0
0.8
0.6
0.4
0.2
0.0
Mole fraction of CS2MOO4_S1


Contour: Mole fraction of CS2MOO4_S1
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of CS2MOO4_S1
0.84
0.72
0.60
0.48
0.36
0.24
0.12
0.00


3D Surface: Mole fraction of BCC_A2
Mole fraction of BCC_A2
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Mole fraction of BCC_A2
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Mole fraction of BCC_A2
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of BCC_A2
0.54
0.48
0.42
0.36
0.30
0.24
0.18
0.12
0.06
0.00


3D Surface: Mole fraction of GAS
Mole fraction of GAS
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Mole fraction of GAS
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Mole fraction of GAS
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of GAS
0.72
0.64
0.56
0.48
0.40
0.32
0.24
0.16
0.08
0.00


3D Surface: Mole fraction of O in GAS
Mole fraction of O in GAS
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Mole fraction of O in GAS
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Mole fraction of O in GAS
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of O in GAS
0.90
0.75
0.60
0.45
0.30
0.15
0.00


3D Surface: Mole fraction of Cs in GAS
Mole fraction of Cs in GAS
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Mole fraction of Cs in GAS
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Mole fraction of Cs in GAS
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of Cs in GAS
0.90
0.75
0.60
0.45
0.30
0.15
0.00


3D Surface: Site fraction of MOO3 on sublattice 1 in GAS
Site fraction of MOO3 on sublattice 1 in GAS
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Site fraction of MOO3 on sublattice 1 in GAS
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Site fraction of MOO3 on sublattice 1 in GAS
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Site fraction of MOO3 on sublattice 1 in GAS
0.24
0.21
0.18
0.15
0.12
0.09
0.06
0.03
0.00


3D Surface: Site fraction of CS2MOO4 on sublattice 1 in GAS
Site fraction of CS2MOO4 on sublattice 1 in GAS
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Site fraction of CS2MOO4 on sublattice 1 in GAS


Contour: Site fraction of CS2MOO4 on sublattice 1 in GAS
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Site fraction of CS2MOO4 on sublattice 1 in GAS
0.90
0.75
0.60
0.45
0.30
0.15
0.00


3D Surface: Mole fraction of Mo in GAS
Mole fraction of Mo in GAS
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
1.0
0.8
0.6
0.4
0.2
0.0
Mole fraction of Mo in GAS


Contour: Mole fraction of Mo in GAS
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mole fraction of Mo in GAS
0.24
0.21
0.18
0.15
0.12
0.09
0.06
0.03
0.00


3D Surface: Cs2MoO4 (gas)
Cs2MoO4 (gas)
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
1.0
0.8
0.6
0.4
0.2
0.0
Cs2MoO4 (gas)


Contour: Cs2MoO4 (gas)
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Cs2MoO4 (gas)
0.315
0.270
0.225
0.180
0.135
0.090
0.045
0.000


3D Surface: MoO3 (gas)
MoO3 (gas)
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
MoO3 (gas)
1.0
0.8
0.6
0.4
0.2
0.0


Contour: MoO3 (gas)
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
MoO3 (gas)
0.00024
0.00021
0.00018
0.00015
0.00012
0.00009
0.00006
0.00003
0.00000


3D Surface: Mo in gaseous phase
Mo in gaseous phase
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mo in gaseous phase


Contour: Mo in gaseous phase
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Mo in gaseous phase
0.09
0.08
0.07
0.06
0.05
0.04
0.03
0.02
0.01
0.00


3D Surface: Cs in gaseous phase
Cs in gaseous phase
1.0
0.8
0.6
0.4
0.2
0.0
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
Chemical potential of O [kJ/mol]
−400
−350
−300
−250
−200
−150
−100
Cs in gaseous phase
1.0
0.8
0.6
0.4
0.2
0.0


Contour: Cs in gaseous phase
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
Cs in gaseous phase
0.45
0.40
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00


3D Surface: MoO4-2 (liquid, sublattice 2)
MoO4-2 (liquid, sublattice 2)
1.0
0.8
0.6
0.4
0.2
0.0
2500
2250
2000
1750
1500
1250
1000
750
500
Temperature [K]
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
1.0
0.8
0.6
0.4
0.2
0.0
MoO4-2 (liquid, sublattice 2)


Contour: MoO4-2 (liquid, sublattice 2)
Temperature [K]
2500
2250
2000
1750
1500
1250
1000
750
500
−400
−350
−300
−250
−200
−150
−100
Chemical potential of O [kJ/mol]
MoO4-2 (liquid, sublattice 2)
0.84
0.72
0.60
0.48
0.36
0.24
0.12
0.00