

Phytoplankton community structure in one sector of Guanabara Bay (RJ, Brazil) during 2011 and 2012

Katia Regina Vieira de Rezende*, Melissa Medeiros Ferreira Hatherly, Cristiane Marques Monteiro Pimenta, Janaina Eduardo, Simone de Castro Vianna, Norberto Mangiavacchi

GESAR, Unidade de Desenvolvimento Tecnológico, Universidade do Estado do Rio de Janeiro
(Rua Fonseca Teles, 121, São Cristóvão, Rio de Janeiro, Brazil)

*Corresponding author: katiarvr@gmail.com

ABSTRACT

This study analyzed the temporal variability of phytoplankton assemblages in the surface waters of Guanabara Bay (RJ, Brazil), at six stations in front of Icaraí Inlet from April/2011 to April/2012. Our results highlight the great contribution of diatoms, dinoflagellates and cyanobacteria, represented by 111 taxa typical of estuarine and coastal areas. The coexistence of benthic and planktonic species suggests considerable hydrodinamism in these waters. All variables were homogeneous ($p > 0.05$) between the stations, but differed between sampling periods. On average, phytoplankton abundance (10^7 cells.L⁻¹) was higher than that of other estuaries and its temporal behavior was closely correlated ($p < 0.01$) with diatoms and cyanobacteria. The richness distribution pattern (7 to 27 taxa) was closely correlated ($p < 0.01$) with dinoflagellates and diatoms. Ninety per cent of all samples presented a low diversity index (< 2.0 bits.cell⁻¹), which indicated the unstable balance of the system, typical of environments subjected to eutrophication. The population structure analysis revealed that 10% of all taxa were resident, 12% visitors and 78% accidental, suggesting the influence of continental and oceanic water influxes. Between the "typical" taxa, the most common were the cyanobacteria of the order Oscillatoriales, the diatoms *Ceratoneis closterium* (= *Cylindrotheca closterium*) and *Leptocylindrus minimus* and the dinoflagellate *Prorocentrum triestinum*.

Descriptors: Tropical estuary, Temporal variation, Species Diversity, Microphytoplankton.

RESUMO

Este trabalho analisou a variabilidade sazonal da comunidade microfitoplanctônica em águas superficiais da Baía da Guanabara (RJ, Brasil) em 6 estações em frente à Enseada de Icaraí, de abril de 2011 a abril de 2012. Os resultados destacaram a alta representatividade de diatomáceas, dinoflagelados e cianobactérias, representados por 111 táxons típicos de ambientes estuarinos/costeiros. A coexistência de espécies bentônicas e planctônicas indicou o alto hidrodinamismo local. Houve homogeneidade ($p > 0,05$) entre as estações de coleta para todas as variáveis, mas diferenças entre campanhas. A densidade média (10^7 cel.L⁻¹) foi superior à de outros sistemas estuarinos e seu comportamento temporal esteve altamente correlacionado ($p < 0,01$) com diatomáceas e cianobactérias. Por sua vez o padrão de distribuição da riqueza (7 a 27 táxons) apresentou alta correlação positiva ($p < 0,01$) com dinoflagelados e diatomáceas. Baixos índices de diversidade ($< 2,0$ bits.cel⁻¹) em 90% das amostras reafirmaram o equilíbrio instável do sistema, típico de ambientes sujeitos à eutrofização. A análise da estrutura das populações estabeleceu que 10% dos táxons são residentes, 12% visitantes e 78% acidentais, reafirmando a influência do aporte continental e/ou águas oceânicas. Entre os táxons "típicos", destacaram-se cianobactérias da Ordem Oscillatoriales, diatomáceas *Ceratoneis closterium* (= *Cylindrotheca closterium*) e *Leptocylindrus minimus* e o dinoflagelado *Prorocentrum triestinum*.

Descritores: Estuário tropical, Variação temporal, Diversidade específica, Microfitoplâncton.

INTRODUCTION

The evaluation of the condition of an ecosystem requires the assessment of its primary productivity and trophic dynamics in view of the structure and ecological functions of its communities. Phytoplankton represents the basis of the main food webs in aquatic ecosystems, and its taxonomic composition and abundance respond to environmental disturbances (i.e. physical processes such as advective currents and turbulence, and chemical composition - nutrients), and to the interaction between species (i.e. competition for resources: light and nutrients) (MARGALEF, 1963; 1978). Consequently, the assessment of an aquatic ecosystem dynamics is relevant not only for the system's production, but also for the possibility of using organisms as an efficient proxy for determining natural and anthropogenic disturbances (LOBO; CALLEGARO; BENDER, 2002).

SIEBURTH, SMETACEK and LENZ (1978) proposed the classification of planktonic cells in three groups according to their size, named pico (0.2-2 μm), nano (2-20 μm) and microplankton (20-200 μm). Because of their small size, these organisms present a short generation time (hours-days), and their rapid response to environmental conditions makes them good indicators of important environmental processes such as eutrophication (HARRIS, 1986; SOMMER, 1989; REYNOLDS; PADISÁK; SOMMER, 1993).

Coastal and estuarine areas present high productivity due to nutrient rich terrestrial inputs and anthropogenic effects on distinct temporal and spatial scales. The great hydrodynamism caused by these impacts increases the ability of these regions to sustain high primary production and metabolic rates of phytoplanktonic cells (CLOERN; FOSTER; KLECKNER, 2014), due to alterations in phytoplankton community structure that are reflected in the marine food web (JI et al., 2007; CLOERN; JASSBY, 2010; LLEBOT et al., 2011).

Guanabara Bay is one of the largest embayments of the Brazilian coast. It is shallow (5-50 m depth) but presents a north-south axis of 30 km, a perimeter of 131 km, an area of 384 km², and 1.87 x 10⁹ m³ of water volume (KJERFVE et al., 1997; KJERFVE, SEELIGER; LACERDA, 2001). The climate of this region is warm and wet all year around (average humidity of 78% and temperature of 23.7° C), with a rainy season during spring-summer (September to March) and a drier period during autumn-winter (April to August), a seasonality that influences the hydrobiology of

the bay. Hydrological characteristics respond to temporal (daily and seasonal) variations of tides and cold fronts (precipitation and winds), which influence the terrestrial freshwater inflow that impacts certain areas of the bay strongly: high precipitation in summer causes an increase of the terrestrial freshwater input, the opposite occurring in winter. The freshwater input is derived from river catchment basins, which receive domestic and industrial effluents (MAYR; TENENBAUM; VILLAC, 1989), and the input of coastal seawater increases during high tide (VALENTIN et al., 1999). Tides present a semi-diurnal regime (AMADOR, 1997) with average amplitude of 0.7 m, ranging from 1.1 m during spring to 0.3 m in neap tide periods (JICA, 1994; VALENTIN et al., 1999; KJERFVE, SEELIGER; LACERDA, 2001). Guanabara Bay is surrounded by large urbanized areas such as the cities of Rio de Janeiro and Niterói (SCHWAMBORN et al., 2004), and by the second largest industrial park in the country, with around 6000 factories 1% of which account for 80% of the industrial pollution poured into the bay (CIDS, 2000). Because of these conditions, Guanabara Bay is considered a polluted eutrophic system (JICA, 1994), despite the processes of autodepuration that occur through interchanges with the ocean, which save the biota from irreversible damage (VALENTIN et al., 1999).

The sampling area is located in front of Icaraí Inlet (22°55'S-43°08'W) in Niterói city, RJ (Figure 1). The waves that enter the inlet, coming from south and southwest, lead to great hydrodynamism in this area, especially during storm surges (SILVA; RESENDE; SANTOS, 1999; SANTOS; SILVA; SALVADOR, 2004).

Phytoplankton from Guanabara Bay has been studied since the early XX century, but most of the studies are scattered in academic thesis and dissertations. The first studies that include analyses of population dynamics as a function of environmental variables were published during the 80's, and generated a broad characterization of the system's hydrobiology (MAYR; TENENBAUM; VILLAC, 1989). Recently, VILLAC and TENENBAUM (2010) have gathered information contained in 57 publications with data obtained between 1913 and 2004 in a state of art manuscript on the phytoplankton biodiversity of Guanabara Bay. The analysis of this information allowed them to conclude that, despite the fact that some of the studies published contain lists of microphytoplankton species, most of them adopt an ecological approach and highlight only the most abundant species. According to this historical study, the inventory of phytoplankton




Figure 1. Map of Guanabara Bay showing location of the sampling area (Niterói, Rio de Janeiro).

species from Guanabara Bay totaled 308 taxa, with the dominance of diatoms (62%) and dinoflagellates (32%), and other groups such as cyanobacteria, euglenophyceans, chlorophyceans, prasinophyceans, silicoflagellates and ebridae were also represented. Other studies revealed that, in Guanabara Bay, phytoplankton assemblages present spatial heterogeneity, and their abundance is comparable to that of intensely polluted estuaries.

Changes in phytoplankton communities due to the effects of natural events or pollutants can be determined by the investigation of the species' composition, cell numbers and diversity indices. Thus, the goal of this work is to describe the abundance and community structure of microphytoplankton in surface waters of one sector of Guanabara Bay, during the period between April 2011 and April 2012. These results broaden the information available on the temporal change of the phytoplankton of this system under the influence of the variation of coastal water quality and anthropogenic activities and may be valuable for future environmental monitoring and assessment programs.

MATERIAL AND METHODS

SAMPLING AND ANALYSIS

Surface water samples were taken using a Van Dorn bottle and then stored in 500 to 1000 ml glass vials and fixed with a Lugol solution (THRONDSSEN, 1978). Sampling was carried out fortnightly (36

samples) from April to June 2011, and monthly during the remaining period (60 samples) at six stations in front of Icaraí inlet (Figure 1). Microphytoplankton abundance and species composition were evaluated in accordance with HASLE (1978). Aliquots of 2 to 10 ml were analyzed by the UTERMÖHL (1958) method, using inverted microscopes (Coleman NIB-100 and Nikon TS100F) with phase contrast and 200x magnification. The phytoplankton counts were based on a minimum of 150 settled units (single cells, chain-forming and filamentous organisms). In addition, the number of cells in each settled unit (SHAW, 1964; TENENBAUM et al., 2001; GUENTHER et al., 2012) was also registered and the results were expressed as cells per liter (cells.L⁻¹).

In order to avoid missing the richness per sample, organisms that could not be identified to the species level were classified into broader taxonomic groups (class, order and family) and identified by their morphotypes according to cells shape and dimension. We would point out that the identification of most taxa requires complex training and the utilization of more advanced microscopy techniques with a higher resolution (SOURNIA, 1978; TOMAS, 1997). The classification systems used for taxonomic denomination were those of ROUND, CRAWFORD and MANN (1990) for Diatoms, FENSOME et al. (1993) for dinoflagellates, KOMÁREK and ANAGNOSTIDIS (1989; 2005) for cyanobacteria and THRONDSSEN (1997) for Chlorophytes.

DATA ANALYSIS

The Kruskal-Wallis non-parametric test was used to test the temporal and spatial variability ($p < 0.05$) in phytoplankton abundance, richness and the Shannon-Wiener Diversity index. The Mann-Whitney non-parametric test was applied to compare co-temporal independent samples during the biweekly sampling period. To establish correlation between variables the Simple Linear Correlation (Pearson's r) test, that determines the extent to which values of the two variables are "proportional" to each other, was used. The analysis of microphytoplankton community structure was performed using three ecological indexes: a) the Constancy index: the taxon was considered "resident" when it was recorded in more than 50% of the samples, "visitor" when it was recorded in 25-50% of the samples and "accidental" when recorded in less than 25% of the samples (DAJOZ, 1983); b) the Shannon-Wiener Species Diversity index and c) Pielou's Evenness index (LEGENDRE; LEGENDRE, 1998). The analysis of Similarity Percentages (SIMPER) identified the taxa responsible for the similarity of samples in each sampling period (CLARKE; WARWICK, 1994). Some analyses (Kruskal-Wallis, Mann-Whitney and Simple Linear Correlation) were carried out using STATISTICA (Version 7), other routines (Shannon-Wiener Diversity, Pielou's Evenness and Simper analysis) were performed using PRIMER (Version 5).

RESULTS

SPECIFIC COMPOSITION, RICHNESS AND ABUNDANCE

A total of 110 taxa were identified, belonging to 4 Divisions (Figure 2; Table 1): Diatoms (55 taxa; 25 species), Dinoflagellates (51 taxa; 25 species), Cyanobacteria (2 taxa) and Chlorophytes (2 taxa).

Richness, Abundance, Diversity Index and Evenness results are presented as the mean values of each sampling period, as the non-parametric tests revealed no differences ($p < 0.05$) between sampling stations. Table 2 presents information on these variables in each sampling period.

The richness distribution pattern (9 to 23 taxa) was highly correlated ($p < 0.01$) with dinoflagellates ($r = 0.77$) and diatoms ($r = 0.50$) which, together, represented 56 to 100% of the total number of taxa, while other groups' contributions were low (< 3 taxa per sample). Richness values lower than the total average (17 ± 3 taxa) were registered in May, October and December 2011, and from January to April 2012. During the biweekly




Figure 2. Percentage Contribution of Diatoms, Dinoflagellates, Cyanobacteria and Chlorophytes considering the entire study period (A) and separated by sampling cruise (B).

sampling period, only May presented a distinct ($p < 0.05$) phytoplankton assemblage, with a low richness of dinoflagellates in the second sampling of the month (Figure 3).

Variations in abundance (7.7×10^5 to 7.0×10^7 cells.L⁻¹) were correlated ($p < 0.01$) with high cell density of diatoms ($R = 0.76$) and cyanobacteria ($R = 0.67$). Values lower than the total mean ($2.0 \times 10^7 \pm 2.2 \times 10^7$ cells.L⁻¹) were registered in most of the sampling periods (Figure 4). During the biweekly sampling period, differences ($p < 0.05$) were observed due to Diatom (April), Dinoflagellate (April and June), Cyanobacteria (May and June) and Euglenophycean (June) variability. Diatom abundances varied from 1.6×10^5 to 1.0×10^8 cells.L⁻¹ ($1.3 \times 10^7 \pm 1.9 \times 10^7$ cells.L⁻¹), and this group was dominant in all sampling periods (mean contribution of 66%), except in April2 and October 2011 and February 2012. Cyanobacteria were represented by the Orders Oscillatoriales and Nostocales, showing abundances between 5.7×10^3 and 4.2×10^7 cells.L⁻¹ ($6.3 \times 10^6 \pm 1.2 \times 10^7$ cell.L⁻¹). Amongst the cyanobacteria, the Order Oscillatoriales was the most numerous throughout the sampling periods, with an average contribution of 79%. The mean abundances of Dinoflagellates and Chlorophytes were of the order of 10^5 cells.L⁻¹ and 10^4 cell.L⁻¹, respectively, with an average contribution lower than 7%.

Table 1. List of taxa found in Guanabara Bay from April 2011 to April 2012 indicating percentage of occurrence (%) in all samples (n = 96), classification by Constancy Index (R = resident; V = visitor and A = accidental) and occurrence by month (n = 13). The following classification systems were adopted: Round, Crawford and Mann (1990) for Diatoms (Bacillariophyta); Fensome et al. (1993) for dinoflagellates (Dinoflagellata); Komárek and Anagnostidis (1989; 2005) for cyanobacteria (Cyanophyta) and Throndsen (1997) for Chlorophytes (Chlorophyta). The morphotypes are not included in this table.

TAXONOMY CATEGORY	Total Occurrence (%) n = 96	Constancy Index	Occurrence by month (n = 13)												
			2011						2012						
			Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr
Division Bacillariophyta															
Class Coscinodiscophyceae															
Order Thalassiosirales															
Family Thalassiosiraceae															
<i>Thalassiosira</i> sp.	18	A	*			*	*		*	*	*			*	*
Family Skeletonemataceae															
<i>Skeletonema</i> cf. <i>costatum</i> (Greville) Cleve	25	A	*		*	*			*	*				*	*
<i>Skeletonema</i> sp.	6	A							*	*					
Order Paraliales															
Family Paraliaceae															
<i>Paralia sulcata</i> (Ehrenberg) Cleve	1	A								*					
Order Coscinodiscales															
Family Coscinodiscaceae															
<i>Coscinodiscus</i> sp.	1	A													*
Order Triceratiales															
Family Triceratiaceae															
<i>Odontella aurita</i> (Lyngbye) C. Agardh	1	A								*					
Order Hemiaulales															
Family Hemiaulaceae															
<i>Cerataulina pelagica</i> (Cleve) Hendey	3	A								*					
<i>Eucampia cornuta</i> (Cleve) Grunow	1	A								*					
<i>Eucampia</i> sp.	1	A								*					
<i>Hemiaulus membranaceus</i> Cleve	1	A								*					
Order Rhizosoleniales															
Family Rhizosoleniaceae															
<i>Dactyliosolen fragilissimus</i> (Bergon) Hasle	1	A													*
<i>Dactyliosolen phuketensis</i> (B. G. Sundström) G. R. Hasle	2	A									*				
<i>Guinardia flaccida</i> (Castracane) H. Peragallo	1	A		*											
<i>Guinardia striata</i> (Stolterfoth) Hasle	2	A				*								*	
<i>Guinardia</i> sp.	15	A			*	*		*							
<i>Proboscia alata</i> (Brightwell) Sundström	2	A			*										*
<i>Rhizosolenia setigera</i> Brightwell	42	V	*			*	*	*	*	*	*			*	*
<i>Rhizosolenia setigera</i> f. <i>pungens</i> (Cleve-Euler) Brunel	2	A									*				
Order Chaetocerotales															
Family Chaetocerotaceae															

Continued Table 1.

<i>Chaetoceros compressus</i> Lauder	3	A								*	*				
<i>Chaetoceros curvisetum</i> Cleve	2	A								*					
<i>Chaetoceros danicus</i> Cleve	6	A					*		*						*
<i>Chaetoceros radicans</i> F. Schütt	2	A					*	*							
<i>Chaetoceros socialis</i> H. S. Lauder	1	A													*
<i>Chaetoceros</i> cf. <i>socialis</i> H. S. Lauder	4	A								*					
<i>Chaetoceros</i> spp.	15	A		*	*		*		*	*	*		*		
Order Leptocylindrales															
Family Leptocylindraceae															
<i>Leptocylindrus danicus</i> Cleve	31	V	*		*	*	*		*	*	*	*	*	*	*
<i>Leptocylindrus minimus</i> Gran	76	R	*	*	*	*	*	*	*	*	*	*	*	*	*
Class Flagellariophyceae															
Order Fragilariales															
Family Fragilariaceae															
<i>Asterionellopsis glacialis</i> (Castracene) Round	5	A					*		*	*				*	
Order Licmophorales															
Familia Licmophoraceae															
<i>Licmophora</i> sp.	1	A				*									
Order Thalassionematales															
Familia Thalassionemataceae															
<i>Thalassionema nitzschioides</i> (Grunow) Mereschkowsky	5	A				*							*	*	
Class Bacillariophyceae															
Order Naviculales															
Family Phaeodactylaceae															
<i>Phaeodactylum tricornutum</i> Bohlin	11	A								*			*	*	*
Family Diploneidaceae															
<i>Diploneis</i> sp.	6	A				*	*	*	*						
Family Naviculaceae															
Complex Tropidoneis	3	A								*					
Family Pleurosigmataceae															
Complex <i>Pleurosigma</i> / <i>Gyrosigma</i>	4	A		*									*	*	
Order Thalassiosiphysales															
Family Catenulaceae															
<i>Amphora</i> sp.	1	A				*									
Order Bacillariales															
Family Bacillariaceae															
Complex <i>C. closterium</i> / <i>Nitzschia longissima</i>	7	A				*	*								
<i>Ceratoneis closterium</i> Ehrenberg (= <i>Cylindrotheca closterium</i>)	97	R	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>Pseudo-nitzschia</i> "complex <i>delicatissima</i> "	14	A	*		*		*	*				*		*	
<i>Pseudo-nitzschia</i> "complex <i>seriata</i> "	4	A								*		*		*	
Division DINOFLAGELLATA															
Class Dinophyceae															
Order Gymnodiniales															
Family Gymnodiniaceae															
<i>Akashiwo sanguinea</i> (K. Hirasaka) G. Hansen & Ø. Moestrup	6	A	*												

Continued Table 1.

<i>Amphidinium</i> spp.	27	V	*	*	*	*	*	*	*	*	*
<i>Gymnodinium</i> spp.	6	A		*							
<i>Gyrodinium</i> cf. <i>spirale</i> (Bergh) Kofoid & Swezy	2	A			*						
<i>Gyrodinium</i> spp.	2	A			*						
Order Gonyaulacales											
Family Goniomomaceae											
<i>Alexandrium</i> spp.	1	A	*								
Order Peridinales											
Family Peridiniaceae											
<i>Scrippsiella</i> cf. <i>spinifera</i> G. Honsell & M. Cabrini	2	A	*								
<i>Scrippsiella</i> cf. <i>trochoidea</i> (Stein) Balech ex Loeblich III	31	V	*	*	*	*					
<i>Scrippsiella</i> sp.	28	V	*		*	*	*	*	*	*	*
Family Congruentiaceae											
<i>Protoperidinium</i> cf. <i>bipes</i> (Paulsen) Balech	5	A			*	*	*	*			
<i>Protoperidinium</i> cf. <i>steinii</i> (Jorgensen) Balech	1	A				*					
<i>Protoperidinium</i> spp.	43	V	*	*	*	*	*	*	*		*
Order uncertain											
Family Oxytoxaceae											
<i>Oxytoxum crassum</i> Schiller	5	A	*		*				*		
<i>Oxytoxum</i> cf. <i>gladiolus</i> Stein	2	A					*		*		
<i>Oxytoxum gracile</i> Schiller	7	A		*		*	*	*		*	
<i>Oxytoxum laticeps</i> Schiller	3	A				*			*		*
<i>Oxytoxum scolopax</i> Stein	2	A	*	*							
<i>Oxytoxum</i> cf. <i>turbo</i> Kofoid	1	A					*				
<i>Oxytoxum</i> sp.	1	A		*							
Order Dinophysiales											
Family Dinophysiaceae											
<i>Dinophysis acuminata</i> Claparède & Lachmann	9	A	*	*	*	*	*	*		*	
<i>Dinophysis fortii</i> Pavillard	3	A	*								
<i>Dinophysis</i> sp.	2	A	*							*	
Family Oxyphysaceae											
<i>Oxyphysis oxytoxoides</i> Kofoid	38	V	*	*	*	*	*	*	*	*	*
Order Prorocentrales											
Family Prorocentraceae											
<i>Prorocentrum balticum</i> (Lohmann) Loeblich	4	A	*		*						*
<i>Prorocentrum compressum</i> (J. W. Bailey) Abé ex Dodge	2	A		*	*						
<i>Prorocentrum dentatum</i> Stein	1	A							*		
<i>Prorocentrum gracile</i> Schütt	5	A	*								
<i>Prorocentrum micans</i> Ehrenberg	47	V	*	*	*	*	*	*	*	*	*
<i>Prorocentrum minimum</i> (Pavillard) J. Schiller	40	V	*	*	*	*	*	*	*	*	*
<i>Prorocentrum scutellum</i> Schröder	2	A	*								
<i>Prorocentrum triestinum</i> J. Schiller	79	R	*	*	*	*	*	*	*	*	*
<i>Prorocentrum</i> sp.	7	A					*		*		*

Continued Table 1.

Class Noctiluciphyceae													
Order Noctilucales													
Family Noctilucaceae													
<i>Pronoctiluca pelagica</i> Fabre-Domérigue	3	A	*									*	
<i>Pronoctiluca spinifera</i> (Lohmann) Schiller	2	A					*		*				
Division Chlorophyta	5	A		*			*						
Class Euglenophyceae	73	R	*	*	*	*	*	*	*	*	*	*	*
<i>Eutreptiella</i> sp.	1	A	*										
Division Cyanophyta													
Class Cyanophyceae													
Order Nostocales	67	R	*	*	*	*		*	*	*	*	*	*
Order Oscillatoriales	91	R	*	*	*	*	*	*	*	*	*	*	*

Table 2. Microphytoplankton (Total and by groups) Richness, Abundance and Ecological Indexes (Species Diversity and Evenness) from April 2011 to April 2012.

Date	Richness					Abundance					Index	
	Total	Diatoms	Dinoflagellates	Cyanobacteria	Chlorophytes	Total	Diatoms	Dinoflagellates	Cyanobacteria	Chlorophytes	Species Diversity	Evenness
Apr1	18	4	12	1	1	35580718	10284643	24712437	410283	173355	0.44	0.15
Apr2	17	2	12	1	1	69754673	45297123	19789149	4639174	29226	0.44	0.15
May1	18	3	13	2	1	4107270	2244201	1392402	430008	40659	0.33	0.14
May2	12	2	6	2	2	6631668	4894936	360093	905809	470830	0.10	0.07
Jun1	19	6	10	2	1	18664938	17702565	189872	486232	286269	1.14	0.42
Jun2	20	5	13	1	1	10407854	9241535	93210	1051068	53111	0.94	0.34
Jul	23	10	11	2	1	56865758	56645156	118060	66902	35640	0.99	0.33
Aug	19	5	12	1	1	28571795	27728657	125740	439361	278036	0.11	0.04
Sep	18	5	9	2	1	13277285	12723868	286800	203216	63401	0.10	0.04
Oct	16	5	8	2	1	10212120	2486926	7372138	246037	107019	1.08	0.37
Nov	18	10	6	1	-	768861	718381	5652	46713	-	1.57	0.48
Dec	13	7	4	2	-	12459305	10876155	1569261	13889	-	0.16	0.07
Jan	9	3	4	2	1	2728105	1885932	531635	306591	3946	1.03	0.51
Feb	16	5	9	2	1	53614363	10990133	42197291	405632	21308	0.81	0.33
Mar	17	9	6	2	1	2168300	851195	1197935	109175	9996	1.07	0.40
Apr	15	7	5	2	1	1454119	345550	1008292	84125	16152	1.49	0.60

COMMUNITY STRUCTURE

According to the Constancy analysis in the study area (Table 1), 78% of the taxa were categorized as ACCIDENTAL (49 diatoms; 36 dinoflagellates; 2 chlorophytes), 10% as RESIDENT (3 diatoms; 5 dinoflagellates, 2 cyanobacteria, 1 chlorophyte), and 12% as VISITORS (3 diatoms; 10 dinoflagellates).

The Shannon-Wiener diversity index varied between 0.10 and 1.57 bits.cell⁻¹ (Table 2), with most values (56%) higher than the mean (0.74 ± 0.50 bits.cell⁻¹; Figure 5). The lowest values were registered in April, May, August, September and December 2011, related to high abundances of certain taxa of Diatoms and Dinoflagellates. Pielou's

evenness varied from 0.04 to 0.60 (0.28 ± 0.18), following the same pattern as the diversity index (Figure 5).

A total of 42 taxa (19 diatoms, 19 dinoflagellates, 2 cyanobacteria and 2 chlorophytes) was defined by SIMPER analysis as the most representative ("typical") in the study area (Table 3), for their contribution of up to 90% of the similarity between the samples of each sampling period. The taxonomic classes with the greatest contributions to the similarity between samples were the dinoflagellates (13-60%) and diatoms (13-55%) (Figure 6). Individual contribution to sample similarity by a singular taxon varied between 2 and 19% in each sampling period. We highlight unidentified cyanobacteria




Figure 3. Total Microphytoplankton, Diatoms and Dinoflagellates Richness from April 2011 to April 2012. Dotted line indicates general mean of Total Richness.




Figure 4. Total Microphytoplankton, Diatoms and Dinoflagellates Abundance (cell.L⁻¹) from April 2011 to April 2012. Dotted line indicates general mean of Total Density.

of the Order Oscillatoriales and Order Nostocales, diatoms *Ceratoneis closterium* (= *Cylindrotheca closterium*) and *Leptocylindrus minimum* and dinoflagellate *Prorocentrum triestinum* for their broad occurrence (> 80%) with high mean abundance (10^4 - 10^7 cells.L⁻¹) in many periods. Table 3 shows the high abundance of certain morphotypes with several shapes and dimensions between 20 and 150 μ m (1 centric diatom, 4 pennate diatoms and 8 dinoflagellates) and brings out the need for taxonomic studies so that species that make a major contribution to community structure may be correctly identified.

When compared with the Constancy index, SIMPER analysis was more effective in determining which taxa made a greater contribution to sample similarity in each




Figure 5. Shannon-Wiener Species Diversity index (bits.cell⁻¹) and Pielou's Evenness index from April 2011 to April 2012. Dotted line indicates general mean of Diversity index.

period. The Constancy index just takes into account the occurrence of each taxon, while SIMPER considers abundance as well as occurrence. This could be observed for the diatom *Skeletonema costatum* classified as ACCIDENTAL by the Constancy index with occurrence in only 25% of the samples. However, this taxon was highlighted by SIMPER for its high mean densities (2.8×10^6 - 2.2×10^7 cells.L⁻¹) that favored a contribution of up to 12% for the similarity of the samples for the periods of June1, June2 and July.

DISCUSSION

The microphytoplankton community of Guanabara Bay was characterized by typical estuarine and coastal species, influenced by a large number of multifactorial abiotic and biotic processes (LLEBOT et al., 2011; CLOERN; FOSTER; KLECKNER, 2014). Among all the factors that may affect the taxonomic composition and temporal variation of phytoplankton in Guanabara Bay are found: tidal cycle, seasonality of water masses of adjacent continental shelf, cold fronts and continental drainage (rainfall, inflow of domestic sewage, etc.)

The elevated number of taxa classified as ACCIDENTAL confirmed the high local hydrodynamism as being a consequence of the environmental factors mentioned above. The processes which caused such turbulence were probably responsible for the presence of benthic species (diatoms *Paralia sulcata*, *Licmophora* sp., *Diploneis* sp., *Amphora* sp.) along with planktonic ones (diatoms *Leptocylindrus danicus*, *L. minimus*,

Table 3. Simper analysis results showing average abundance (cell.L⁻¹) of each taxon from April 2011 to April 2012. Values underlined represent taxa with contribution $\geq 10\%$. Taxa were ordered by the sum (Σ) of average abundance considering the entire sampling period. The Morphotypes are included.

	Σ	2011	2012															
		Apr1	Apr2	May1	May2	Jun1	Jun2	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	
DIATOMS																		
<i>Ceratoneis closterium</i>	1.5 x 10 ⁸	268795	45288034	2204603	4875816	4900030	6361924	34217077	27582734	12664516	375628		10722583	1450707	107327	15027	34250	
<i>Skeletonema cf. costatum</i>	3.7 x 10 ⁷					12054996	2771530	22341109										
<i>Leptocylinthus minutus</i>	2.5 x 10 ⁷	9827038		24869		488356	13543		97592		1992860	53319	124846	423125	10596338	690848	238276	
<i>Rhizosolenia setigera</i>	4.2 x 10 ⁵										10755	377664	8050			12286	9575	
<i>Phaeodactylum tricornutum</i>	2.7 x 10 ⁵														269895			
<i>Gomardtia</i> sp.	1.7 x 10 ⁵					135964	31867											
<i>Chaetoceros cf. socialis</i>	8.2 x 10 ⁴											82115						
<i>Leptocylinthus denticus</i>	4.3 x 10 ⁴							13255				14908				14872		
<i>Chaetoceros</i> spp.	3.0 x 10 ⁴											29791						
Complex <i>C. closterium</i> / <i>Nitzschia longissima</i>	2.2 x 10 ⁴							22222										
<i>Pseudo-nitzschia</i> "delicatissima complex"	2.2 x 10 ⁴									22107								
<i>Thalassiosira</i> sp.	1.7 x 10 ⁴									11087		13311	3630					
<i>Skeletonema</i> sp.	1.1 x 10 ⁴																	
Complex <i>Tropidoneis</i>	1.2 x 10 ³											1222						
MORPHOTYPES																		
Centric diatom (20-50 μm)	1.1 x 10 ⁵			7062				20207	24962			61032	947				13048	
Pennate diatom (linear; 50-100 μm)	5.9 x 10 ⁴															58953		
Pennate diatom (lanceolate; 20-50 μm)	2.3 x 10 ⁴									6174						16701		
Pennate diatom (linear; 100-150 μm)	1.6 x 10 ⁴															16464		
Pennate diatom (retangular; 20-50 μm)	4.7 x 10 ⁴							4658										
DINOFLLAGELLATES																		
<i>Akashiwo sanguinea</i>	4.2 x 10 ⁶		4200474															
<i>Prorocentrum triestinum</i>	8.2 x 10 ⁵	51624	36760		30273	116844	254136	7780	119102	45344	93210	23541		23938	8681		10838	
<i>Prorocentrum micans</i>	1.8 x 10 ⁵	126192	6904				27087				18854							

[illegible]




Figure 6. Percentual Contribution of phytoplankton groups determined by similarity percentage analysis (SIMPER).

Thalassionema nitzschioides, *Coscinodiscus* spp., *Pseudo-nitzschia* spp., *Skeletonema* cf. *costatum*; Dinoflagellates *Alexandrium* sp.; *Protoperidinium* spp.) (RICARD, 1987; HASLE; SYVERTSEN, 1997). It is important to state that benthic microalgae are important primary producers in shallow aquatic ecosystems and that their production exceeds that of planktonic microalgae in certain periods (BERGESCH; ODEBRECHT; ABREU, 1995; BRANDINI; FERNANDES, 1996). The relevance of a taxon in the study area was determined by its high contribution to sample similarity in each season. Thus SIMPER analysis confirmed the representativity of “RESIDENT” taxa as well as the occasional contribution of others during the study period.

The organisms identified in the present study had already been observed in Guanabara Bay due to their wide distribution in the system (VILLAC; TENENBAUM, 2010). The low frequency and abundance of *Skeletonema costatum* called the authors' attention since this diatom has been consistently reported as opportunist in eutrophic environments. This pattern change may be associated with the location of the sampling stations close to the bay's entrance, a less polluted site due to the contribution of more saline, cleaner and clearer coastal water (SANTOS et al., 2007). Similar results were found by GUENTHER et al. (2012) in a short temporal scale investigation also in the entrance of Guanabara Bay during summer 2004, in which *S. costatum* was not considered an abundant taxon. Likewise, SANTOS et al. (2007) registered higher densities of this species in the inner portion when compared with the entrance of the bay. These outcomes suggest that the hydrodynamic conditions found in different areas of Guanabara Bay may exercise great influence on the representation of *S.*

costatum. Another important aspect to be considered is that this diatom was the subject of a taxonomic review revealing that the genus biodiversity at any given place is most likely underestimated and may include more than one species (SARNO et al., 2007; KOISTRA et al., 2008). For instance, the morphology of *Skeletonema* species was examined in coastal waters of southern Brazil and the authors came to the conclusion that this genus is highly diverse in that geographical area, with the occurrence of four confirmed species: *S. costatum*, *S. pseudocostatum*, *S. potamos* and *S. tropicum* (BERGESCH; GARCIA; ODEBRECHT, 2009).

The recurrent observation of potentially harmful species in the bay requires some attention, specially concerning diatoms of the genus *Pseudo-nitzschia* due to its ability to produce a powerful neurotoxin known as domoic acid (VILLAC; TENENBAUM, 2001; VILLAC; DOUCETTE; KACZMARSKA, 2010). Some studies have shown that *Pseudo-nitzschia* abundance is related to a major entry of nutrients into the water column (PARSON; DORTCH, 2002). Some other potentially harmful species were also found in Guanabara Bay, such as the diatoms *Cerataulina pelagica*, *Leptocylindrus danicus*, *L. minimus* and Dinoflagellates *Akashiwo sanguinea*, *Oxyphysis oxytoxoides*, *Prorocentrum micans*, *P. balticum*, *Dinophysis acuminata*, *Scrippsiella* cf. *trochoidea* (FRYXELL; VILLAC, 1999; HALLEGRAEFF; ANDERSON; CEMBELLA, 2003).

The average cell density (10^7 cel.L⁻¹), described in this study, were higher than those found in other Brazilian coastal estuaries: Sepetiba Bay-RJ (10^4 - 10^6 cel.L⁻¹; TENENBAUM et al. 2004), Estuary of Paraíba do Norte River-PB (10^3 - 10^6 cel.L⁻¹; SASSI, 1991), Paranaguá Bay-PR (10^4 - 10^6 cel.L⁻¹; BRANDINI, 1985; BRANDINI; THAMM, 1994), São Sebastião Channel-SP (10^5 - 10^6 cel.L⁻¹; GIANESSELLA et al., 1999). However, studies in Guanabara Bay carried out between 1913 and 2004 showed similar values (10^5 - 10^9 cel.L⁻¹) (VILLAC; TENENBAUM, 2010). These differences may be related to different environmental conditions related to changes in time and space as well as to human intervention (artificial eutrophication, dredging, navigation, etc.).

Diatoms and dinoflagellates were the predominant groups and together were responsible for more than 96% of the specific composition, in accordance with the literature that indicates their predominance in Brazilian coastal regions (BRANDINI et al., 1997). The dominance of diatoms in shallow coastal regions reflects the instability

of this environment, where turbulence homogenizes the water column, increases nutrient concentration in the euphotic zone and reduces cell sinking (MARGALEF, 1978; SMETACEK, 1988; LLEBOT et al., 2011). The high representativeness (40%) of chain-forming diatoms (species of *Chaetoceros*, *Leptocylindrus*, *Hemiaulus*, *Skeletonema*, *Pseudo-nitzschia*, *Eucampia*) has been related to estuarine and coastal environments (FERNANDES; BRANDINI, 2004; BÖTTJER; MORALES, 2005). According to REYNOLDS; PADISÁK and SOMMER (1993) this morphology provides a larger surface for light capture and represents an advantage over environments with high suspended solid concentrations. Laboratory assays also suggest that these organisms thrive in the environment as a result of a lower grazing pressure by microzooplankton (BODE et al., 2005).

The combination of different nutrition strategies (autotrophic and mixotrophic) may give dinoflagellates a competitive advantage over other organisms, even in the conditions of limited light that are very common in estuaries and river deltas (LALLI; PARSONS, 1993). Some taxa such as species of the genus *Protoperidium* and of the Order Gymnodiniales, considered heterotrophs (STEIDINGER; TANGEN, 1997), have also been frequently observed in samples. The high contribution of organisms of the Order Gymnodiniales to richness and abundance confirms VILLAC and TENENBAUM (2010) statement that Guanabara Bay dinoflagellates, although frequent and highly representative, are underestimated by virtue of the sampling and analysis procedures used in most of the studies. These dinoflagellates have extremely fragile cells that are deformed or destroyed by the commonly used fixative substances. So that a more precise identification of most taxa demands a different methodology from fixation to sample handling combined with the utilization of more advanced microscopy techniques with higher resolution.

The cyanobacteria, of the Orders Oscillatoriales and Nostocales, were also very important in phytoplankton's attaining up to 99% of abundance in some samples. These results were expected and suggest that the increase in filamentous cyanobacteria density is a response to rainy periods and high levels of eutrophication (SANTOS et al., 2007; VILLAC; TENENBAUM, 2010).

The high concentration of euglenophyceans in the study area can be associated with inland polluted waters (LIMA; TENENBAUM; VALENTIN, 2010; GUENTHER et al. 2012), whereas these organisms require organically enriched water for growth (LEE, 2008).

The presence of the diatoms *Proboscia alata*, *Rhizosolenia setigera*, *Hemiaulus membranaceus*, *Guinardia striata*, *Eucampia cornuta*, *Dactyliosolen phuketensis* and the dinoflagellates *Prorocentrum balticum*, *Pronoctiluca pelagica*, *Oxytoxum gracile* indicates the influence of the Tropical Water that flows along the Brazilian continental shelf mixed with Coastal Water (HASLE; SYVERSTSEN, 1997; STEIDINGER; TANGEN, 1997).

The low diversity index ($< 2.0 \text{ bits.cell}^{-1}$), in 90% of samples, is a typical characteristic of systems in unstable equilibrium such as estuaries or polluted environments subject to local eutrophication (LLEBOT et al., 2011). Water mass enrichment processes first induce the proliferation of a reduced number of species and consequently a profound reduction in the species diversity index (MARGALEF, 1958; MARGALEF, 1980).

CONCLUSION

The composition and abundance of the phytoplankton of a certain region is certainly associated with the local hydrography. The interaction between water movements and phytoplankton organisms results from a combination of environmental factors. Thus the hydrodynamic properties of each region play an important role in the temporal variability and structure of phytoplankton populations (MARGALEF, 1978; ESTRADA; BERDALET, 1997). The information produced by this study will, therefore, add to knowledge of the Guanabara Bay system and will be useful for management purposes and also for the regulation of land use in the area surrounding the bay.

ACKNOWLEDGEMENTS

The authors would like to thank the Centro de Biologia Experimental Oceanus for field support and the provision of data. The authors are also grateful to their colleagues Priscila Kienteca Lange and Marcio Lincoln da Silva for their attention and kindness in translating this manuscript.

REFERENCES

- AMADOR, E. S. Baía de Guanabara e ecossistemas periféricos: homem e natureza. Rio de Janeiro: Reproarte. 1997. 539 p.
- BERGESCH, M.; ODEBRECHT, C.; ABREU, P. C. O. Microalgas do estuário da Lagoa dos Patos: interação entre o sedimento e a coluna de água. Oecol. Bras., v.1, p. 273-289, 1995.

- BERGESCH, M.; GARCIA, M.; ODEBRECHT, C. Diversity and morphology of *Skeletonema* species in Southern Brazil, Southwestern Atlantic Ocean. J. Phycol., v. 45, n. 6, p. 1348-1352, 2009.
- BODE, A.; ÁLVAREZ-OSSORIO, T. GONZÁLEZ, N. LORENZO, J. RODRÍGUEZ, C. VARELA, M. VARELA, M.; M. Seasonal variability of plankton blooms in the Ria de Ferrol (NW Spain): II. Plankton abundance, composition and biomass. Estuarine, Coastal Shelf Sci., v. 63, n. 1/2, p. 285-300, 2005.
- BÖTTJER, D.; MORALES, C. E. Microzooplankton grazing in a coastal embayment off Concepcion, Chile, (~36° S) during non-upwelling conditions. J. Plankton. Res., v. 27, n. 4, p. 383-391, 2005.
- BRANDINI, F. P. Seasonal succession of the phytoplankton in the Bay of Paranaguá (Paraná State - Brazil). Rev. Bras. Biol., n. 45, n. 4, p. 687-694, 1985.
- BRANDINI, F. P.; FERNANDES, L. F. Microalgae of the continental shelf off Paraná State, Southeastern Brazil: a review of studies. Rev. Bras. Oceanogr., v. 44, n. 1, p. 69-80, 1996.
- BRANDINI, F. P.; THAMM, C. A. C. Variações diárias e sazonais do fitoplâncton e parâmetros ambientais na Baía de Paranaguá. Nerítica, v. 8, n.1/2, p. 55-72, 1994.
- BRANDINI, F. P.; LOPES, R. M.; GUTSEIT, K. S.; SPACH, H. L.; SASSI, R. Planctonologia na Plataforma Continental do Brasil: diagnose e revisão bibliográfica. Brasília: Publicação do Ministério do Meio Ambiente, CIRM e Fundação de Estudos do Mar, 1997. 196 p.
- CIDS - Centro Internacional de Desenvolvimento Sustentável; Baía de Guanabara - Dossiê Sócio-Ambiental, Rio de Janeiro, 2000.
- CLARKE, K. R.; WARWICK, R. M. Change in marine communities: an approach to statistical analysis and interpretation. Plymouth: Plymouth Marine Laboratory, 1994. 144 p.
- CLOERN, J. E., FOSTER, S. Q., KLECKNER, A. E. Phytoplankton primary production in the world's estuarine-coastal ecosystems. Biogeosciences, v. 11, p. 2477-2501, 2014.
- CLOERN, J. E., JASSBY, A. D. Patterns and scales of phytoplankton variability in estuarine-coastal ecosystems. Estuar. Coasts., v. 33, n. 2, p. 230-241, 2010.
- DAJOZ, R. Ecologia geral. Petrópolis: Vozes, 1983. 472 p.
- ESTRADA, M.; BERDALET, E. Phytoplankton in a turbulent world. Sci. Mar., v. 61, supl. 1, p. 125-140, 1997.
- FENSOME, R. A.; TAYLOR, F. J. R.; NORRIS, G.; SARJEANT, W. A. S.; WHARTON, I. D.; WILLIAMS, G. L. A classification of living and fossil dinoflagellates. Hanover: Sheridan Press, 1993. 350 p.
- FERNANDES, L. F.; BRANDINI, F. P. Diatom associations in the shelf waters off Paraná State, Southern Brazil: annual variations in relation to environmental factors. Braz. J. Oceanogr., v. 52, n. 1, p. 19-34, 2004.
- FRYXELL, G. A.; VILLAC, M. C. Toxic diatoms. In: STORMER, E. F.; SMOLL, J. P. (Eds.). The Diatoms, Applications for the Environmental and Earth Sciences. 2.ed. Cambridge: Cambridge University Press, 1999. p. 419-428.
- GIANESELLA, S. M. F.; KUTNER, M. B. B.; SALDANHA-CORREIA, F. M. P.; POMPEU, M. Assessment of plankton community and environmental conditions in São Sebastião Channel prior to the construction of a produced water outfall. Rev. Bras. Oceanogr., v. 47, n. 1, p. 29-46, 1999.
- GUENTHER, M.; LIMA, I.; MUGRABE, G.; TENENBAUM, D. R.; GONZALEZ-RODRIGUEZ, E.; VALENTIN, J. L. Small time scale plankton structure variations at the entrance of a tropical eutrophic bay (Guanabara Bay, Brazil) Braz. J. Oceanogr., v. 60, n. 4, p. 405-414, 2012.
- HALLEGRAEFF, G. M.; ANDERSON, D. M.; CEMBELLA, A. D. Manual on harmful marine microalgae. Monographs on Oceanographic Methodology, 11. Paris: United Nations Educational, Scientific and Cultural Organization (UNESCO) Publishing, 2003. 793 p.
- HARRIS, G. P. Phytoplankton ecology: structure, function and fluctuation. London: Chapman & Hall, 1986. 384 p.
- HASLE, G. R. Using the inverted microscope. In: SOURNIA, A. (Ed.). Phytoplankton Manual. Paris: UNESCO, 1978. p. 191-196.
- HASLE G. A.; SYVERTSEN E. E. Marine diatoms. In: TOMAS, C. R. (Ed.). Identifying Marine Phytoplankton. New York: Academic Press, 1997. p. 5-386.
- JI, R.; DAVIS, C.; CHEN, C.; TOWNSEND, D.; MOUNTAIN, D.; BEARDSLEY, R. Influence of ocean freshening on shelf phytoplankton dynamics. Geophys. Res. Lett., 2007.
- JICA. The study of the Guanabara Bay Ecosystem. Japan International Cooperation Agency. Tokyo: Kokusai Kogyo Co., Ltd, 1994.
- KJERFVE, B.; RIBEIRO, C. H. A.; DIAS, G. T. M.; FILIPPO, A. M.; QUARESMA, V. S. Oceanographic characteristics of an impacted coastal bay: Baía de Guanabara, Rio de Janeiro, Brazil. Cont. Shelf Res., v. 17, n. 3, p. 1609-1643, 1997.
- KJERFVE, B.; SEELIGER, U.; LACERDA, L. D. A. Summary of natural and human-induced variables on coastal marine ecosystems of Latin America. In: SEELIGER, U.; KJERFVE, B. (Org.). Coastal Marine Ecosystems of Latin America. Heidelberg, Berlin: Springer Verlag, 2001. p.341-353.
- KOOISTRA, W. H. C. F., SARNO, D., BALZANO, S., GU, H., ANDERSEN, R. A.; ZINGONE, A. Global diversity and biogeography of *Skeletonema* species (Bacillariophyta). Protist, v. 159, n. 2, p.177-193, 2008.
- KOMÁREK, J.; ANAGNOSTIDIS, K. Modern approach to the classification system of cyanophytes (Nostocales). Arch. Hydrobiol. Suppl., v. 82, n. 3, p. 247-345, 1989.
- KOMÁREK, J. ANAGNOSTIDIS, K. Süßwasserflora von Mitteleuropa. Cyanoprokaryota 2. Teil: Oscillatoriales. In: BÜDEL, B.; KIENITZ, L.; GÄRTNER, G.; SCHAGERL, M. (Eds.). Süßwasserflora von Mitteleuropa 19.: München: Elsevier Spektrum Akademischer Verlag, 2005. 759 p.
- LALLI, C. M.; PARSONS, T. R. Biological Oceanography: an introduction. New York: Elsevier Butterworth-Heinemann, 1993. 301 p.
- LEE, R. E. Phycology. 4.ed. Cambridge: Cambridge University Press, 2008. p. 245.
- LEGENDRE, L.; LEGENDRE, P. Numerical Ecology. Developments in environmental Modelling, 20. New York: Elsevier, 1998. 853 p.

- LIMA, I. V.; TENENBAUM, D. R.; VALENTIN, J. L. Variação em Micro-escala temporal do Nano e Microplâncton da baía de Guanabara (RJ). In: XIII Congresso Brasileiro de Ficologia, 2010, Paraty, RJ. Anais do XIII Congresso Brasileiro de Ficologia, 2010.
- LLEBOT, C.; SOLÉ, J.; DELGADO, M.; FERNÁNDEZ-TEJEDOR, M.; JORDI CAMP, J.; ESTRADA, M. Hydrographical forcing and phytoplankton variability in two semi-enclosed estuarine bays. *J. Mar. Syst.*, v. 86, n. 3/4, p. 69-86, 2011.
- LOBO, E. A.; CALLEGARO, V. L. M.; BENDER, E. P. Utilização de Algas Diatomáceas Epilíticas como Indicadoras da Qualidade da Água em Rios e Arroios da Região Hidrográfica do Guaíba, RS, Brasil. Santa Cruz do Sul: EDUNISC, 2002. 127 p.
- MARGALEF, R. La diversidad. In: MARGALEF, R. (Org.). Ecología. Barcelona: Omega, 1980. p. 359-382.
- MARGALEF, R. Life-forms of phytoplankton as survival alternatives in an unstable environment. *Oceanol. Acta*, v.1, n. 4, p. 493-509. 1978.
- MARGALEF, R. Succession in marine populations. In: VIRA, R. (Ed.) *Advancing Frontiers of plant Sciences 2*. New Delhi: Institute for Advanced Science Culture, 1963. p. 137-88.
- MARGALEF, R. Temporal succession and spatial heterogeneity in phytoplankton. In: BUZZATI-TRAVERSO, A. (Ed.) *Perspectives in Marine Biology*. Berkeley: University of California Press, 1958. p. 323-349.
- MAYR, L. M.; TENENBAUM, D. R.; VILLAC, M. C. Hydrobiological characterization of Guanabara bay. In: MAGOON, O.; NEVES, C. (Eds.). *Coastlines of Brazil*. New York: American Society of Civil Engineers, 1989. p. 124-138.
- PARSON, M. L.; DORTCH, Q. Sedimentological evidence of an increase in *Pseudo-nitzschia* (Bacillariophyceae) abundance in response to coastal eutrophication. *Limnol. Oceanogr.*, v. 47, n. 2, p. 551-558, 2002.
- REYNOLDS, C.S.; PADISÁK, J.; SOMMER, U. Intermediate disturbance in the ecology of phytoplankton and the maintenance of species diversity: a synthesis. *Hydrobiologia*, v. 249, p. 183-188, 1993.
- RICARD, M. Atlas du Phytoplancton Marin. Diatomophycées. Paris: Editions Du Centre National de la Recherche Scientifique, 1987. 297 p.
- ROUND, F. E.; CRAWFORD, R. M.; MANN, D. G. The diatoms: biology and morphology of the genera. Cambridge: Cambridge University Press, 1990. 747 p.
- SANTOS, C. L.; SILVA, M. A. M.; SALVADOR, M. V. S. Dinâmica sazonal e os efeitos das ressacas nas praias de Niterói/RJ. *Rev. Bras. Geoci.*, v. 34, n. 3, p. 355-360, 2004.
- SANTOS, V. S.; VILLAC, M. C.; TENENBAUM, D. R.; PARANHOS, R. Auto-and heterotrophic nanoplankton and filamentous bacteria of Guanabara bay (RJ, Brazil): estimates of cell/filament numbers versus carbon content. *Braz. J. Oceanogr.*, v. 55, p. 133-143, 2007.
- SARNO, D.; KOOISTRA, W. C. H. F.; BALZANO, S.; HARGRAVES, P. E.; ZINGONE, A. Diversity in the genus *Skeletonema* (Bacillariophyceae): III. Phylogenetic position and morphological variability of *Skeletonema costatum* and *Skeletonema grevillei*, with the description of *Skeletonema ardens* sp. nov. *J. Phycol.*, v. 43, p. 156-170, 2007.
- SASSI, R. Phytoplankton and environmental factors in the Paraíba do Norte river estuary, northeastern Brazil: composition, distribution and quantitative remarks. *Bol. Inst. Oceanogr.*, v. 39, n. 2, p. 93-115, 1991.
- SCHWAMBORN, R.; BONECKER, S. L. C.; GALVÃO, I. B.; SILVA, T. A.; NEUMANN-LEITÃO, S. Mesozooplankton grazing under conditions of extreme eutrophication in Guanabara Bay, Brazil. *J. Plankton Res.*, v. 26, n. 9, p. 983-992, 2004.
- SHAW, A. B. Time in stratigraphy. New York: McGraw-Hill Book Company, 1964.
- SIEBURTH, J. M.; SMETACEK, V.; LENZ, J. Pelagic ecosystem structure: heterotrophic compartments of plankton and their relationship to plankton size fractions. *Limnol. Oceanogr.*, v. 23, n. 6, p. 1256-1263, 1978.
- SILVA, M. A. M.; RESENDE, M. C. C. M.; SANTOS, C. L. Um estudo sobre a dinâmica das praias de Niterói (Baía de Guanabara, RJ). *Anais da Academia Brasileira de Ciências*, v. 71, n. 4, p. 962-967, 1999.
- SMETACEK, V. Plankton characteristics. In: *Ecosystems of the world: continental shelves*. v. 2. Amsterdam: Elsevier, 1988. p. 93-130.
- SOMMER, U. Plankton Ecology: succession in plankton communities. Berlin: Springer Verlag, 1989. 369 p.
- SOURNIA, A. Phytoplankton Manual. Monogr. oceanogr. Methodol., U.N., v. 6, 1978.
- STEIDINGER, K. A.; TANGEN, K. Dinoflagellates. In: TOMAS, C. R. (Ed.). *Identifying Marine Phytoplankton*. New York: Academic Press, 1997. p. 387-584.
- TENENBAUM, D. R.; VILLAC, M. C.; GOMES, E. A. T.; CUPELO, A. C.; SANTOS, V. S. A new "sight" on microbial plankton ecology: coastal x oceanic system in Brazil. In: FARIA, B. M.; FARJALLA, V. F.; ESTEVES, F. A. (Eds.). *Aquatic microbial ecology in Brazil. Series Oecologia Brasiliensis*, v. IX. Rio de Janeiro: PPGE-UFRJ, 2001. p. 133-152.
- TENENBAUM, D. R.; VILLAC, M. C.; VIANA, S. C.; MATOS, M.; HATHERLY, M.; LIMA, I. V.; MENEZES, M. Phytoplankton identification atlas Sepetiba Bay, Brazil. Rio de Janeiro/London: IOC, 2004b. 100 p.
- THRONDSSEN, J., 1997. The Planktonic Marine Flagellates. In: TOMAS, C.R. (ed.) *Identifying Marine Phytoplankton*. New York: Academic Press, 1997. p. 591-729.
- THRONDSSEN, J. Preservation and storage. In: SOURNIA, A., (Ed.). *Phytoplankton manual*. Monogr oceanogr. Methodol., 1978. p. 69-74.
- TOMAS, C. R. *Identifying Marine Phytoplankton*. New York: Academic Press, 1997. 583 p.
- UTERMÖHL, H. Perfeccionamiento del metodo cuantitativo de fitoplancton. *Comun. Assoc. Int. Limnol. Teor. Apl.*, v. 9, p. 1-89, 1958.
- VALENTIN, J. L.; TENENBAUM, D. R.; BONECKER, A. C. T.; BONECKER, S. L. C.; NOGUEIRA, C. R.; VILLAC, M. C. O sistema planctônico da Baía de Guanabara: síntese do conhecimento. In: SILVA, S. H. G.; LAVRADO, H. P. (Eds.). *Ecologia dos ambientes costeiros do estado do Rio de Janeiro. Série Oecologia Brasiliensis*. Rio de Janeiro: PPGE-UFRJ, 1999. p. 35-59.
- VILLAC, M. C.; TENENBAUM, D. R. The phytoplankton of Guanabara Bay, Brazil. I. Historical account of its biodiversity. *Biota Neotrop.*, v. 10, p. 271-293, 2010.

VILLAC, M. C.; TENENBAUM, D. R. The coastal *Pseudo-nitzschia* from the state of Rio de Janeiro, Brazil. In International Conference on Harmful Algal Blooms. Hobart, v. 1, p. 34-37, 2001.

VILLAC, M. C.; DOUCETTE, G. J.; KACZMARSKA, I. Toxic marine diatoms. In: STOERMER, E. F.; SMOLL, J. P. (Eds.). The Diatoms, Applications for the Environmental and Earth Sciences. 2. ed. Cambridge: Cambridge University Press, 2010. p. 536-547.