

180 x 265 mm

5 mm

53 mm

18,5 mm

18 mm

2,5 mm

4 mm

1 mm

2,5 mm

6,5 mm

4 mm

5 mm

4 mm

3,5 mm

3,5 mm

7,5 mm

original article

Cambria Bold

9 pt/ 10,8 pt - CO M0 Y0 K100

BJPT

Brazilian Journal of  
Physical Therapy

Prospective monitoring and self-report of previous falls  
among older women at high risk of falls and fractures:  
a study of comparison and agreement

Cambria Bold

14 pt/ 16,8 pt - CO M0 Y0 K100

Patrícia A. Garcia<sup>1</sup>, João M. D. Dias<sup>2</sup>, Sílvia L. A. Silva<sup>3</sup>,  
Rosângela C. Dias<sup>2</sup>

Cambria Bold

8 pt/ 9,6 pt - CO M0 Y0 K100

ABSTRACT | **Background:** The identification of the occurrence of falls is an important step for screening and for  
assessing the risk of falls for the elderly. The methods of monitoring these events are susceptible to recording biases,  
and the choice of the most accurate method remains challenging. **Objectives:** (i) To investigate the agreement between  
retrospective self-reporting and prospective monitoring of methods of recording falls, and (ii) to compare the retrospective  
self-reporting of falls and the prospective monitoring of falls and recurrent falls over a 12-month period among older  
women at high risk of falls and fractures. **Method:** A total of 118 community-dwelling older women with low bone  
density were recruited. The incidence of falls was monitored prospectively in 116 older women (2 losses) via monthly  
phone calls over the course of a year. At the end of this monitoring period, the older women were asked about their recall  
of falls in the same 12-month period. **Results:** The agreement between the two methods was analyzed, and the sensitivity and  
specificity of self-reported previous falls in relation to the prospective monitoring were calculated. **Results:** There was  
moderate agreement between the prospective monitoring and the retrospective self-reporting of falls in classifying fallers  
(Kappa=0.595) and recurrent fallers (Kappa=0.589). The limits of agreement were 0.35±1.66 falls. The self-reporting of  
prior falls had a 67.2% sensitivity and a 94.2% specificity in classifying fallers among older women and a 50% sensitivity  
and a 98.9% specificity in classifying recurrent fallers. **Conclusion:** Self-reporting of falls over a 12-month period  
underestimated 32.8% of falls and 50% of recurrent falls. The findings recommend caution if one is considering replacing  
monthly monitoring with annual retrospective questioning.

Cambria Bold

8 pt/ 9,6 pt - CO M0 Y0 K80

**Keywords:** aged; bone density; accidental falls; rehabilitation; mental recall.

Times New Roman Regular

9 pt/ 10,8 pt - CO M0 Y0 K100

HOW TO CITE THIS ARTICLE

Cambria Bold

8 pt/ 9,6 pt - CO M0 Y0 K80

Garcia PA, Dias JMD, Silva SLA, Dias RC. Prospective monitoring and self-report of previous falls among older women  
at high risk of falls and fractures: a study of comparison and agreement. Braz J Phys Ther. 2015 May-June; 19(3):217-225.  
<http://dx.doi.org/10.1590/bjpt-rbf.2014.0095>

Times New Roman Regular

8,5 pt/ 11 pt - CO M0 Y0 K100

• Introduction

Cambria Bold

12 pt/ 14,4 pt - CO M0 Y0 K100

Falls are a common problem with high prevalence among the  
elderly population, even among those who are active  
and healthy, and constitute one of the major preventable  
geriatric syndromes<sup>1</sup>. Among the community-dwelling  
elderly, approximately 30% suffer a fall each year,  
and half experience recurrent falls<sup>2</sup>. Elderly women  
with osteoporosis and having a high risk of fractures  
exhibit an even higher frequency of falls (51.1%)<sup>3</sup>.  
A significant portion of these falls results in injuries  
(36%)<sup>4</sup>, fractures (3.4% to 19%)<sup>2,4,5</sup>, and the need for  
medical assistance (8 to 19%)<sup>4,5</sup> and affects lifestyle  
choices, creating a high socio-economic burden<sup>6</sup>.  
Additionally, experiencing one or more falls in the  
course of one year significantly increases the chances

Cambria Bold

12 pt/ 14,4 pt - CO M0 Y0 K100

of the occurrence of new episodes in the following  
year among the community-dwelling elderly<sup>4,5</sup> and  
postmenopausal women<sup>1,7</sup>.  
Thus, the surveillance of falls among the elderly  
represents a priority health issue<sup>6</sup>, which is why  
questioning the occurrence of falls has been  
used in clinical/scientific decision making<sup>8-10</sup>. Several  
methods have been suggested for monitoring the  
occurrence of falls among the community-dwelling  
elderly, including questions asking individuals to recall  
these events at several intervals by means of telephone,  
face-to-face or mail interviews, information obtained  
from medical records, and/or prospective records using  
falls calendars or diaries<sup>8,9,11-13</sup>. However, the elderly

Times New Roman Regular

10 pt/ 12,5 pt - CO M0 Y0 K100

<sup>1</sup>Colegiado de Fisioterapia, Universidade de Brasília (UnB), Brasília, DF, Brasil  
<sup>2</sup>Programa de Pós-graduação em Ciências da Reabilitação, Escola de Educação Física, Fisioterapia e Terapia Ocupacional, Universidade Federal de  
Minas Gerais (UFMG), Belo Horizonte, Minas Gerais, Brasil  
<sup>3</sup>Universidade Federal de Alfenas (Unifal), Alfenas, Minas Gerais, Brasil  
Received: June 20, 2014 Revised: Sept. 09, 2014 Accepted: Feb. 04, 2015

Cambria Regular

7 pt/ 8,4 pt - CO M0 Y0 K100

Braz J Phys Ther. 2015 May-June; 19(3):217-225

Cambria Regular

7 pt/ 8,4 pt - CO M0 Y0 K100

217

Times New Roman Regular

8 pt/ 9,6 pt - CO M0 Y0 K100


<http://dx.doi.org/10.1590/bjpt-rbf.2014.0095>

Cambria Regular

7 pt/ 8 pt - CO M0 Y0 K80

6 mm

editora cubo  
soluções para o universo acadêmico



180 x 265 mm

common than over-reporting. Of the 64 elderly women who reported falls during the monthly monitoring, 32.8% denied having fallen in the self-reporting of previous falls. Of the 52 participants who reported no falls during the monthly monitoring, 5.76% reported falls in the self-reporting of previous falls. Similarly, of the 24 elderly women who reported recurrent falls during prospective monitoring, 50% denied recurrent falls in the retrospective self-report. Additionally, of the 92 participants who denied having fallen two or more times during the prospective monitoring, 1.1% reported recurrent falls in the self-report of the 12 previous months.

However, the ratio of elderly women who incorrectly reported not having fallen in the previous 12 months decreased with the increase in the number of falls during the year of retrospective follow-up: 40% under-reporting among participants who reported one fall, 30.76% among those who reported two falls, 16.6% among those who reported three falls, and no under-reporting among those who had fallen four to six times during the study.

**Table 3.** Agreement between prospective monitoring and retrospective self-reporting of falls and recurrent falls over 12 months (n=116).

Retrospective Self-report	Prospective Monitoring		Total
	Non-faller (0 fall)	Faller (≥1 fall)	
Non-faller (0 fall)	49	21	70
Faller (≥ 1 fall)	3	43	46
Total	52	64	116

	Non-recurrent faller (≤1 fall)	Recurrent faller (≥2 falls)	Total
Non-recurrent faller (≤1 fall)	91	12	103
Recurrent faller (≥2 falls)	1	12	13
Total	92	24	116


Retrospective self-report of falls: Sensitivity = 43/64 = 67.2% (CI 95% 55.0 – 77.4). Specificity = 49/52 = 94.2% (CI 95% 84.4 – 98.0). Percentage agreement = (49 + 43)/116 = 79.31%. *Kappa* = 0.595, p=0.001 (CI 95% 0.458 – 0.732). Retrospective self-reporting of recurrent falls: Sensitivity = 12/24 = 50% (CI 95% 31.4 – 68.6). Specificity = 91/92 = 98.9% (CI 95% 94.1 – 99.8). Percentage agreement = (91 + 12)/116 = 88.79%. *Kappa* = 0.589, p=0.001 (CI 95% 0.395-0.783).

**Table 4.** Sensitivity and specificity of retrospective self-reporting of falls and of recurrent falls for sample subgroups.

Subgroups	Falls (≥1 fall)		Recurrent Falls (≥2 falls)	
	Sensitivity (CI 95%)	Specificity (CI 95%)	Sensitivity (CI 95%)	Specificity (CI 95%)
No injuries after falls (n=19)	63.1% (41.0 – 80.8)	0%	60% (23.1 – 88.2)	92.8% (68.5 – 98.7)
Injuries after falls (n=44)	68.2% (53.4 – 80.0)	0%	47.4% (27.3 – 68.3)	100% (86.7 – 100.0)

### Comparison between retrospective self-reporting and prospective monitoring of falls for sample subgroups

The analyses of the sensitivity and specificity of retrospective self-reporting of falls and recurrent falls with respect to prospective monitoring regarding subgroups of elderly women who may or may not have suffered injurious falls are shown in Table 4. A better recall capacity of falls in the previous



**Figure 1.** Bland-Altman diagram: comparison of prospective monthly monitoring over 12 months and annual retrospective self-reporting (previous 12 months). UL = upper limit; LL = Lower Limit; Mean difference = 0.35. SD = 0.83. Limit of agreement = 0.35±1.66.

recommend a more simple definition of falls to allow for a broader understanding<sup>11</sup> allowing more attention to be paid to the correct understanding of the definition by patients. Thus, the proper recognition of the patient’s history of falls would allow clinicians and researchers to develop strategies to reduce the incidence of falls and injuries and to preserve mobility among elderly patients<sup>14</sup>, especially among those at high risk of falls and fractures.

● Conclusion

The method of retrospective self-reporting of falls in the previous 12 months exhibited moderate agreement and limited accuracy with respect to the method of prospective monitoring of falls among elderly women at high risk of falls and fractures. The retrospective self-reporting of falls was more sensitive with respect to prospective monitoring among women with injurious falls than among those with no post-fall injuries. These findings call for caution when substituting monthly monitoring with retrospective questioning and indicate the importance of including associated clinical-functional information for decision-making in clinical-scientific settings.

● References

1. Ganz DA, Bao Y, Shekelle PG, Rubenstein LZ. Will my patient fall? JAMA. 2007;297(1):77-86. <http://dx.doi.org/10.1001/jama.297.1.77>. PMID:17200478.
2. Cruz DT, Ribeiro LC, Vieira MT, Teixeira MT, Bastos RR, Leite IC. Prevalência de quedas e fatores associados em idosos. Rev Saude Publica. 2012;46(1):138-46. <http://dx.doi.org/10.1590/S0034-89102011005000087>. PMID:22183513.
3. Silva RB, Costa-Paiva L, Oshima MM, Morais SS, Pinto-Neto AM. Frequência de quedas e associação com parâmetros estabilométricos em mulheres na pós-menopausa com e sem osteoporose. Rev Bras Ginecol Obstet. 2009;31(10):496-502. <http://dx.doi.org/10.1590/S0100-72032009001000005>. PMID:19942997.
4. Swanenburg J, Bruin ED, Uebelhart D, Mulder T. Falls prediction in elderly people: a 1-year prospective study. Gait Posture. 2010;31(3):317-21. <http://dx.doi.org/10.1016/j.gaitpost.2009.11.013>. PMID:20047833.
5. Sai AJ, Gallagher JC, Smith LM, Logsdon S. Fall predictors in the community dwelling elderly: a cross sectional and prospective cohort study. J Musculoskelet Neuronal Interact. 2010;10(2):142-50. PMID:20516631.
6. Peel N. Validating recall of falls by older people. Accid Anal Prev. 2000;32(3):371-2. [http://dx.doi.org/10.1016/S0001-4575\(99\)00066-4](http://dx.doi.org/10.1016/S0001-4575(99)00066-4). PMID:10776852.

7. Ersoy Y, MacWalter RS, Durmus B, Altay ZE, Baysal O. Predictive effects of different clinical balance measures and the fear of falling on falls in postmenopausal women aged 50 years and over. Gerontology. 2009;55(6):660-5. <http://dx.doi.org/10.1159/000235652>. PMID:19690394.
8. Kunkel D, Pickering RM, Ashburn AM. Comparison of retrospective interviews and prospective diaries to facilitate fall reports among people with stroke. Age Ageing. 2011;40(2):277-80. <http://dx.doi.org/10.1093/ageing/afq177>. PMID:21242191.
9. Alexandre TS, Meira DM, Rico NC, Mizuta SK. Accuracy of Timed Up and Go Test for screening risk of falls among community-dwelling elderly. Rev Bras Fisioter. 2012;16(5):381-8. <http://dx.doi.org/10.1590/S1413-35552012005000041>. PMID:22858735.
10. Perracini MR, Teixeira LF, Ramos JL, Pires RS, Najas MS. Fall-related factors among less and more active older outpatients. Rev Bras Fisioter. 2012;16(2):166-72. <http://dx.doi.org/10.1590/S1413-35552012005000009>. PMID:22378477.
11. Hauer K, Lamb SE, Jorstad EC, Todd C, Becker C, PROFANE-Group. Systematic review of definitions and methods of measuring falls in randomised controlled fall prevention trials. Age Ageing. 2006;35(1):5-10. <http://dx.doi.org/10.1093/ageing/afi218>. PMID:16364930.
12. Hannan MT, Gagnon MM, Aneja J, Jones RN, Cupples LA, Lipsitz LA, et al. Optimizing the tracking of falls in studies of older participants: comparison of quarterly telephone recall with monthly falls calendars in the MOBILIZE Boston Study. Am J Epidemiol. 2010;171(9):1031-6. <http://dx.doi.org/10.1093/aje/kwq024>. PMID:20360242.
13. Ganz DA, Higashi T, Rubenstein LZ. Monitoring falls in cohort studies of community-dwelling older people: effect of the recall interval. J Am Geriatr Soc. 2005;53(12):2190-4. <http://dx.doi.org/10.1111/j.1532-5415.2005.00509.x>. PMID:16398908.
14. Hale WA, Delaney MJ, Cable T. Accuracy of patient recall and chart documentation of falls. J Am Board Fam Pract. 1993;6(3):239-42. PMID:8503294.
15. Sanders KM, Hayles AL, Kotowicz MA, Nicholson GC. Monitoring falls in cohort studies of community-dwelling older women. J Am Geriatr Soc. 2009;57(4):733-4. <http://dx.doi.org/10.1111/j.1532-5415.2009.02205.x>. PMID:19392967.
16. Mackenzie L, Byles J, D’Este C. Validation of self-reported fall events in intervention studies. Clin Rehabil. 2006;20(4):331-9. <http://dx.doi.org/10.1191/0269215506cr947oa>. PMID:16719031.
17. Pinto-Neto AM, Soares A, Urbanetz AA, Souza ACA, Ferrari AEM, Amaral B, et al. Consenso Brasileiro de Osteoporose 2002. Rev Bras Reumatol. 2002;42(6):343-54.
18. Organização Mundial de Saúde - OMS. Envelhecimento ativo: uma política de saúde. Brasília; 2005.
19. Chaves MLF, Godinho CC, Porto CS, Mansur L, Carthery-Goulart MT, Yassuda MS, et al. Cognitive, funcitonal and behavioral assessment. Dement Neuropsychol. 2011;5(3):153-66.
20. Brucki SM, Nitrini R, Caramelli P, Bertolucci PH, Okamoto IH. Sugestões para o uso do mini-exame do estado mental no Brasil. Arq Neuropsiquiatr. 2003;61(3B):777-81. <http://dx.doi.org/10.1590/S0004-282X2003000500014>. PMID:14595482.
21. Neri AL, Ongaratto LL, Yassuda MS. Mini-Mental State Examination sentence writing among community-dwelling

