
Cornell notes Turner style
Instructions

- Cut out the notes on the right hand side and **paste** into the **right hand side** of your notebook page.
- **Include** the **vertical black line or redraw this line** – giving yourself a large left hand margin
- **Read** the notes several times
- **Ask** yourself **three questions**:
  1. Can I recognise all the content (information) in the notes?
  2. In the notes is there a process or procedure I need to know?
  3. Do I understand the idea/meaning/argument outlined in the notes. That is: do I "get it"?
- **Do the following**
  - **a.** Within the notes themselves, **underline** all the **content** you should learn or memorise.
  - **b.** In the left margin **label** each section of the notes as either a **definition**, **example**, **explanation**, **process**, or **evidence** (and you could make up your own categories)
  - **c.** With each label in step b, write a key word or phrase which tells us what the label is referring to. An example would be…




(note – the example is not for these notes
- The key words are highlighted)

  - **d.** At the end of the notes **write a summary underneath the notes**. The summary should be **short** – it is to **demonstrate your understanding** of the notes, not rehash the content. Often you should try to draw a diagram or mind map.

Sex Linked Inheritance


Sex-linked inheritance involves genes present on a specific portion of the sex chromosomes (the 23rd pair).

Male =



If a gene is present on this part of the sex chromosome, a male can only have 1 allele, not two.

Female =



Females being two X chromosomes can have two alleles (normal pattern of inheritance).

This has implications in predicting inheritance as the male will possess only one allele – males have no backup copy of the gene. Males are therefore more susceptible to diseases or conditions caused by recessive genes on this part of the X chromosome, e.g. haemophilia, colour blindness.

Example:

Haemophilia is a disease caused by a sex-linked recessive gene. If a woman who is heterozygous for haemophilia (that is, she does not have haemophilia but is a carrier) has a child with a man with haemophilia. What are the chances any son or daughter will have the diseases?

* Punnet squares are still used, but used differently and <u>interpreted differently</u>.

Male \ Female	X_H	X_h
X_H	$X_H X_h$	$X_H X_h$
Y	$X_H Y$	$X_h Y$

- **<u>Daughters</u>** (XX) – 100% heterozygous not haemophilia.
- **<u>Males</u>** (XY)
  50% recessive phenotype (haemophilia)
  50% dominant phenotype