

# Markov-switching (discrete-time) Hawkes process

S. Robin

joint work with A. Bonnet

LPSM, Sorbonne université

STOR-i, Lancaster, Jun. 2025

# 'Motivation'

## Counting process


Overnight recording of bat cries in continuous time



# 'Motivation'

## Counting process

Overnight recording of bat cries in continuous time




# 'Motivation'

## Counting process

Overnight recording of bat cries in continuous time

- Can we detect changes in the distribution of events?




# 'Motivation'

## Counting process

Overnight recording of bat cries in continuous time

- ▶ Can we detect changes in the distribution of events?
- ▶ Can we associate each time period with some underlying behavior?




# 'Motivation'

## Counting process

Overnight recording of bat cries in continuous time

- ▶ Can we detect changes in the distribution of events?
- ▶ Can we associate each time period with some underlying behavior?




Modelling. Latent Markov switching process.

# Point process

# Point process

Reminder.




- ▶  $(T_k)_{k \geq 1}$  a random collection of points
- ▶ Count process  $H(t) = \sum_{k \geq 1} \mathbb{I}\{T_k \leq t\}$
- ▶ Intensity function  $\lambda(t)$ : immediate probability of observing an event at time  $t$



# Point process

Reminder.



- ▶  $(T_k)_{k \geq 1}$  a random collection of points
- ▶ Count process  $H(t) = \sum_{k \geq 1} \mathbb{I}\{T_k \leq t\}$
- ▶ Intensity function  $\lambda(t)$ : immediate probability of observing an event at time  $t$

## Examples

- ▶ Homogeneous Poisson process:  $\lambda(t) \equiv \lambda$
- ▶ Heterogeneous Poisson process:  $\lambda(t) = \text{deterministic function}$
- ▶ Hawkes process:  $\lambda(t) = \text{function of the past events} = \text{random function}$

# Outline

## (Discrete) Hawkes process

### Continuous-time Hawkes process

#### Discrete-time Hawkes process

#### Markovian representation

## Discrete Markov switching Hawkes process

### Model


### Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

# Univariate Hawkes process



(Conditional) intensity function for the Hawkes process [Haw71]:

$$\lambda(t) = \lambda(t \mid \mathcal{H}_t) = \lambda_0 + \sum_{T_k < t} h(t - T_k)$$

- ▶  $\lambda_0$  = baseline
- ▶  $h$  = kernel = influence of past events

## Self-exciting exponential Hawkes process


$$\lambda(t) = \lambda_0 + \sum_{T_k < t} a e^{-b(t-T_k)}$$

**Self exciting:** Each event increases the probability of observing another event

# Self-exciting exponential Hawkes process


$$\lambda(t) = \lambda_0 + \sum_{T_k < t} a e^{-b(t-T_k)}$$

**Self exciting:** Each event increases the probability of observing another event



- ▶ Exponential kernel function  $h(t) = a e^{-bt}$
- ▶  $a \geq 0$  to ensure that  $\lambda$  is non negative
- ▶  $a/b < 1$  to ensure stationarity
- ▶ Applications: sismology, epidemiology, vulcanology, neurosciences, ecology, ...

## Cluster representation [HO74]



- ▶ Immigrants arrive at rate  $\lambda_0$
- ▶ Each immigrant or descendant produces new individuals at rate  $h(t - T)$

# Outline

## (Discrete) Hawkes process

Continuous-time Hawkes process

**Discrete-time Hawkes process**

Markovian representation

## Discrete Markov switching Hawkes process

Model

Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

# Discrete-time Hawkes process

## Continuous time exponential Hawkes process

$$\lambda(t) = \lambda_0 + \sum_{T_k < t} a e^{-b(t-T_k)}$$




# Discrete-time Hawkes process

## Continuous time exponential Hawkes process

$$\lambda(t) = \lambda_0 + \sum_{T_k < t} a e^{-b(t-T_k)}$$

## Discretization [Seo15,Kir16,Kir17]

- ▶  $I_k = [\tau_{k-1}; \tau_k]$  with  $\tau_k = k\Delta$
- ▶  $H_k = H(I_k)$  the number of events on  $I_k$



- ▶ Distribution of  $(H_k)_{k \geq 1}$ ?

# Decomposition of the count

$H_k$  = number of events on  $I_k = [\tau_{k-1}; \tau_k]$

$$H_k \triangleq B_k + \sum_{\ell \leq k-1} \sum_{T \in I_\ell} M_T(I_k) + R_k$$

where

- ▶  $B_k$  = number of immigrants within  $I_k$ :

$$B_k \sim \mathcal{P}(\mu)$$

with  $\mu = \lambda_0 \Delta$ ,

- ▶  $M_T(I_k)$  = number of descendants descendants of  $T < \tau_k$  within  $I_k$ :

$$M_T(I_k) \sim \mathcal{P} \left( \int_{I_k} a e^{-b(t-T)} dt \right) = \mathcal{P} \left( \alpha e^{-b(\tau_k - T)} \right)$$

with  $\alpha = a(e^{b\Delta} - 1)/b$ ,

- ▶  $R_k$  = number of descendants of points  $T \in I_k$  within  $I_k$

# Discrete time Hawkes process

When  $\Delta$  is small:

- ▶  $R_k \simeq 0$
- ▶ For  $T \in I_\ell$ :  $e^{-b(\tau_k - T)} \simeq e^{-b(\tau_k - \tau_\ell)} = \beta^{k-\ell}$  with  $\beta = e^{-b\Delta}$ , so

$$\sum_{\ell \leq k-1} \sum_{T \in I_\ell} M_T(I_k) \stackrel{\Delta}{\simeq} \sum_{\ell \leq k-1} \sum_{T \in I_\ell} \mathcal{P}(\alpha \beta^{k-\ell}) \stackrel{\Delta}{=} \mathcal{P} \left( \sum_{\ell \leq k-1} H_{k-\ell} \alpha \beta^{\ell-1} \right)$$

# Discrete time Hawkes process

When  $\Delta$  is small:

- ▶  $R_k \simeq 0$
- ▶ For  $T \in I_\ell$ :  $e^{-b(\tau_k - T)} \simeq e^{-b(\tau_k - \tau_\ell)} = \beta^{k-\ell}$  with  $\beta = e^{-b\Delta}$ , so

$$\sum_{\ell \leq k-1} \sum_{T \in I_\ell} M_T(I_k) \stackrel{\Delta}{\simeq} \sum_{\ell \leq k-1} \sum_{T \in I_\ell} \mathcal{P}(\alpha \beta^{k-\ell}) \stackrel{\Delta}{=} \mathcal{P} \left( \sum_{\ell \leq k-1} H_{k-\ell} \alpha \beta^{\ell-1} \right)$$

Discrete-time Hawkes process  $Y = \{Y_k\}_{k \leq 1}$ .

$$Y_k \mid (Y_\ell)_{\ell \leq k-1} \sim \mathcal{P} \left( \mu + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

See [Kir16] for the convergence toward a continuous-time Hawkes process.

# Outline

## (Discrete) Hawkes process

Continuous-time Hawkes process

Discrete-time Hawkes process

Markovian representation

## Discrete Markov switching Hawkes process

Model

Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

# Markovian representation

Discrete-time Hawkes process  $Y = \{Y_k\}_{k \leq 1}$ .

$$Y_k \mid (Y_\ell)_{\ell \leq k-1} \sim \mathcal{P} \left( \mu + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

→  $(Y_k)_{k \geq 1}$  is not a Markov chain (infinite memory).

# Markovian representation

Discrete-time Hawkes process  $Y = \{Y_k\}_{k \leq 1}$ .

$$Y_k \mid (Y_\ell)_{\ell \leq k-1} \sim \mathcal{P} \left( \mu + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

$\rightarrow (Y_k)_{k \geq 1}$  is not a Markov chain (infinite memory).

Markovian representation.

# Markovian representation

Discrete-time Hawkes process  $Y = \{Y_k\}_{k \leq 1}$ .

$$Y_k \mid (Y_\ell)_{\ell \leq k-1} \sim \mathcal{P} \left( \mu + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

→  $(Y_k)_{k \geq 1}$  is not a Markov chain (infinite memory).

Markovian representation.

► Define

$$U_1 = 0, \quad U_k = \sum_{\ell=1}^k \alpha \beta^{\ell-1} Y_{k-\ell},$$



# Markovian representation

Discrete-time Hawkes process  $Y = \{Y_k\}_{k \leq 1}$ .

$$Y_k \mid (Y_\ell)_{\ell \leq k-1} \sim \mathcal{P} \left( \mu + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

→  $(Y_k)_{k \geq 1}$  is not a Markov chain (infinite memory).

Markovian representation.

- Define

$$U_1 = 0, \quad U_k = \sum_{\ell=1}^k \alpha \beta^{\ell-1} Y_{k-\ell},$$

- we have for  $k \geq 1$  (with  $U_0 = Y_0 = 0$ )

$$U_k = \alpha Y_{k-1} + \beta U_{k-1}, \quad Y_k \mid U_k \sim \mathcal{P}(\mu + U_k).$$

# Markovian representation

Discrete-time Hawkes process  $Y = \{Y_k\}_{k \leq 1}$ .

$$Y_k \mid (Y_\ell)_{\ell \leq k-1} \sim \mathcal{P} \left( \mu + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

→  $(Y_k)_{k \geq 1}$  is not a Markov chain (infinite memory).

Markovian representation.

- Define

$$U_1 = 0, \quad U_k = \sum_{\ell=1}^k \alpha \beta^{\ell-1} Y_{k-\ell},$$

- we have for  $k \geq 1$  (with  $U_0 = Y_0 = 0$ )

$$U_k = \alpha Y_{k-1} + \beta U_{k-1}, \quad Y_k \mid U_k \sim \mathcal{P}(\mu + U_k).$$

→  $((Y_k, U_k))_{k \geq 1}$  forms a Markov Chain.

# Graphical model

Discrete time Hawkes process.

$$(Y_k)_{k \geq 1} \sim \text{Discrete Hawkes}(\mu, \alpha, \beta)$$

$$U_1 = 0, \quad U_k = \alpha Y_{k-1} + \beta U_{k-1}, \quad Y_k \sim \mathcal{P}(\mu + U_k)$$


# Graphical model

Discrete time Hawkes process.

$$(Y_k)_{k \geq 1} \sim \text{Discrete Hawkes}(\mu, \alpha, \beta)$$

$$U_1 = 0, \quad U_k = \alpha Y_{k-1} + \beta U_{k-1}, \quad Y_k \sim \mathcal{P}(\mu + U_k)$$

Graphical model:



$(U_k)_{k \geq 1} = \text{memory}$ ,  $(Y_k)_{k \geq 1} = \text{observed process}$ .

$$\begin{aligned} p(U_k, Y_k \mid (U_\ell, Y_\ell)_{\ell \leq k-1}) &= p(U_k, Y_k \mid U_{k-1}, Y_{k-1}) \\ &= p(U_k \mid U_{k-1}, Y_{k-1}) p(Y_k \mid U_k) \end{aligned}$$

# Outline

## (Discrete) Hawkes process

Continuous-time Hawkes process

Discrete-time Hawkes process

Markovian representation

## Discrete Markov switching Hawkes process

Model

Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

# Discrete time Hawkes HMM

Model:  $Q$  hidden states

- ▶ Hidden path:  $(Z_k)_{k \geq 1}$  homogeneous Markov chain with  $Q$  states, transition matrix  $\pi$  and initial distribution  $\nu$ :

$$(Z_k)_{k \geq 1} \sim MC_Q(\nu, \pi)$$

# Discrete time Hawkes HMM

**Model:**  $Q$  hidden states

- ▶ Hidden path:  $(Z_k)_{k \geq 1}$  homogeneous Markov chain with  $Q$  states, transition matrix  $\pi$  and initial distribution  $\nu$ :

$$(Z_k)_{k \geq 1} \sim MC_Q(\nu, \pi)$$

- ▶ Observed counts: for  $k \geq 1$  and

$$(Y_k \mid (Y_\ell)_{\ell \leq k-1}, Z_k = q) \sim \mathcal{P} \left( \mu_q + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

or, for  $k \geq 1$  (with  $U_0 = Y_0 = 0$ )

$$U_k = \alpha Y_{k-1} + \beta U_{k-1},$$

$$Y_k \mid U_k \sim \mathcal{P}(\mu_{Z_k} + U_k)$$

# Discrete time Hawkes HMM

**Model:**  $Q$  hidden states

- ▶ Hidden path:  $(Z_k)_{k \geq 1}$  homogeneous Markov chain with  $Q$  states, transition matrix  $\pi$  and initial distribution  $\nu$ :

$$(Z_k)_{k \geq 1} \sim MC_Q(\nu, \pi)$$

- ▶ Observed counts: for  $k \geq 1$  and

$$(Y_k \mid (Y_\ell)_{\ell \leq k-1}, Z_k = q) \sim \mathcal{P} \left( \mu_q + \sum_{\ell=1}^{k-1} \alpha \beta^{\ell-1} Y_{k-\ell} \right)$$

or, for  $k \geq 1$  (with  $U_0 = Y_0 = 0$ )

$$U_k = \alpha Y_{k-1} + \beta U_{k-1},$$

$$Y_k \mid U_k \sim \mathcal{P}(\mu_{Z_k} + U_k)$$


**Assumptions:**

- ▶ The immigration rate  $\mu$  varies with the hidden state
- ▶ The distribution of the number of offspring  $(\alpha, \beta)$  does not vary with the hidden state



# Discrete time Hawkes HMM


Graphical model:



$(Z_k)_{k \geq 1}$  = **hidden path**,  $(U_k)_{k \geq 1}$  = **memory**,  $(Y_k)_{k \geq 1}$  = **observed process**.

# Discrete time Hawkes HMM

## Graphical model:



$(Z_k)_{k \geq 1}$  = **hidden path**,  $(U_k)_{k \geq 1}$  = **memory**,  $(Y_k)_{k \geq 1}$  = **observed process**.

## Remarks:

- ▶ The memory of the past is 'stored' in the variable  $U_k$ , which can still be computed recursively ( $U_k = \alpha Y_{k-1} + \beta U_{k-1}$ )
- ▶ The Markovian property still holds if the influence of the past varies with the hidden state ( $\alpha \rightarrow \alpha_k, \beta \rightarrow \beta_k$ ).

# Outline

## (Discrete) Hawkes process

Continuous-time Hawkes process

Discrete-time Hawkes process

Markovian representation

## Discrete Markov switching Hawkes process

Model

Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

# Identifiability

**Proposition:** The model parameter  $\theta = (\nu, \pi, (\mu_q)_{1 \leq q \leq Q}, \alpha, \beta)$  is identifiable from the joint distribution  $p_\theta^{Y_1, Y_2, Y_3}$ :

$$\theta' \neq \theta \quad \Rightarrow \quad p_{\theta'}^{Y_1, Y_2, Y_3} \neq p_\theta^{Y_1, Y_2, Y_3}.$$

---

<sup>1</sup>The generic technique from [AMR09] does not apply here.

# Identifiability

**Proposition:** The model parameter  $\theta = (\nu, \pi, (\mu_q)_{1 \leq q \leq Q}, \alpha, \beta)$  is identifiable from the joint distribution  $p_{\theta}^{Y_1, Y_2, Y_3}$ :

$$\theta' \neq \theta \quad \Rightarrow \quad p_{\theta'}^{Y_1, Y_2, Y_3} \neq p_{\theta}^{Y_1, Y_2, Y_3}.$$

**Sketch of proof.** Finite Poisson mixtures are identifiable [Tei61], so, because<sup>1</sup>

$$p_{\theta}^{Y_1, Y_2, Y_3}(x, y, z) = \sum_{1 \leq q, \ell, m \leq Q} \nu_q \pi_{q\ell} \pi_{\ell m} \mathcal{P}(x; \mu_q) \mathcal{P}(y; \mu_{\ell} + \alpha x) \mathcal{P}(z; \mu_m + \alpha \beta x + \alpha y),$$

---

<sup>1</sup>The generic technique from [AMR09] does not apply here.

# Identifiability

**Proposition:** The model parameter  $\theta = (\nu, \pi, (\mu_q)_{1 \leq q \leq Q}, \alpha, \beta)$  is identifiable from the joint distribution  $p_{\theta}^{Y_1, Y_2, Y_3}$ :

$$\theta' \neq \theta \quad \Rightarrow \quad p_{\theta'}^{Y_1, Y_2, Y_3} \neq p_{\theta}^{Y_1, Y_2, Y_3}.$$

**Sketch of proof.** Finite Poisson mixtures are identifiable [Tei61], so, because<sup>1</sup>

$$p_{\theta}^{Y_1, Y_2, Y_3}(x, y, z) = \sum_{1 \leq q, \ell, m \leq Q} \nu_q \pi_{q\ell} \pi_{\ell m} \mathcal{P}(x; \mu_q) \mathcal{P}(y; \mu_{\ell} + \alpha x) \mathcal{P}(z; \mu_m + \alpha \beta x + \alpha y),$$

1.  $\nu$  and  $\mu$  can be identified from  $p_{\theta}(Y_1)$ , [sum over  $y$  and  $z$ ]

---

<sup>1</sup>The generic technique from [AMR09] does not apply here.

# Identifiability

**Proposition:** The model parameter  $\theta = (\nu, \pi, (\mu_q)_{1 \leq q \leq Q}, \alpha, \beta)$  is identifiable from the joint distribution  $p_{\theta}^{Y_1, Y_2, Y_3}$ :

$$\theta' \neq \theta \quad \Rightarrow \quad p_{\theta'}^{Y_1, Y_2, Y_3} \neq p_{\theta}^{Y_1, Y_2, Y_3}.$$

**Sketch of proof.** Finite Poisson mixtures are identifiable [Tei61], so, because<sup>1</sup>

$$p_{\theta}^{Y_1, Y_2, Y_3}(x, y, z) = \sum_{1 \leq q, \ell, m \leq Q} \nu_q \pi_{q\ell} \pi_{\ell m} \mathcal{P}(x; \mu_q) \mathcal{P}(y; \mu_{\ell} + \alpha x) \mathcal{P}(z; \mu_m + \alpha \beta x + \alpha y),$$

1.  $\nu$  and  $\mu$  can be identified from  $p_{\theta}(Y_1)$ ,      [sum over  $y$  and  $z$ ]
2. then  $\alpha$  can be identified from  $p_{\theta}(Y_2 \mid Y_1 = 1)$ ,      [fix  $x = 1$ , sum over  $z$ ]

---

<sup>1</sup>The generic technique from [AMR09] does not apply here.

# Identifiability

**Proposition:** The model parameter  $\theta = (\nu, \pi, (\mu_q)_{1 \leq q \leq Q}, \alpha, \beta)$  is identifiable from the joint distribution  $p_{\theta}^{Y_1, Y_2, Y_3}$ :

$$\theta' \neq \theta \quad \Rightarrow \quad p_{\theta'}^{Y_1, Y_2, Y_3} \neq p_{\theta}^{Y_1, Y_2, Y_3}.$$

**Sketch of proof.** Finite Poisson mixtures are identifiable [Tei61], so, because<sup>1</sup>

$$p_{\theta}^{Y_1, Y_2, Y_3}(x, y, z) = \sum_{1 \leq q, \ell, m \leq Q} \nu_q \pi_{q\ell} \pi_{\ell m} \mathcal{P}(x; \mu_q) \mathcal{P}(y; \mu_{\ell} + \alpha x) \mathcal{P}(z; \mu_m + \alpha \beta x + \alpha y),$$

1.  $\nu$  and  $\mu$  can be identified from  $p_{\theta}(Y_1)$ , [sum over  $y$  and  $z$ ]
2. then  $\alpha$  can be identified from  $p_{\theta}(Y_2 \mid Y_1 = 1)$ , [fix  $x = 1$ , sum over  $z$ ]
3. then  $\beta$  can be identified from  $p_{\theta}(Y_3 \mid Y_1 = 1, Y_2 = 0)$ , [fix  $x = 1, y = 0$ ]

<sup>1</sup>The generic technique from [AMR09] does not apply here.



# Identifiability

**Proposition:** The model parameter  $\theta = (\nu, \pi, (\mu_q)_{1 \leq q \leq Q}, \alpha, \beta)$  is identifiable from the joint distribution  $p_\theta^{Y_1, Y_2, Y_3}$ :

$$\theta' \neq \theta \quad \Rightarrow \quad p_{\theta'}^{Y_1, Y_2, Y_3} \neq p_\theta^{Y_1, Y_2, Y_3}.$$

**Sketch of proof.** Finite Poisson mixtures are identifiable [Tei61], so, because<sup>1</sup>

$$p_\theta^{Y_1, Y_2, Y_3}(x, y, z) = \sum_{1 \leq q, \ell, m \leq Q} \nu_q \pi_{q\ell} \pi_{\ell m} \mathcal{P}(x; \mu_q) \mathcal{P}(y; \mu_\ell + \alpha x) \mathcal{P}(z; \mu_m + \alpha \beta x + \alpha y),$$

1.  $\nu$  and  $\mu$  can be identified from  $p_\theta(Y_1)$ ,      [sum over  $y$  and  $z$ ]
2. then  $\alpha$  can be identified from  $p_\theta(Y_2 \mid Y_1 = 1)$ ,      [fix  $x = 1$ , sum over  $z$ ]
3. then  $\beta$  can be identified from  $p_\theta(Y_3 \mid Y_1 = 1, Y_2 = 0)$ ,      [fix  $x = 1, y = 0$ ]
4. then  $\pi$  can be identified from the joint mixture      [sum over  $z$ ]

$$p_\theta^{Y_1, Y_2}(x, y) = \sum_{1 \leq q, \ell \leq Q} \nu_q \pi_{q\ell} \mathcal{P}(x; \mu_q) \mathcal{P}(y; \mu_\ell + \alpha x),$$

which is proven identifiable.

---

<sup>1</sup>The generic technique from [AMR09] does not apply here.

# Inference

**Aim:** Infer the parameter  $\theta$

$$\hat{\theta} = \arg \max_{\theta} \log p_{\theta}(Y)$$

# Inference

**Aim:** Infer the parameter  $\theta$

$$\hat{\theta} = \arg \max_{\theta} \log p_{\theta}(Y)$$

**EM algorithm for HMM:** [DLR77,CMR05]

$$\theta^{(h+1)} = \underbrace{\arg \max_{\theta}}_{\text{M step}} \underbrace{\mathbb{E}_{\theta^{(h)}}}_{\text{E step}} [\log p_{\theta}(Y, Z) \mid Y]$$

- ▶ E step: Evaluate  $Q(\theta \mid \theta^{(h)}) = \mathbb{E}_{\theta^{(h)}} [\log p_{\theta}(Y, Z) \mid Y]$  (forward-backward recursion)
- ▶ M step: Gradient descent, computing  $\nabla_{\theta} Q(\theta \mid \theta^{(h)})$  by recursion

# Inference

## Classification:

Marginal:

$$\hat{Z}_k = \arg \max_q P_{\hat{\theta}}\{Z_k = q \mid Y\},$$

Joint (Viterbi):

$$\hat{Z} = \arg \max_z P_{\hat{\theta}}\{Z = z \mid Y\}$$

# Inference

## Classification:

$$\text{Marginal:} \quad \hat{Z}_k = \arg \max_q P_{\hat{\theta}}\{Z_k = q \mid Y\},$$

$$\text{Joint (Viterbi):} \quad \hat{Z} = \arg \max_z P_{\hat{\theta}}\{Z = z \mid Y\}$$

## Model selection: Penalized likelihood

$$AIC_Q = \log p_{\hat{\theta}_Q}(Y) - D_Q,$$

$$BIC_Q = \log p_{\hat{\theta}_Q}(Y) - D_Q \frac{\log(N)}{2}$$

with  $D_Q = \text{number of parameters} = 2 + Q^2$  and  $N = \text{number of time bins}$ .

# Outline

## (Discrete) Hawkes process

- Continuous-time Hawkes process

- Discrete-time Hawkes process

- Markovian representation

## Discrete Markov switching Hawkes process

- Model

- Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

## Simulation design ( $Q = 3$ )

- ▶ **Baseline continuous parameters:**  $m^0 = [10, 200, 1000]$ ,  $a^0 = 40$ ,  $b = 160$

## Simulation design ( $Q = 3$ )

- ▶ Baseline continuous parameters:  $m^0 = [10, 200, 1000]$ ,  $a^0 = 40$ ,  $b = 160$
- ▶ Increasing signal:  $\lambda = 0.5, 1, 1.5, 2$

$$a = \lambda a^0, \quad m = \lambda m^0.$$



## Simulation design ( $Q = 3$ )

► **Baseline continuous parameters:**  $m^0 = [10, 200, 1000]$ ,  $a^0 = 40$ ,  $b = 160$

► **Increasing signal:**  $\lambda = 0.5, 1, 1.5, 2$

$$a = \lambda a^0, \quad m = \lambda m^0.$$

► **Simulated process:**

$$(H_t)_{0 \leq t \leq 1} \sim \text{Heterogeneous } \textit{Continuous} \text{ Hawkes}(a, b^0, m)$$

## Simulation design ( $Q = 3$ )

- ▶ **Baseline continuous parameters:**  $m^0 = [10, 200, 1000]$ ,  $a^0 = 40$ ,  $b = 160$

- ▶ **Increasing signal:**  $\lambda = 0.5, 1, 1.5, 2$

$$a = \lambda a^0, \quad m = \lambda m^0.$$

- ▶ **Simulated process:**


$$(H_t)_{0 \leq t \leq 1} \sim \text{Heterogeneous } \textit{Continuous} \text{ Hawkes}(a, b^0, m)$$

- ▶ **Discretized process:**  $n = H(1)$

$$N = c n, \quad c = 0.5, 1, 2, 4,$$


$$Y_k = H\left(\left[\frac{k-1}{N}; \frac{k}{N}\right]\right), \quad k = 1, \dots, N.$$

→ not a discrete-time Hawkes process as defined earlier

Simulation results ( $Q^* = 3$ )Parameter estimates. Distribution of  $\hat{\theta} - \theta^*$  ( $Q = Q^*$ )


# Simulation results ( $Q^* = 3$ )

Model selection: BIC. Distribution of  $BIC_Q - BIC_1$




# Simulation results ( $Q^* = 3$ )

Model selection: AIC. Distribution of  $AIC_Q - AIC_1$



# Simulation results ( $Q^* = 3$ )

Classification. MAP / Viterbi (+ comput. time)



## Simulation conclusions

- ▶ Inference easier when more signal (large  $\lambda$ )!!!
- ▶ Inference easier with thinner discretization step (large  $N$ )  
But at the price of a higher computational cost
- ▶ BIC does not capture the right number of states  
Sequences not simulated according to the model
- ▶ AIC does, with reasonable signal ( $\lambda$ ) and discretization ( $N$ )  
Blind to the simulation shift from the model?

## Simulation conclusions

- ▶ Inference easier when more signal (large  $\lambda$ )!!!
- ▶ Inference easier with thinner discretization step (large  $N$ )  
But at the price of a higher computational cost
- ▶ BIC does not capture the right number of states  
Sequences not simulated according to the model
- ▶ AIC does, with reasonable signal ( $\lambda$ ) and discretization ( $N$ )  
Blind to the simulation shift from the model?

## Practical recommendations.

Take  $N = 2n$  and use AIC



# Outline

## (Discrete) Hawkes process

- Continuous-time Hawkes process

- Discrete-time Hawkes process

- Markovian representation

## Discrete Markov switching Hawkes process

- Model

- Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

# Bat cries

**Data set.** 1555 overnight recordings all over France

# Bat cries

**Data set.** 1555 overnight recordings all over France


**Poisson vs Hawkes / Homogeneous vs HMM.** Best model based on AIC

	Poisson	Hawkes	Total
Homogeneous	34	353	387
Hidden Markov	24	1144	1168
Total	58	1497	1555


- ▶ Memory (95%) and heterogeneity (75%) are present in most sequences
- ▶ Hawkes-HMM best fits almost 3 sequences out of 4.

# Example

Hawkes HMM ( $\hat{Q} = 3$ )




Poisson HMM ( $\hat{Q} = 4$ )



- ▶ Poisson-HMM needs many state changes to account for self-excitation
- ▶ Hawkes-HMM state changes do not correspond to slope changes

## States and species

The number of bat species was also recorded



- The number of states does not match the number of species

# Outline

## (Discrete) Hawkes process

- Continuous-time Hawkes process

- Discrete-time Hawkes process

- Markovian representation

## Discrete Markov switching Hawkes process

- Model

- Identifiability & Inference

## Simulation study

## Illustrations

## Discussion

# Summary

## What we did.

- ▶ The discretized Hawkes process with exponential kernel is a Markov model
  - ⇒ The discretized Markov switching Hawkes process with exponential kernel is a hidden Markov model
- ▶ The standard EM machinery applies to achieve maximum likelihood inference.
- ▶ Not shown: initialization based on existing estimation procedures for homogeneous Hawkes ([Che21],[CL22]) and Poisson HMM.

# Discussion

What we did not do.

- ▶ Goodness-of-fit: 'Poissonisation' (on-going).
- ▶ Model selection: derive a proper (BIC?) criterion accounting for the discretization step.
- ▶ Understand the inferred latent states in terms of animal behavior, biogeography, species, . . .



# Discussion













## What we did not do.

- ▶ Goodness-of-fit: 'Poissonisation' (on-going).
- ▶ Model selection: derive a proper (BIC?) criterion accounting for the discretization step.
- ▶ Understand the inferred latent states in terms of animal behavior, biogeography, species, . . .

## In parallel. With C. Dion-Blanc, D. Hawat and E. Lebarbier

- ▶ Efficient change-point detection ('segmentation') in (marked) Poisson & Hawkes processes.  
→ Dynamic programming applies [DBHLR24].
- ▶ Segmentation-classification of Poisson processes.

## References

-  Allman, C. Matias, and J.A. Rhodes. Identifiability of parameters in latent structure models with many observed variables. *The Annals of Statistics*, pages 3099–3132, 2009.
-  Cheysson. *hawkesbow: Estimation of Hawkes Processes from Binned Observations*, 2021. R package version 1.0.2.
-  Cheysson and G. Lang. Spectral estimation of hawkes processes from count data. *The Annals of Statistics*, 50(3):1722–1746, 2022.
-  Cappé, E. Moulines, and T. Rydén. *Inference in Hidden Markov Models*. Springer, 2005.
-  Orion-Blanc, D. Hawat, E Lebarbier, and S Robin. Multiple change-point detection for Poisson processes. Technical Report 2302.09103, arXiv, 2024.
-  Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B*, 39:1–38, 1977.
-  Hawkes. Spectra of some self-exciting and mutually exciting point processes. *Biometrika*, 58(1):83–90, 1971.
-  Hawkes and D. Oakes. A cluster process representation of a self-exciting process. *Journal of applied probability*, 11(3):493–503, 1974.
-  Kirchner. Hawkes and  $\text{INAR}(\infty)$  processes. *Stochastic Processes and their Applications*, 126(8):2494–2525, 2016.
-  Kirchner. An estimation procedure for the hawkes process. *Quantitative Finance*, 17(4):571–595, 2017.
-  Seol. Limit theorems for discrete hawkes processes. *Statistics & Probability Letters*, 99:223–229, 2015.
-  Teicher. Identifiability of mixtures. *The Annals of Mathematical Statistics*, 32(1):244–248, 1961.

# Discrete HMM

Conversion formulas from continuous to discrete Hawkes

$$\alpha = \frac{a(e^{b\Delta} - 1)}{b}, \quad \beta = e^{-b\Delta}$$

# Discrete HMM

Conversion formulas from continuous to discrete Hawkes


$$\alpha = \frac{a(e^{b\Delta} - 1)}{b}, \quad \beta = e^{-b\Delta}$$

## 3-step initialization

- ▶ Homogeneous Hawkes for the reproduction parameters  $\alpha$  and  $\beta$  (hawkesbow R package [Che21])
- ▶ Poisson-HMM for the rates  $\mu_1, \dots, \mu_Q$  and transition  $\pi$
- ▶ Correction  $\mu_k \rightarrow \tilde{\mu}_k$  to account for reproduction rate


# Simulation results ( $Q^* = 1$ , $N = cn$ , $m^0 = 400$ )

Parameter estimates. Distribution of  $\hat{\theta} - \theta^*$  ( $Q = Q^*$ )




# Simulation results ( $Q^* = 1$ , $N = cn$ , $m^0 = 400$ )

Model selection: BIC. Distribution of  $BIC_Q - BIC_1$




# Simulation results ( $Q^* = 1$ , $N = cn$ , $m^0 = 400$ )

Model selection: AIC. Distribution of  $AIC_Q - AIC_1$



# Simulation results ( $Q^* = 2$ , $N = cn$ , $m^0 = [10, 800]$ )


Parameter estimates. Distribution of  $\hat{\theta} - \theta^*$  ( $Q = Q^*$ )






# Simulation results ( $Q^* = 2$ , $N = cn$ , $m^0 = [10, 800]$ )

Model selection: BIC. Distribution of  $BIC_Q - BIC_1$




# Simulation results ( $Q^* = 2$ , $N = cn$ , $m^0 = [10, 800]$ )

Model selection: AIC. Distribution of  $AIC_Q - AIC_1$



# Simulation results ( $Q^* = 2$ , $N = cn$ , $m^0 = [10, 800]$ )

## Classification. MAP / Viterbi (+ comput. time)



## Model comparison for bat cries sequences

Poisson vs Hawkes / Homogeneous vs HMM. Best model based on BIC

	Poisson	Hawkes	Total
Homogeneous	132	775	907
Hidden Markov	21	627	648
Total	153	1402	1555

# States and locations

