

# Change-point detection in a Poisson process

S. Robin

joint work with E. Lebarbier, C. Dion-Blanc [DBLR23]

Sorbonne université

Stats au sommet, Rochebrune, Mar. 2022

# Example

Bat cries (night of the 17 jul. 2019)



## Example

Point process on  $t \in [0, 1]$ .


Event times:

$$0 < T_1 < \dots T_i < \dots T_n < 1$$

Counting process:

$$N(t) = \sum_{i=1}^n \mathbb{I}\{T_i \leq t\}$$

Bat cries (night of the 17 jul. 2019)<sup>a</sup>



<sup>a</sup>source: Vigie-Chiro program, Y. Bas, CESCO-MNHN

## Example

Point process on  $t \in [0, 1]$ .

Event times:

$$0 < T_1 < \dots T_i < \dots T_n < 1$$


Counting process:

$$N(t) = \sum_{i=1}^n \mathbb{I}\{T_i \leq t\}$$

Poisson Process.

$$\{N(t)\}_{0 \leq t \leq 1} \sim PP(\lambda(t))$$

Bat cries (night of the 17 jul. 2019)<sup>a</sup>



<sup>a</sup>source: Vigie-Chiro program, Y. Bas, CESCO-MNHN

Intensity function  $\lambda(t)$ :

$$\lambda(t) = \lim_{\Delta t \rightarrow 0} \frac{\mathbb{P}\{N(t + \Delta t) - N(t) = 1\}}{\Delta t},$$

$$\mathbb{E}N(s) - \mathbb{E}N(t) = \int_t^s \lambda(u) \, du$$

# Change-point detection

Piecewise constant intensity function.

Change-points

$$(\tau_0 =) 0 < \tau_1 \cdots < \tau_{K-1} < 1 (= \tau_K)$$


For  $t \in I_k = ]\tau_{k-1}, \tau_k]$ :

$$\lambda(t) = \lambda_k$$

→ Continuous piecewise linear cumulated intensity function

$$\Lambda(0, t) = \int_0^t \lambda(s) \, ds.$$

Bat cries (night of the 17 jul. 2019)<sup>a</sup>



<sup>a</sup>source: Vigie-Chiro program, Y. Bas, CESCO-MNHN

# Change-point detection

Piecewise constant intensity function.

Bat cries (night of the 17 jul. 2019)<sup>a</sup>

Change-points


$$(\tau_0 =) 0 < \tau_1 \cdots < \tau_{K-1} < 1 (= \tau_K)$$

For  $t \in I_k = ]\tau_{k-1}, \tau_k]$ :

$$\lambda(t) = \lambda_k$$

→ Continuous piecewise linear cumulated intensity function

$$\Lambda(0, t) = \int_0^t \lambda(s) ds.$$



<sup>a</sup>source: Vigie-Chiro program, Y. Bas, CESCO-MNHN

Aim.

- ▶ Segmentation: estimate  $(\tau, \lambda)$  in a reasonably fast manner
- ▶ Model selection: choose  $K$

# Change-point detection

Piecewise constant intensity function.

Kilauea eruptions

Change-points

$$(\tau_0 =) 0 < \tau_1 \cdots < \tau_{K-1} < 1 (= \tau_K)$$

For  $t \in I_k = ]\tau_{k-1}, \tau_k]$ :

$$\lambda(t) = \lambda_k$$

→ Continuous piecewise linear cumulated intensity function

$$\Lambda(0, t) = \int_0^t \lambda(s) \, ds.$$



Aim.

- ▶ Segmentation: estimate  $(\tau, \lambda)$  in a reasonably fast manner
- ▶ Model selection: choose  $K$

# Change-point detection

Piecewise constant intensity function.

Kilauea eruptions (from 1750 to 1984)<sup>a</sup>

Change-points


$$(\tau_0 =) 0 < \tau_1 \cdots < \tau_{K-1} < 1 (= \tau_K)$$

For  $t \in I_k = ]\tau_{k-1}, \tau_k]$ :

$$\lambda(t) = \lambda_k$$

→ Continuous piecewise linear cumulated intensity function

$$\Lambda(0, t) = \int_0^t \lambda(s) \, ds.$$



<sup>a</sup>source: [HB17]

Aim.

- ▶ Segmentation: estimate  $(\tau, \lambda)$  in a reasonably fast manner
- ▶ Model selection: choose  $K$



# Change-point detection

Piecewise constant intensity function.

Change-points

$$(\tau_0 =) 0 < \tau_1 \cdots < \tau_{K-1} < 1 (= \tau_K)$$


For  $t \in I_k = ]\tau_{k-1}, \tau_k]$ :

$$\lambda(t) = \lambda_k$$

→ Continuous piecewise linear cumulated intensity function

$$\Lambda(0, t) = \int_0^t \lambda(s) \, ds.$$

Kilauea eruptions (from 1750 to 1984)<sup>a</sup>



<sup>a</sup>source: [HB17]

Aim.

- ▶ Segmentation: estimate  $(\tau, \lambda)$  in a reasonably fast manner
- ▶ Model selection: choose  $K$

# Outline

## Estimation

## Model selection

## Illustrations

## Extensions

## Maximum-likelihood segmentation

First useful property of Poisson processes: Independence of disjoint intervals.

## Maximum-likelihood segmentation

First useful property of Poisson processes: Independence of disjoint intervals.

(Neg-log-)likelihood. Denoting

- ▶  $\Delta\tau_k$  the length of the  $k$ -th interval ( $= \tau_k - \tau_{k-1}$ ),
- ▶  $\Delta N_k$  the number of events within the  $k$ -th interval ( $= N(\tau_k) - N(\tau_{k-1})$ ):

$$-\log p_{\tau, \lambda}(N) = \sum_{k=1}^K \lambda_k \Delta\tau_k - \Delta N_k \log \lambda_k,$$

## Maximum-likelihood segmentation

First useful property of Poisson processes: Independence of disjoint intervals.

(Neg-log-)likelihood. Denoting

- ▶  $\Delta\tau_k$  the length of the  $k$ -th interval ( $= \tau_k - \tau_{k-1}$ ),
- ▶  $\Delta N_k$  the number of events within the  $k$ -th interval ( $= N(\tau_k) - N(\tau_{k-1})$ ):

$$-\log p_{\tau,\lambda}(N) = \sum_{k=1}^K \lambda_k \Delta\tau_k - \Delta N_k \log \lambda_k,$$

Additive contrast. General form = sum over the segments

$$\gamma(\tau, \lambda) = \sum_{k=1}^K C(\Delta N_k, \Delta\tau_k, \lambda_k)$$

## Maximum-likelihood segmentation

First useful property of Poisson processes: Independence of disjoint intervals.

(Neg-log-)likelihood. Denoting

- ▶  $\Delta\tau_k$  the length of the  $k$ -th interval ( $= \tau_k - \tau_{k-1}$ ),
- ▶  $\Delta N_k$  the number of events within the  $k$ -th interval ( $= N(\tau_k) - N(\tau_{k-1})$ ):

$$-\log p_{\tau, \lambda}(N) = \sum_{k=1}^K \lambda_k \Delta\tau_k - \Delta N_k \log \lambda_k,$$

Additive contrast. General form = sum over the segments

$$\gamma(\tau, \lambda) = \sum_{k=1}^K C(\Delta N_k, \Delta\tau_k, \lambda_k)$$

Optimization problem.

$$(\hat{\tau}, \hat{\lambda}) = \arg \min_{\tau \in \mathcal{T}^K, \lambda \in (\mathbb{R}^+)^K} \gamma(\tau, \lambda).$$

## Minimizing the contrast function

**Optimal  $\lambda$ .** Because the contrast is additive, we may define

$$\hat{\lambda}_k = \hat{\lambda}_k(\tau) = \arg \min_{\lambda_k \in \mathbb{R}^+} C(\Delta N_k, \Delta \tau_k, \lambda_k)$$

e.g.  $\hat{\lambda}_k = \Delta N_k / \Delta \tau_k$  if  $\gamma = -\log p_\theta$ .

## Minimizing the contrast function

**Optimal  $\lambda$ .** Because the contrast is additive, we may define

$$\hat{\lambda}_k = \hat{\lambda}_k(\tau) = \arg \min_{\lambda_k \in \mathbb{R}^+} C(\Delta N_k, \Delta \tau_k, \lambda_k)$$

e.g.  $\hat{\lambda}_k = \Delta N_k / \Delta \tau_k$  if  $\gamma = -\log p_\theta$ .

**Optimal  $\tau$ .** We are left with the minimization problem

$$\hat{\tau} = \arg \min_{\tau \in \mathcal{T}^K} \hat{\gamma}(\tau), \quad \text{where} \quad \hat{\gamma}(\tau) = \gamma(\tau, \hat{\lambda}(\tau))$$

where  $\mathcal{T}^K$  is the continuous segmentation space:

$$\mathcal{T} = \left\{ \tau \in [0, 1]^{K+1} : 0 = \tau_0 < \tau_1 < \dots < \tau_{K-1} < \tau_K = 1 \right\}.$$



## Minimizing the contrast function

**Optimal  $\lambda$ .** Because the contrast is additive, we may define

$$\hat{\lambda}_k = \hat{\lambda}_k(\tau) = \arg \min_{\lambda_k \in \mathbb{R}^+} C(\Delta N_k, \Delta \tau_k, \lambda_k)$$

e.g.  $\hat{\lambda}_k = \Delta N_k / \Delta \tau_k$  if  $\gamma = -\log p_\theta$ .

**Optimal  $\tau$ .** We are left with the minimization problem

$$\hat{\tau} = \arg \min_{\tau \in \mathcal{T}^K} \hat{\gamma}(\tau), \quad \text{where} \quad \hat{\gamma}(\tau) = \gamma(\tau, \hat{\lambda}(\tau))$$


where  $\mathcal{T}^K$  is the continuous segmentation space:

$$\mathcal{T} = \left\{ \tau \in [0, 1]^{K+1} : 0 = \tau_0 < \tau_1 < \dots < \tau_{K-1} < \tau_K = 1 \right\}.$$

**Main issue:** The contrast  $\hat{\gamma}(\tau)$  is **neither convex nor continuous** wrt  $\tau$ .

# Shape of the contrast fonction

Observed  $N(t)$ :  $n = 10$ ,





---

<sup>1</sup>gray borders come by pair

# Shape of the contrast fonction

Observed  $N(t)$ :  $n = 10$ ,




Contrast  $\hat{\gamma}(\tau)$  for  $K = 3$  segments:

$$\tau = (\tau_1, \tau_2).$$

One 'block' =  
one specific value for the vector <sup>1</sup>

$$\Delta N = (\Delta N_1, \Delta N_2, \Delta N_3)$$



<sup>1</sup>gray borders come by pair

## Partitioning the segmentation space

**Partitioning the number of events.** Define  $\mathcal{N}^K = \left\{ \nu \in \mathbb{N}^K : \sum_{k=1}^K \nu_k = n \right\}$ .

→  $\nu_k$  = given number of events in segment  $k$ .

## Partitioning the segmentation space

**Partitioning the number of events.** Define  $\mathcal{N}^K = \left\{ \nu \in \mathbb{N}^K : \sum_{k=1}^K \nu_k = n \right\}$ .

→  $\nu_k$  = given number of events in segment  $k$ .


**Partitioning the segmentation space.** For  $\nu \in \mathcal{N}_K$ , define

$$\mathcal{T}_\nu^K = \left\{ \tau \in \mathcal{T}^K : \Delta N = \nu \right\}.$$

→  $\mathcal{T}_\nu^K$  = set of segmentation satisfying the prescribed  $\nu = (\nu_1, \dots, \nu_K)$ .

We have

$$\min_{\tau \in \mathcal{T}^K} \hat{\gamma}(\tau) = \min_{\nu \in \mathcal{N}^K} \min_{\tau \in \mathcal{T}_\nu^K} \hat{\gamma}(\tau).$$



# Optimal segmentation

**Proposition 1.** If  $K \leq n$  and if  $\hat{\gamma}(\tau)$  is strictly concave wrt  $\tau \in \mathcal{T}_\nu^K$  for each  $\nu \in \mathcal{N}^K$ , then

$$\hat{\tau} = \arg \min_{\tau \in \mathcal{T}^K} \hat{\gamma}(\tau) \subset \{T_1^-, T_1, T_2^-, T_2, \dots, T_n^-, T_n\}.$$

# Optimal segmentation

**Proposition 1.** If  $K \leq n$  and if  $\hat{\gamma}(\tau)$  is strictly concave wrt  $\tau \in \mathcal{T}_\nu^K$  for each  $\nu \in \mathcal{N}^K$ , then

$$\hat{\tau} = \arg \min_{\tau \in \mathcal{T}^K} \hat{\gamma}(\tau) \subset \{T_1^-, T_1, T_2^-, T_2, \dots, T_n^-, T_n\}.$$

**Proposition 2.** If each  $\hat{C}(\nu_k, \Delta\tau_k) := C(\nu_k, \Delta\tau_k, \hat{\lambda}_k)$  is strictly concave wrt  $\Delta\tau_k$ ,  $\hat{\gamma}(\tau)$ , then is strictly concave wrt  $\tau \in \mathcal{T}_\nu^K$ .

## Optimal segmentation

**Proposition 1.** If  $K \leq n$  and if  $\hat{\gamma}(\tau)$  is strictly concave wrt  $\tau \in \mathcal{T}_\nu^K$  for each  $\nu \in \mathcal{N}^K$ , then

$$\hat{\tau} = \arg \min_{\tau \in \mathcal{T}^K} \hat{\gamma}(\tau) \subset \{T_1^-, T_1, T_2^-, T_2, \dots, T_n^-, T_n\}.$$

**Proposition 2.** If each  $\hat{C}(\nu_k, \Delta\tau_k) := C(\nu_k, \Delta\tau_k, \hat{\lambda}_k)$  is strictly concave wrt  $\Delta\tau_k$ ,  $\hat{\gamma}(\tau)$ , then is strictly concave wrt  $\tau \in \mathcal{T}_\nu^K$ .

**Consequence.**  $\hat{\tau}$  can be obtained by dynamic programming over the  $2n + 2$  possible change-points

$$\mathcal{S} = \{0, T_1^-, T_1, T_2^-, T_2, \dots, T_n^-, T_n, 1\}$$

with complexity at most  $O(n^2)$ .



## Admissible contrasts

**Poisson contrast.**  $\hat{C}_P(\nu_k, \Delta\tau_k) = \nu_k(1 - \log \nu_k + \log \Delta\tau_k)$  is concave wrt  $\Delta\tau$ .

## Admissible contrasts

**Poisson contrast.**  $\hat{C}_P(\nu_k, \Delta\tau_k) = \nu_k(1 - \log \nu_k + \log \Delta\tau_k)$  is concave wrt  $\Delta\tau$ .

**Poisson-Gamma model.** For each segment  $1 \leq k \leq K$ :

$$\Lambda_k \text{ iid } \sim \mathcal{G}\text{am}(a, b), \quad \{N(t)\}_{t \in I_k} \mid \Lambda_k \sim PP(\Lambda_k).$$

## Admissible contrasts

**Poisson contrast.**  $\hat{C}_P(\nu_k, \Delta\tau_k) = \nu_k(1 - \log \nu_k + \log \Delta\tau_k)$  is concave wrt  $\Delta\tau$ .

**Poisson-Gamma model.** For each segment  $1 \leq k \leq K$ :

$$\Lambda_k \text{ iid } \sim \mathcal{G}\text{am}(a, b), \quad \{N(t)\}_{t \in I_k} \mid \Lambda_k \sim PP(\Lambda_k).$$

Contrast for one segment:

$$C_{PG}(\Delta N_k, \Delta\tau_k) = \text{cst} - \log \Gamma(a + \Delta N_k) + (a + \Delta N_k) \log(b + \Delta\tau_k)$$

→ Strictly concave wrt  $\Delta\tau_k$ .

## Desirable contrast

**Remark.** The Poisson contrast  $\hat{C}_P(\nu_k, \Delta\tau_k) = \nu_k(1 - \log \nu_k + \log \Delta\tau_k)$  satisfies

$$\hat{C}_P(\nu_k = 1, \Delta\tau_k = 0) = -\infty.$$

- ▶ The optimal solution will involve segments with null length and containing only one event.
- ▶ 'Undesirable' contrast.

## Desirable contrast

**Remark.** The Poisson contrast  $\hat{C}_P(\nu_k, \Delta\tau_k) = \nu_k(1 - \log \nu_k + \log \Delta\tau_k)$  satisfies

$$\hat{C}_P(\nu_k = 1, \Delta\tau_k = 0) = -\infty.$$

- ▶ The optimal solution will involve segments with null length and containing only one event.
- ▶ 'Undesirable' contrast.

**Poisson-Gamma contrast.**  $C_{PG}(\nu_k, \Delta\tau_k) = -\log \Gamma(a + \nu_k) + (a + \nu_k) \log(b + \Delta\tau_k)$ .

- ▶ Satisfies the concavity property ( $\rightarrow$  admissible),
- ▶ but avoids segments with null length ( $\rightarrow$  desirable).

# Outline

Estimation

Model selection

Illustrations

Extensions

# Model selection

Second useful property of Poisson processes: Thinning.

- ▶  $\{N(t)\} \sim PP(\lambda(t))$
- ▶ Sample event times (with prob.  $v$ )
- ▶ Store the remaining events



# Model selection

Second useful property of Poisson processes: Thinning.

▶  $\{N(t)\} \sim PP(\lambda(t))$

▶ Sample event times (with prob.  $\nu$ )

▶ Store the remaining events



$$\{N^L(t)\} \sim PP(\nu\lambda(t)), \quad \{N^T(t)\} \sim PP((1-\nu)\lambda(t)), \quad \{N^L(t)\} \perp \{N^T(t)\}$$



# Model selection

Second useful property of Poisson processes: Thinning.

▶  $\{N(t)\} \sim PP(\lambda(t))$

▶ Sample event times (with prob.  $v$ )

▶ Store the remaining events



$$\{N^L(t)\} \sim PP(v\lambda(t)), \quad \{N^T(t)\} \sim PP((1-v)\lambda(t)), \quad \{N^L(t)\} \perp \{N^T(t)\}$$

**Consequence.** If  $\{N(t)\}_{0 \leq t \leq 1} \sim PP(\lambda(t))$ , with  $\lambda(t)$  piecewise constant with change-points  $\tau = (\tau_k)$  and intensities  $\lambda = (\lambda_k)$ , then

# Model selection

Second useful property of Poisson processes: Thinning.

▶  $\{N(t)\} \sim PP(\lambda(t))$

▶ Sample event times (with prob.  $\nu$ )

▶ Store the remaining events



$$\{N^L(t)\} \sim PP(\nu\lambda(t)), \quad \{N^T(t)\} \sim PP((1-\nu)\lambda(t)), \quad \{N^L(t)\} \perp \{N^T(t)\}$$

**Consequence.** If  $\{N(t)\}_{0 \leq t \leq 1} \sim PP(\lambda(t))$ , with  $\lambda(t)$  piecewise constant with change-points  $\tau = (\tau_k)$  and intensities  $\lambda = (\lambda_k)$ , then

- ▶  $\lambda^L(t)$  piecewise constant with change-points  $(\tau_k)$  and intensities  $(\nu\lambda_k)$ ,
- ▶  $\lambda^T(t)$  piecewise constant with change-points  $(\tau_k)$  and intensities  $((1-\nu)\lambda_k)$ ,
- ▶  $\{N^L(t)\} \perp \{N^T(t)\}$ .

## Cross validation

Sampling event times provides two independent Poisson processes with **same change-points**.

## Cross validation

Sampling event times provides two independent Poisson processes with **same change-points**.

**Cross-validation procedure.** For  $1 \leq K \leq K_{\max}$ ,

► Repeat for  $1 \leq m \leq M$  :

1 – Sample the event times to form  $\{N^{L,m}(t)\}$  (learn) and  $\{N^{T,m}(t)\}$  (test),

2 – Estimate  $\hat{\tau}^{L,m}$  and  $\hat{\lambda}^{L,m}$  from  $\{N^{L,m}(t)\}$ ,

3 – Compute the contrast  $\gamma_K^{T,m} = \gamma \left( \{N^T(t)\}; \hat{\tau}^{L,m}, \frac{1-v}{v} \hat{\lambda}^{L,m} \right)$ .

## Cross validation

Sampling event times provides two independent Poisson processes with **same change-points**.

**Cross-validation procedure.** For  $1 \leq K \leq K_{\max}$ ,

► Repeat for  $1 \leq m \leq M$  :

1 – Sample the event times to form  $\{N^{L,m}(t)\}$  (learn) and  $\{N^{T,m}(t)\}$  (test),

2 – Estimate  $\hat{\tau}^{L,m}$  and  $\hat{\lambda}^{L,m}$  from  $\{N^{L,m}(t)\}$ ,

3 – Compute the contrast  $\gamma_K^{T,m} = \gamma \left( \{N^{T,m}(t)\}; \hat{\tau}^{L,m}, \frac{1-v}{v} \hat{\lambda}^{L,m} \right)$ .

► Compute

$$\bar{\gamma}_K = \frac{1}{M} \sum_{m=1}^M \gamma_K^{T,m}$$

## Cross validation

Sampling event times provides two independent Poisson processes with **same change-points**.

**Cross-validation procedure.** For  $1 \leq K \leq K_{\max}$ ,

► Repeat for  $1 \leq m \leq M$  :

1 – Sample the event times to form  $\{N^{L,m}(t)\}$  (learn) and  $\{N^{T,m}(t)\}$  (test),

2 – Estimate  $\hat{\tau}^{L,m}$  and  $\hat{\lambda}^{L,m}$  from  $\{N^{L,m}(t)\}$ ,

3 – Compute the contrast  $\gamma_K^{T,m} = \gamma\left(\{N^T(t)\}; \hat{\tau}^{L,m}, \frac{1-v}{v} \hat{\lambda}^{L,m}\right)$ .

► Compute

$$\bar{\gamma}_K = \frac{1}{M} \sum_{m=1}^M \gamma_K^{T,m}$$

► Select

$$\hat{K} = \arg \min_K \bar{\gamma}_K$$

# Outline

Estimation

Model selection

**Illustrations**

Extensions

## Practical implementation.

**Contrasts.** During the CV process, we use

- ▶ a sampling rate of  $\nu = 4/5$ ,
- ▶ the Poisson-Gamma contrast  $\gamma_{PG}$  for the learning step and
- ▶ the Poisson contrast  $\gamma_P$  for the test step.



## Practical implementation.

**Contrasts.** During the CV process, we use

- ▶ a sampling rate of  $\nu = 4/5$ ,
- ▶ the Poisson-Gamma contrast  $\gamma_{PG}$  for the learning step and
- ▶ the Poisson contrast  $\gamma_P$  for the test step.

**Hyper-parameters.** For an observed path  $\{N(t)\}_{0 \leq t \leq 1}$  with  $n$  events, we use

$$a = 1, \quad b = 1/n$$

to fit the observed total number of events.

## Practical implementation.

**Contrasts.** During the CV process, we use

- ▶ a sampling rate of  $\nu = 4/5$ ,
- ▶ the Poisson-Gamma contrast  $\gamma_{PG}$  for the learning step and
- ▶ the Poisson contrast  $\gamma_P$  for the test step.

**Hyper-parameters.** For an observed path  $\{N(t)\}_{0 \leq t \leq 1}$  with  $n$  events, we use

$$a = 1, \quad b = 1/n$$

to fit the observed total number of events.

**R package `CptPointProcess`** available on [github.com/Elebarbier/CptPointProcess](https://github.com/Elebarbier/CptPointProcess).

## Some simulations

**Simulation setting.**  $K = 6$  segments with varying length. Tuning parameters:

- ▶  $\bar{\lambda}$  average intensity ( $\rightarrow$  total number of events),
- ▶  $\lambda_R$  = height of the steps ( $\rightarrow$  contrast between segments).


# Some simulations

**Simulation setting.**  $K = 6$  segments with varying length. Tuning parameters:


- ▶  $\bar{\lambda}$  average intensity ( $\rightarrow$  total number of events),
- ▶  $\lambda_R$  = height of the steps ( $\rightarrow$  contrast between segments).

**Results.** Choose  $K$  via CV, then refit the parameters to the whole dataset.


Model selection  
 $\hat{K}$



Change-point location  
Hausdorff( $\hat{\tau}, \tau^*$ )




Cumulated intensity  
 $\ell_2(\hat{\Lambda}, \Lambda^*)$




# Kilauea eruptions

$n = 63$  eruptions reported between the mid 18th and the late 20th century.

Model selection via CV



Resulting segmentation



# Outline

Estimation

Model selection

Illustrations

Extensions

# Extensions

## Marked Poisson Process.

- ▶  $\{Y(t)\}_{0 \leq t \leq 1} \sim MPP(\lambda(t), \mu(t))$ :

$$\{N(t)\}_{0 \leq t \leq 1} \sim PP(\lambda(t)), \quad \text{at each } T_i: X_i \sim \mathcal{F}(\mu(T_i))$$

- ▶ Works the same way, provided that concavity holds.
- ▶ Bat cries: Mark = bat species or cry duration.
- ▶ Poisson-Gamma events + Exponential-Gamma durations is both admissible and desirable.

## Extensions

### Marked Poisson Process.

- ▶  $\{Y(t)\}_{0 \leq t \leq 1} \sim MPP(\lambda(t), \mu(t))$ :

$$\{N(t)\}_{0 \leq t \leq 1} \sim PP(\lambda(t)), \quad \text{at each } T_i: X_i \sim \mathcal{F}(\mu(T_i))$$

- ▶ Works the same way, provided that concavity holds.
- ▶ Bat cries: Mark = bat species or cry duration.
- ▶ Poisson-Gamma events + Exponential-Gamma durations is both admissible and desirable.

### Segmentation-clustering.

- ▶ Each segment belongs to a class  $1 \leq q \leq Q$  (with probability  $\pi_q$  and intensity  $\lambda_k = \ell_q$ ),
- ▶ Combination of EM and DP algorithms [PRLD07],
- ▶ Bat cries: Class = animal behaviour (hunt, transit, ...)



## Extensions

### Marked Poisson Process.

- ▶  $\{Y(t)\}_{0 \leq t \leq 1} \sim MPP(\lambda(t), \mu(t))$ :

$$\{N(t)\}_{0 \leq t \leq 1} \sim PP(\lambda(t)), \quad \text{at each } T_i: X_i \sim \mathcal{F}(\mu(T_i))$$

- ▶ Works the same way, provided that concavity holds.
- ▶ Bat cries: Mark = bat species or cry duration.
- ▶ Poisson-Gamma events + Exponential-Gamma durations is both admissible and desirable.




### Segmentation-clustering.

- ▶ Each segment belongs to a class  $1 \leq q \leq Q$  (with probability  $\pi_q$  and intensity  $\lambda_k = \ell_q$ ),
- ▶ Combination of EM and DP algorithms [PRLD07],
- ▶ Bat cries: Class = animal behaviour (hunt, transit, ...)

### And also.

- ▶ Theoretically grounded model selection criterion (BIC),
- ▶ Consistency of the estimated change-points,
- ▶ Other desirable contrasts, ...

# References I

-  Lion-Blanc, E Lebarbier, and S Robin. Multiple change-point detection for poisson processes. Technical Report 2302.09103, arXiv, 2023.
-  Ho and M Bhaduri. A quantitative insight into the dependence dynamics of the Kilauea and Mauna Loa volcanoes, Hawaii. *Mathematical Geosciences*, 49(7):893–911, 2017.
-  F. Picard, S. Robin, E Lebarbier, and J-J Daudin. A segmentation/clustering model for the analysis of array CGH data. *Biometrics*, 63(3):758–766, 2007.

# Appendix

Number of elements in the partition of the segmentation space.


$$|\mathcal{N}_K| = \sum_{h=\lfloor (K-1)/2 \rfloor}^K \binom{n-1}{h-1} \binom{h+1}{K-h}$$

# Appendix






Number of elements in the partition of the segmentation space.

$$|\mathcal{N}_K| = \sum_{h=\lfloor (K-1)/2 \rfloor}^K \binom{n-1}{h-1} \binom{h+1}{K-h}$$

Simulations: Shape of the intensity function  $\lambda(t)$ .  $K = 6$ ,  $\bar{\lambda} = 100$ ,  $\lambda_R = 1, 3, 8$ .



# Poisson process: Mauna Loa eruptions

 $n = 40$ 
 $K = 2$ 

 $K = 3$ 

 $K = 4$ 

 $K = 5$ 

 $K = 6$ 


## Marked Poisson process: Etna eruptions

**Count and marks:** Events = eruptions, marks = duration of each eruption.

**Model.**

- ▶ Piecewise-constant intensity Poisson process for the events
- ▶ Exponential distribution (with segment specific parm.) for the durations


## Marked Poisson process: Etna eruptions

**Count and marks:** Events = eruptions, marks = duration of each eruption.

**Model.**


- Piecewise-constant intensity Poisson process for the events
- Exponential distribution (with segment specific parm.) for the durations

**CV for the selection of  $K$ .**




# Marked Poisson process: Etna eruptions

Poisson ( $\hat{K} = 2$ )  
Events




Marked Poisson ( $\hat{K} = 2$ )  
Events Marks






# Marked Poisson process: Etna eruptions

Poisson ( $\hat{K} = 2$ )  
Events



Marked Poisson ( $\hat{K} = 2$ )  
Events Marks



Marked Poisson ( $K = 3$ )  
Events Marks

