



Extrafloral nectaries exhibit dual ecological functions in a plant from the Brazilian Cerrado

Renan Fernandes Moura^{1,2} · Liégý Resende dos Santos³ · Vitor Miguel da Costa Silva³ · Roberth Fagundes⁴ · Kleber Del-Claro^{3,5}

Received: 12 October 2022 / Accepted: 30 March 2023 / Published online: 20 April 2023
© The Author(s), under exclusive licence to Springer Nature B.V. 2023

Abstract

Extrafloral nectaries (EFNs) are nectar-secreting plant glands not related to pollination. Several not mutually exclusive hypotheses explain the ecological function of EFNs. We focused on the (1) protection hypothesis, which predicts the attraction of natural enemies by the extrafloral nectar, providing indirect defense to plants against herbivores, and the (2) ant-distracting hypothesis, which predicts the distraction of ants away from sap-sucking herbivores secreting sugary rewards (honeydew). We aimed to test both hypotheses simultaneously to understand the relative ecological roles of EFNs in a model plant from the Brazilian Cerrado. We experimentally manipulated plant groups according to the presence or absence of EFNs and hemipterans during two years of field study (2018 and 2019). We found some support for both hypotheses. Extrafloral nectaries reduced the damage caused by herbivores by attracting predatory ants, although plants with both EFNs and hemipterans showed the highest herbivory reduction in 2018. The presence of neither EFNs nor hemipterans was associated with increased fruit production. The honeydew, which had higher sugar concentrations than the extrafloral nectar, was used by ants for longer periods. However, more than 70% of the ants were observed feeding on the extrafloral nectar instead of honeydew. We suggest that the relatively high quantity of EFNs can compensate for their lower quality and still reduce the number of ants attending and protecting hemipterans, ultimately mitigating plant damage. In summary, EFNs may fit multiple functions depending on the ecological context.

Keywords Brazilian savanna · Coevolution · Evolutionary ecology · Insect-plant plant defenses · Mutualism

Introduction

Extrafloral nectaries (EFNs) are plant glands that secrete nectar, a sugar-rich aqueous solution that also contains small amounts of amino acids, lipids, alkaloids, and volatile organic compounds (Bentley 1977; Koptur 1992; Blüthgen et al. 2004a). EFNs occur in more than 110 families of plants

and 876 genera (Weber et al. 2015). In the Cerrado biome (Brazilian savanna), 25% of woody species have EFNs, representing about 30% of all trees found in that biome (Oliveira and Freitas 2004). Despite the similar chemical composition to floral nectar, extrafloral nectar usually has a higher sugar concentration (Koptur 1994) and is not related to pollination (Del-Claro et al. 2016). Due to their nutritional value (Blüthgen et al. 2004a; Byk and Del-Claro 2011; Calixto et al. 2021a), EFNs attract several animals, such as ants, spiders, and wasps (Nahas et al. 2012; Stefani et al. 2015; Moura et al. 2021). Some animals, especially ants, often attack and remove arthropod herbivores from those plants (Fagundes et al. 2017), reducing herbivory levels (Oliveira and Pie 1998; Schoederer et al. 2010; Gomes et al. 2021; Moura et al. 2022; Moura and Del-Claro 2023) and increasing fruit and seed production (Rosumek et al. 2009; Nascimento and Del-Claro 2010).

Extrafloral nectaries are thus mainly considered to be part of a plant's indirect defense system (Pearse et al. 2020),

Communicated by Avery L Russell.

✉ Renan Fernandes Moura
renanfmoura@gmail.com

¹ Dead Sea and Arava Science Center, Masada, Israel

² Tel Aviv University, Tel Aviv-Yafo, Israel

³ Universidade de São Paulo, São Paulo, SP, Brazil

⁴ Universidade da Integração Internacional da Lusofonia Afro-Brasileira, Redenção, CE, Brazil

⁵ Universidade Federal de Uberlândia, Uberlândia, MG, Brazil

as predicted by the plant protection hypothesis (Bentley 1977; Calixto et al. 2018; Moura et al. 2021). Despite the enormous amount of evidence supporting this claim (e.g., Rosumek et al. 2009), other hypotheses propose alternative roles for the EFNs (Del-Claro et al. 2016), since ants do not always defend plants from herbivores. Such inconsistency is not surprising, since the outcomes of ant–plant interactions vary according to different ecological factors, including the involved plant (Trager et al. 2010; Baker-Méio and Marquis 2012), herbivore (Alves-Silva et al. 2015), and ant species (Ness et al. 2006; Del-Claro and Marquis 2015). For instance, many ant species occurring on EFN-bearing plants are simply unable to prey on or remove herbivores due to physical limitations (e.g., small size) or lack of aggressiveness (Fagundes et al. 2017). In addition, the quality and quantity of extrafloral nectar production strongly vary in time and space and among species (Lange et al. 2017; Calixto et al. 2021b), directly affecting ant community composition, visiting frequency, and, consequently, the effectiveness of ant protection (Pacelle et al. 2019; Moura and Del-Claro 2023). For example, well-protected plants usually offer high-quality rewards (Pacelle et al. 2019) and exhibit just a single or a few dominant ant species that are highly aggressive (Miller 2007).

A second hypothesis through which EFNs can be beneficial for plants is the ant-distracting hypothesis. This hypothesis was first proposed by Becerra and Venable (1989) and suggests that EFNs may have evolved in response to the negative effects of honeydew-producing hemipterans (trophobionts) on plants (Pierce et al. 2002). Some hemipterans offer sugary rewards (e.g., honeydew) to lure ants and use them as bodyguards against predators, parasitoids, and pathogens (Stadler and Dixon 2005; Nielsen et al. 2010). Thus, under the protection of tending ants, hemipteran populations may grow dramatically (Rice and Eubanks 2013), spreading plant diseases and inflicting high damage to leaves and reproductive structures (Rico-Gray and Castro 1996; Delabie 2001; Nelson and Mooney 2022). Some researchers have acknowledged that EFNs can be as rich or richer in nutrients than honeydew (Engel et al. 2001; Katayama et al. 2013; Chanam et al. 2015) and hypothesized that EFNs may serve as an alternative resource to disrupt ant–hemipteran mutualisms. This particular role of EFNs could be valuable for plants since developing chemical defenses against sap-sucking insects imposes many physiological constraints (Züst and Agrawal 2016). Furthermore, the peak production of EFNs is relatively predictable in space and time (Lange et al. 2017), and the extrafloral nectar is usually more abundant and easier to consume than honeydew (Becerra and Venable 1989). In comparison to the protection hypothesis, however, the ant-distracting hypothesis has been much less tested, and there is no consensus on the relative importance of the latter to ant–plant–hemipteran interactions (Del-Claro

and Oliveira 1993; Katayama and Suzuki 2003; Blüthgen et al. 2004a). The outcomes of ant–plant–hemipteran interactions are also highly contextual. For instance, ants may simply fail to protect hemipterans and eventually prey on them (Billick et al. 2007; Schifani et al. 2023). However, some studies further suggest that, in certain contexts, trophobiont hemipterans are ecologically analogous to EFNs and provide indirect plant defenses against non-tended herbivores via ant attraction (Moura et al. 2021).

Considering the high variability of studies evaluating the effects of EFNs on plants (Trager et al. 2010), testing alternative hypotheses for the role of EFNs may help explain why some published studies have not found mutualistic effects between ants and plants with EFNs. While many studies have investigated alternatives to the protection hypothesis (Del-Claro et al. 2016), we are not aware of any study that has experimentally investigated the effects of multiple hypotheses. Here, we investigated the ecological role of EFNs by experimentally testing both the protection and ant-distracting hypotheses in a system involving a native extrafloral-bearing species from the Brazilian Cerrado that commonly hosts honeydew-producing hemipterans.

Material and methods


Study area

This study took place at the Clube Caça e Pesca Itororó (CCPIU) reserve, west of Uberlândia city (18° 59' S, 48° 18' W), Minas Gerais. The site was chosen for having an extensive area of 628 ha, mainly characterized by typical Cerrado vegetation (Brazilian Savanna). This is the largest remnant of the natural Cerrado within the limits of Uberlândia, bearing trees and shrub species ranging between 2 and 8 m in height. The climate is characterized by a dry (May–September) and a wet (October–April) season, with an average temperature of 22.7 °C (Velasque and Del-Claro 2016).

Study species

We chose *Banisteriopsis malifolia* (Malpighiaceae) (Fig. 1a, b) as a model species. It is a shrub species (about 2 m tall) that bears two extrafloral nectaries (EFNs) on the base of each leaf (Fig. 1e). This species is abundant and easy to access at the study site, which allowed us to carry out the proposed experiments simultaneously. Trophobiont hemipterans are commonly found on the reproductive parts of this plant (Fig. 1c), and many ants feed on the honeydew produced by them (Fig. 1d) (Alves-Silva 2011; Alves-Silva et al. 2013). Other common herbivores are endophytic beetles that

Fig. 1 Pictures of *Banisteriopsis malifolia* (Malpighiaceae) (a), a common shrub with different ant resources (honeydew-producing hemipterans and extrafloral nectaries) (b). Trophobiont hemipterans on flowers (c), ants feeding on extrafloral nectar (d), and a pair of extrafloral nectaries (e)



consume flower buds (Torezan-Silingardi 2011; Mendes-Silva et al. 2021).

Sampling procedure

Most of the methods described below were carried out in two flowering periods of *Banisteriopsis malifolia*: between March and April 2018 and 2019, although we performed some additional experiments in 2021 and 2022 (see below). During these periods, we used different plant individuals to test each hypothesis. All marked individuals of *B. malifolia* were at least three meters apart, were between 1.5 and 2 m in height, and had leaves in the sprouting stage. After establishing all treatments, we performed manipulative experiments to test the ecological hypothesis of the EFNs. For logistical and time-optimization reasons, we selected two study sites 100 m apart, one for each hypothesis test. We studied 90 *B. malifolia* plants to test the protection and ant-distracting hypotheses (30 and 60 plants, respectively). Voucher specimens of all observed ant species were collected for posterior identification in the laboratory.

Protection hypothesis

Between 2018 and 2019, we randomly divided 30 plants into two groups ($n=15$): no ants (exclusion) and Control. The study site was visited three times a week, and each plant

was observed for 5 min. We manually removed all ants from plants in the Exclusion group and used a non-toxic sticky resin (Tanglefoot®), applied on masking tape positioned on the base of the plant stem, to restrict the access of ants to the EFNs. In the Control group, the resin was only partially applied to the stem so ants could freely pass. In both groups, we marked three branches on each plant, and in each branch, we counted the number of ants and herbivores. In each branch, young leaves were tagged (six per branch) to estimate leaf area loss after two months of observation. Leaves were photographed on the tagging day and after the two-month period. Leaf area loss was estimated with ImageJ software (Schneider et al. 2012). The percentage of leaf area loss was calculated by dividing the total leaf area by the estimated area loss.

Ant-distracting hypothesis

In 2018 and 2019, we studied 60 plants divided into four groups: Exclusion, Hemipteran (*Enchenopa* sp., Membracidae), EFN, and EFN + Hemipteran ($n=15$ for each group). For each plant, we randomly selected three branches to work with. The Exclusion group had all hemipterans manually removed from the selected branches, while all EFNs were obstructed with the aid of colorless nail polishes. The EFN group had all hemipterans removed, but the EFNs were preserved. The Hemipteran group had all hemipterans

preserved, while all EFNs were obstructed. Finally, the EFN + Hemipteran group had all components preserved and was characterized as our control. In this group, we applied colorless nail polish just beside the EFNs. After establishing all plant groups, we performed one-minute observations on each plant. During the observations, we counted the number of ants on the EFNs and the number of hemipterans found on the sampled branches. For all sampled branches, fruit production was evaluated by the end of the experiment, and the percentage of leaf herbivory was calculated according to the methods used to test hypothesis 1 (above).

In 2021, we repeated the same treatments: EFN, Hemipteran, and EFN + Hemipteran ($n = 10$ for each group) to assess the time ants spent using each resource (honeydew or extrafloral nectar) and moving around the plant. For this, we randomly selected one ant from each plant and observed it for 3 min. During this observation period, we recorded how much time the ants spent tending trophobiont hemipterans, feeding on the EFNs, and moving around the plant.

Sugar concentrations in honeydew and extrafloral nectar

In 2022, we replicated treatments EFN, Hemipteran, and EFN + Hemipteran ($n = 10$ for each group) to measure the sugar concentrations of the honeydew and extrafloral nectar. Branches with Hemipterans and/or EFNs were bagged with polyester fabrics between 7:00 and 8:00 am. Polyester bags minimize water evaporation from nectar and ease the sampling process of extrafloral nectar and honeydew without interference from ants and other arthropods. After 24 h, we returned to the field to collect the nectar and honeydew. Hemipterans were held laterally with flexible forceps and lightly pressed for honeydew secretion, which was collected with a graduated microcapillary tube (5 μ L). The same types of tubes were used to collect nectar from EFNs. Finally, we used a manual refractometer (Eclipse model) to measure the sugar concentrations of the honeydew and the extrafloral nectar of each plant (see Lange et al. 2017, for details).

Data analyses

All tests were performed using R statistical software (R Core Team 2020, version 4.0.0). We used analysis of variance (ANOVA) and linear mixed model (LMM) procedures to test the two hypotheses in two years of study (2018 and 2019). To test the protection hypothesis, we conducted two one-way ANOVAs between the Control (no manipulation) and Exclusion (no ants) groups using leaf herbivory proportion and the production of fruits as response variables. To evaluate the ant-distracting hypothesis, we first conducted a set of two-way ANOVAs testing for the effects of both the presence of EFNs and hemipterans (EFN, Hemipteran,

Exclusion, and Control groups) on leaf herbivory proportion and fruit production. We also evaluated interaction effects (EFNs \times hemipteran factors) and performed Tukey post-hoc tests when necessary. Afterward, we performed two LMMs evaluating the mean proportion of ants observed attending either EFNs or hemipterans on plants from three groups: EFN + Hemipteran (Control), EFN, and Hemipteran. We used plant identity and the number of hemipterans as random factors: $y \sim x + (1|\text{plant ID}) + (1|\text{no hemipterans})$ (Bates et al. 2015). The goal here was to verify whether ants prefer a specific resource when both occur in the same plant (EFN + Hemipteran group) and how ants use those resources when they occur apart (EFN and Hemipteran groups), while controlling for potential effects of the number of hemipterans. Using the same plant groups, we performed another LMM to test for differences in the time ants spent using each resource type (honeydew or extrafloral nectar) and moving on the plant. Finally, we used a t-test to analyze the possible differences between the sugar concentrations of honeydew and extrafloral nectar.


We performed a set of analyses of similarity (ANOSIM; Clarke 1993) to test for differences in the species composition of ants among plant groups (Control, Hemipteran, and EFN) from each year of the study. The ANOSIM set was performed using the “vegan” package of R (Oksanen et al. 2013). We first used the Hellinger transformation on our data (square root of the sample total standardized data). Then, the Bray–Curtis distance was calculated, and 999 permutations were performed based on the number of visiting ants per plant (Santos and Moura 2021). From the total number of individuals ($n = 45$), we excluded plants that were not visited by any ant (2018: $n = 6$; 2019: $n = 3$). After ANOSIM, we generated a set of nonmetric multidimensional scaling (NMDS) graphs to visualize the results.

All statistical assumptions were verified according to Zuur et al. (2010). Data normality was assessed using box-plots, histograms, and Lilliefors normality tests. Variance homogeneity was verified using the function “var. test” from the “stats” package and the function “simulateResiduals” from the “DHARMA” package. Variability signs and values in the Results section depict the standard error of the mean (SEM).

Results

We observed a total of 10 morphospecies of ants visiting *B. malifolia*. The most common ants were *Camponotus* species and *Ectatomma tuberculatum*.

Fig. 2 Percentage of herbivory (a and b) and fruit production (c and d) in groups of *Banisteriopsis malifolia* (Malpighiaceae) where ants were maintained (Control; n = 15) or excluded (No ant; n = 15) in 2018 and 2019. Distinct letters indicate statistically different groups according to $\alpha = 5\%$



Protection hypothesis

In 2018 and 2019, leaf herbivory was greater in plants without ants (Exclusion group) (2018: $F_{1,28} = 27.78$; $p = 0.001$; 2019: $F_{1,28} = 13.97$; $p < 0.001$; Fig. 2a and b). However, fruit production did not differ between the groups (2018: $F_{1,28} = 0.031$; $p = 0.86$; 2019: $F_{1,28} = 0.30$; $p = 0.58$; Fig. 2c and d).

Ant-distracting hypothesis


In 2018, the highest herbivory was found in plants without EFNs and hemipterans (Exclusion), while the lowest herbivory was observed in plants bearing both EFNs and hemipterans (2018: $F_{2,57} = 14.40$; $p < 0.001$; Fig. 3a). In 2019, the EFN + Hemipteran and EFN groups had herbivory values close to zero (2019: $F_{2,57} = 56.24$; $p < 0.001$; Fig. 3b). In both years, we did not observe significant differences in fruit production (2018: $F_{2,57} = 0.56$; $p = 0.57$; Fig. 3c; 2019: $F_{2,57} = 2.68$; $p = 0.077$; Fig. 3d).

Ants showed a preference for EFNs compared to hemipterans. In both years, about 70% of ants, on average, were observed feeding on EFNs when both EFNs

and hemipterans occurred on the same plant (Control), while less than 30% of them were observed feeding on hemipterans (2018: $F_{1,28} = 17.38$; $p < 0.001$; Fig. 4a; 2019: $F_{1,28} = 16.08$; $p < 0.001$; Fig. 4c). Interestingly, ants showed no preference when resources occurred separately on plants (only hemipterans or only EFNs) (2018: $F_{1,28} = 0.66$; $p = 0.43$; Fig. 4b; 2019: $F_{1,28} = 1.69$; $p = 0.20$; Fig. 4d).

When we compared ant activity between plant groups (Control, Hemipteran, and EFN) and the time spent using each resource type (honeydew or nectar) or moving on the plant, we found that ant activity differed according to the plant group and food resource ($F_{2,129} = 4.05$; $p = 0.019$; Fig. 5). In the Control and Hemipteran groups, the ants spent most of their time feeding on honeydew (Control: Honeydew = $74.5\% \pm 4.32$ SEM, Control: Nectar = $6\% \pm 1.66$ SEM; Fig. 5a; Hemipteran: Honeydew = $82\% \pm 2.26$ SEM; Fig. 5b). In the EFN group, the ants spent almost the same amount of time foraging or feeding on extrafloral nectar (Nectar = $52\% \pm 4.3$ SEM; Fig. 5c).

Fig. 3 Mean percentage of leaf herbivory (a and b) and fruit production (c and d) observed in four plant groups of *Banisteriopsis malifolia* (Malpigiaceae) during the two years of the study (n per group = 15). Hemip = only honeydew-producing hemipterans; EFN + Hemip = extrafloral nectaries and hemipterans on the same plant; Exclusion = both hemipterans and extrafloral nectaries excluded; EFN = only extrafloral nectaries. Distinct letters indicate statistically different groups according to $\alpha = 5\%$



Sugar concentrations in honeydew and extrafloral nectar

We observed that honeydew had a higher sugar concentration, on average, than extrafloral nectar (Honeydew: $24.25\% \pm 1.61$ SEM, $n = 20$; EFNs: $18.71\% \pm 0.65$ SEM, $n = 20$; $t_{2,38} = 3.19$, $p = 0.003$).

Ant species composition

We found no differences in ant species composition among plant groups (Control, Hemipteran, and EFN) either in 2018 (ANOSIM: $R = 0.022$, $p = 0.27$; Fig. 6a) or 2019 (ANOSIM: $R = -0.064$, $p = 0.99$; Fig. 6b).


Discussion

Our results provide partial support for both the protection and ant-distracting hypotheses, indicating that EFNs may possess multiple ecological roles. The protection hypothesis was supported, as we observed increased herbivory values in the absence of ants (Fig. 2a, and c), although the influence of ants was not converted into a higher reproductive output for *Banisteriopsis malifolia* (Fig. 2c and d). In the first year of

the study, we also observed that the simultaneous presence of hemipterans and EFNs was associated with the lowest values of foliar herbivory (Fig. 3a), although differences in the reproductive output were not detected (Fig. 3b and d). We found some support for the ant-distracting hypothesis since more than 70% of ants were registered feeding on the extrafloral nectar instead of honeydew when both resources were offered together (Fig. 4a and c). When offered separately (i.e., different plant groups), the resources were equally used by ants (Fig. 4b and d). However, we also observed that the ants fed for longer periods of time on honeydew, which contained, on average, 29.6% more sugar than the extrafloral nectar (Fig. 5).

The complex interactions involving plants, hemipterans, and ants partially explain why our study supported, at least partially, both hypotheses. EFNs may have evolved multiple ecological roles in regions where plants with EFNs co-occur with sap-sucking insects (Chanam et al. 2015). In addition to the protection role, EFNs can distract ants that could potentially be attending hemipterans. While hemipterans have also been associated with a reduction in herbivory, we speculate that EFNs may help control hemipteran populations, thus minimizing plant damage due to overpopulation, as large numbers of sap-sucking hemipterans are associated with diseases and increased plant damage (Delabie 2001;

Fig. 4 Proportion of visiting ants using resources (honeydew or extrafloral nectar) of *Banisteriopsis malifolia* (Malpighiaceae) exhibiting the simultaneous presence of extrafloral nectaries and hemipterans (a and c; EFN + Hemipteran group; $n = 15$) and in plants where only one type of resource was offered (b and d; Hemipteran or EFN groups; n per group = 15). Distinct letters indicate statistically different groups according to $\alpha = 5\%$



Moura et al. 2021). Furthermore, EFNs might reduce the number of ants visiting flowers, thus minimizing potential conflicts between ants and pollinators. Otherwise, hemipterans are commonly found on flowers; thus, attracted ants may interfere with pollinator services (Ibarra-Isassi and Oliveira 2018).

EFNs and hemipterans exhibited additive protective effects on plants in the first year of our study. In 2018, we observed that the lowest herbivory values were associated with plants producing both extrafloral nectar and honeydew, although we did not observe any differences in fruit production. This similar productivity might be explained by the fact that *B. malifolia* is constantly attacked by endophytic beetles that consume flower buds and bypass ant defenses (Torezan-Silingardi 2011; Mendes-Silva et al. 2021). Nevertheless, it is important to notice that we did not measure seed production, so we do not know whether herbivore damage could have caused changes in the number of viable seeds, for instance. Although trophobiont hemipterans can eventually provide benefits to plants (see Styrsky and Eubanks 2010; Vilela and Del-Claro 2018), their positive effects do

not always stack when combined with other resources, such as EFNs (Savage and Rudgers 2013). The effects of combining hemipterans and EFNs are thus largely dependent on the ecological context, such as resource availability, the involved ant species, and herbivore pressure (Davidson et al. 2003; Rudgers et al. 2010; Chanam et al. 2015; Lange et al. 2017). Such conditional scenarios may explain why plants with only EFNs already showed virtually no herbivore damage in the second year of study. Furthermore, as we observed that the ant community composition did not change depending on the offered resource, eventual differences among plant treatments in terms of resource use and protection can be mostly attributed to unequal ant recruitment.

The greater abundance of ants feeding on extrafloral nectar than honeydew was somewhat surprising, given the higher sugar concentration in the honeydew, although the available evidence in the literature is conflicting. While some studies have shown that honeydew tends to attract more ants, as they are usually richer in nutrients than EFNs (Blüthgen et al. 2004a)—they may even contain addictive chemicals (Kudo et al. 2021)—other studies have demonstrated that




Fig. 5 Time spent by ants moving around the plant and feeding on honeydew or extrafloral nectar from the three groups of *Banisteriopsis malifolia* (Malpighiaceae): Control (no manipulation; $n = 10$), Hemipteran (only honeydew-producing hemipterans; $n = 10$), and EFN (only extrafloral nectaries; $n = 10$) (a–c, respectively). Distinct letters indicate statistically different groups according to $\alpha = 5\%$

ants are less attracted by the complex sugars found in honeydew (Blüthgen and Fiedler 2004a). Although 70% of the total ants were observed feeding on EFNs instead of hemipterans when both resources were available, we showed that the ants spent more time tending hemipterans than feeding on EFNs. We believe that the relative abundance of EFNs and hemipterans on *B. malifolia* might explain this apparent paradox. While high-quality resources can trigger intense competition among ants, having enough power to structure whole ant communities (Blüthgen et al. 2004b; Orivel et al. 2018), they are not the sole meaningful factor influencing




Fig. 6 Species composition of ants visiting three plant groups of *Banisteriopsis malifolia* (Malpighiaceae) in 2018 (a) and 2019 (b). Control=no manipulation ($n_{2018} = 14$; $n_{2019} = 14$); Hemipteran=only honeydew-producing hemipterans ($n_{2018} = 11$; $n_{2019} = 14$); EFN=only extrafloral nectaries ($n_{2018} = 14$; $n_{2019} = 14$). Ant species composition was not different among the groups ($\alpha = 5\%$) either in 2018 (ANOSIM: $R = 0.022$, $p = 0.27$) or 2019 (ANOSIM: $R = -0.064$, $p = 0.99$). Red triangles represent the ant species distribution

foraging decisions among ants. Ants make foraging decisions based on their preferences and the availability and distribution of resources (Davidson et al. 2003; Blüthgen and Fiedler 2004b; Savage and Rudgers 2013). Even though EFNs produced nectar with lower sugar concentrations than honeydew, they were more abundant on the plant (two EFN glands per leaf) and produced a larger reward volume (personal observation). Thus, ants might be willing to use the first encountered food resource, regardless of the quality, which by a simple matter of probability will most likely be the extrafloral nectar. The apparent collective preference for EFNs might reflect their high availability on the plant. This also explains why ants used extrafloral nectar and honeydew equally when no alternatives were offered. This reasoning

is strengthened by the fact that plants with either EFNs or hemipterans were visited by similar ant communities, suggesting that the observed feeding preferences cannot be attributed to interspecific differences.

From an evolutionary perspective, the quantitative investment of *B. malifolia* in EFNs might be beneficial in two ways. First, abundant EFNs may still distract ants from hemipterans—and minimize their potential damage to plants—despite the putative inferior nutritious value of extrafloral nectar compared to honeydew. We hypothesize that resources of medium quality may foster ant protection since ants have to move around the plant in search of additional resources. Otherwise, resources of high quality could reduce ant movement (see Fig. 5b and c) and consequently the defensive potential of ants. Second, having more EFNs scattered on plants means that more ants can simultaneously visit the plant, which may improve indirect defenses. We also argue that a strong preference for hemipterans over EFNs would most likely be an unstable evolutionary strategy—for both ants and plants—since ants can stimulate the overgrowth of hemipteran populations, rapidly reducing plant fitness and ultimately ant fitness. Still, our results suggest that keeping hemipteran populations under control by reducing hemipteran attendance to stable levels via ant-distracting mechanisms may actually improve plant defenses. In summary, *B. malifolia* investment in EFNs is in line with both the protection and ant-distracting hypotheses.

We concluded that EFNs may suit more than a single ecological role in plants (e.g., indirect defense), strengthening the importance of these structures for plant populations and possibly plant communities. We also concluded that honeydew-producing hemipterans are not intrinsically negative for plants, and under certain circumstances, they may even foster indirect defenses provided by ants. As for future perspectives, the next step is to understand how the relative quantity, productivity, and positioning of both EFNs and honeydew-producing hemipterans affect ant–plant–hemipteran interactions.

Acknowledgements We thank the Laboratório de Ecologia Comportamental e de Interações (LECI), two anonymous reviewers, and Dr. Avery Russell for interesting discussions and suggestions that improved the quality of this manuscript. LRS thanks the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for funding – Finance Code 001.

Declarations

Competing interests The authors declare no competing interests.

References

- Alves-Silva E (2011) Post fire resprouting of *Banisteriopsis malifolia* (Malpighiaceae) and the role of extrafloral nectaries on the associated ant fauna in a Brazilian Savanna. *Sociobiology* 58:327
- Alves-Silva E, Bächtold A, Barônio GJ et al (2015) Ant–herbivore interactions in an extrafloral nectaried plant: are ants good plant guards against curculionid beetles? *J Nat Hist* 49:841–851
- Alves-Silva E, Barônio GJ, Torezan-Silingardi HM, Del-Claro K (2013) Foraging behavior of *Brachygastra lecheguana* (Hymenoptera: Vespidae) on *Banisteriopsis malifolia* (Malpighiaceae): extrafloral nectar consumption and herbivore predation in a tending ant system. *Entomol Sci* 16:162–169
- Baker-Méio B, Marquis RJ (2012) Context-dependent benefits from ant–plant mutualism in three sympatric varieties of *Chamaecrista desvauxii*. *J Ecol* 100:242–252
- Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4. *J Stat Softw* 67(1):1–48
- Becerra JXI, Venable DL (1989) Extrafloral nectaries: a defense against ant–Homoptera mutualisms? *Oikos* 55(2):276–280
- Bentley BL (1977) The protective function of ants visiting the extrafloral nectaries of *Bixa orellana* (Bixaceae). *J Ecol* 65(1):27–38
- Billick I, Hammer S, Reithel JS, Abbot P (2007) Ant–aphid interactions: are ants friends enemies or both? *Ann Entomol Soc Am* 100(6):887–892
- Blüthgen N, Fiedler K (2004a) Preferences for sugars and amino acids and their conditionality in a diverse nectar-feeding ant community. *J Anim Ecol* 73:155–166
- Blüthgen N, Fiedler K (2004b) Competition for composition: lessons from nectar-feeding ant communities. *Ecology* 85:1479–1485
- Blüthgen N, Gottsberger G, Fiedler K (2004a) Sugar and amino acid composition of ant-attended nectar and honeydew sources from an Australian rainforest. *Austral Ecol* 29:418–429
- Blüthgen N, Stork NE, Fiedler K (2004b) Bottom-up control and co-occurrence in complex communities: honeydew and nectar determine a rainforest ant mosaic. *Oikos* 106:344–358
- Byk J, Del-Claro K (2011) Ant–plant interaction in the Neotropical savanna: direct beneficial effects of extrafloral nectar on ant colony fitness. *Popul Ecol* 53:327–332
- Calixto ES, Lange D, Del-Claro K (2018) Protection mutualism: an overview of ant–plant interactions mediated by extrafloral nectaries. *Oecol Aust* 22:410–425
- Calixto ES, Lange D, Del-Claro K (2021a) Net benefits of a mutualism: influence of the quality of extrafloral nectar on the colony fitness of a mutualistic ant. *Biotropica* 53:846–856
- Calixto ES, Lange D, Moreira X, Del-Claro K (2021b) Plant species specificity of ant–lant mutualistic interactions: differential predation of termites by *Camponotus crassus* on five species of extrafloral nectaries plants. *Biotropica* 53(5):1406–1414
- Chanam J, Kasinathan S, Pramanik GK et al (2015) Foliar extrafloral nectar of *Humboldtia brunonis* (Fabaceae), a paleotropical ant-plant, is richer than phloem sap and more attractive than honeydew. *Biotropica* 47:1–5
- Clarke KR (1993) Non-parametric multivariate analyses of changes in community structure. *Aust J Ecol* 18:117–143
- Davidson DW, Cook SC, Snelling RR, Chua TH (2003) Explaining the abundance of ants in lowland tropical rainforest canopies. *Science* 300:969–972
- Del-Claro K, Marquis RJ (2015) Ant species identity has a greater effect than fire on the outcome of an ant protection system in Brazilian Cerrado. *Biotropica* 47:459–467
- Del-Claro K, Oliveira PS (1993) Ant–Homoptera interaction: do alternative sugar sources distract tending ants? *Oikos* 68(2):202–206

- Del-Claro K, Rico-Gray V, Torezan-Silingardi HM et al (2016) Loss and gains in ant–plant interactions mediated by extrafloral nectar: fidelity, cheats, and lies. *Insects Soc* 63:207–221
- Delabie JHC (2001) Trophobiosis between Formicidae and Hemiptera (Sternorrhyncha and Auchenorrhyncha): an overview. *Neotrop Entomol* 30:501–516
- Engel V, Fischer MK, Wäckers FL, Völkl W (2001) Interactions between extrafloral nectaries, aphids and ants: are there competition effects between plant and homopteran sugar sources? *Oecologia* 129:577–584
- Fagundes R, Dáttilo W, Ribeiro SP et al (2017) Differences among ant species in plant protection are related to production of extrafloral nectar and degree of leaf herbivory. *Biol J Linn Soc* 122:71–83
- Gomes IJMT, Solar R, Blüthgen N et al (2021) Multiple effects of mutualistic ants improve the performance of a Neotropical ant-plant: a long-term study with the *Cecropia-Azteca* system. *Basic Appl Ecol* 57:78–90
- Ibarra-Isassi J, Oliveira PS (2018) Indirect effects of mutualism: ant–treehopper associations deter pollinators and reduce reproduction in a tropical shrub. *Oecologia* 186(3):691–701
- Katayama N, Hembry DH, Hojo MK, Suzuki N (2013) Why do ants shift their foraging from extrafloral nectar to aphid honeydew? *Ecol Res* 28:919–926
- Katayama N, Suzuki N (2003) Bodyguard effects for aphids of *Aphis craccivora* Koch (Homoptera: Aphididae) as related to the activity of two ant species, *Tetramorium caespitum* Linnaeus (Hymenoptera: Formicidae) and *Lasius niger* L. (Hymenoptera: Formicidae). *Appl Entomol Zool* 38:427–433
- Koptur S (1992) Plants with extrafloral nectaries and ants in Everglades habitats. *Florida Entomol* 75(1):38–50
- Koptur S (1994) Floral and extrafloral nectars of Costa Rican Inga trees: a comparison of their constituents and composition. *Biotropica* 26(3):276–284
- Kudo T, Aonuma H, Hasegawa E (2021) A symbiotic aphid selfishly manipulates attending ants via dopamine in honeydew. *Sci Rep* 11:1–8
- Lange D, Calixto ES, Del-Claro K (2017) Variation in extrafloral nectary productivity influences the ant foraging. *PLoS ONE* 12:e0169492
- Mendes-Silva I, Queiroga D, Calixto ES et al (2021) Multiple cues guarantee successful predation by a Neotropical wasp. *Behaviour* 1:1–13
- Miller TEX (2007) Does having multiple partners weaken the benefits of facultative mutualism? A test with cacti and cactus-tending ants. *Oikos* 116:500–512
- Moreira VSS, Del-Claro K (2005) The outcomes of an ant–treehopper association on *Solanum lycocarpum* St. Hill: increased membracid fecundity and reduced damage by chewing herbivores. *Neotrop Entomol* 34:881–887
- Moura RF, Colberg E, Alves-Silva E, Mendes-Silva I, Fagundes R, Stefani V, Del-Claro K (2021) Biotic defenses against herbivory. In: Del-Claro K, Torezan-Silingardi HM (eds) Plant–animal interactions: source of biodiversity. Springer, Cham, pp 93–118
- Moura RF, Couto CMV, Del-Claro K (2022) Ant nest distribution and richness have opposite effects on a Neotropical plant with extrafloral nectaries. *Ecol Entomol* 47:626–635
- Moura RF, Del-Claro K (2023) Plants with extrafloral nectaries share indirect defenses and shape the local arboreal ant community. *Oecologia* 201:73–82
- Nahas L, Gonzaga MO, Del-Claro K (2012) Emergent impacts of ant and spider interactions: herbivory reduction in a tropical savanna tree. *Biotropica* 44:498–505
- Nascimento EA, Del-Claro K (2010) Ant visitation to extrafloral nectaries decreases herbivory and increases fruit set in *Chamaecrista debilis* (Fabaceae) in a Neotropical savanna. *Flora-morphology. Distrib Funct Ecol Plants* 205:754–756
- Nelson AS, Mooney KA (2022) The evolution and ecology of interactions between ants and honeydew-producing hemipteran insects. *Annu Rev Ecol Evol Syst* 53(1):379–402
- Ness JH, Morris WF, Bronstein JL (2006) Integrating quality and quantity of mutualistic service to contrast ant species protecting *Ferocactus wislizeni*. *Ecology* 87:912–921
- Nielsen C, Agrawal AA, Hajek AE (2010) Ants defend aphids against lethal disease. *Biol Lett* 6(2):205–208
- Oksanen J, Blanchet FG, Kindt R, et al (2013) Package ‘vegan.’ *Community Ecol Packag version* 2:1–295
- Oliveira PS, Freitas AVL (2004) Ant–plant–herbivore interactions in the neotropical cerrado savanna. *Naturwissenschaften* 91:557–570
- Oliveira PS, Pie MR (1998) Interaction between ants and plants bearing extrafloral nectaries in cerrado vegetation. *An Soc Entomol Bras* 27:161–176
- Orivel J, Klimes P, Novotny V, Leponce M (2018) Resource use and food preferences in understory ant communities along a complete elevational gradient in Papua New Guinea. *Biotropica* 50:641–648
- Pacelle FT, Costa FV, Neves FS et al (2019) Nectar quality affects ant aggressiveness and biotic defense provided to plants. *Biotropica* 51:196–204
- Pearse IS, LoPresti E, Schaeffer RN et al (2020) Generalising indirect defence and resistance of plants. *Ecol Lett* 23:1137–1152
- Pierce NE, Braby MF, Heath A et al (2002) The ecology and evolution of ant association in the Lycaenidae (Lepidoptera). *Annu Rev Entomol* 47:733–771
- R Core Team (2020) R: a language and environment for statistical computing. R Foundation for Statistical Computing. <http://www.r-project.org/>
- Rico-Gray V, Castro G (1996) Effect of an ant–aphid interaction on the reproductive fitness of *Paullinia fuscescens* (Sapindaceae). *Southwest Nat* 41(4):434–440
- Rice KB, Eubanks MD (2013) No enemies needed: cotton aphids (Hemiptera: Aphididae) directly benefit from red imported fire ant (Hymenoptera: Formicidae) tending. *Fla Entomol* 96(3):929–932
- Rosumek FB, Silveira FAO, de Neves FS et al (2009) Ants on plants: a meta-analysis of the role of ants as plant biotic defenses. *Oecologia* 160:537–549
- Rudgers JA, Savage AM, Rúa MA (2010) Geographic variation in a facultative mutualism: consequences for local arthropod composition and diversity. *Oecologia* 163:985–996
- Santos LR, Moura RF (2021) Local and temporal effects on ant richness, recruitment and resource dominance in two Cerrado areas. *Sociobiology* 68:e6204–e6204
- Savage AM, Rudgers JA (2013) Non-additive benefit or cost? Disentangling the indirect effects that occur when plants bearing extrafloral nectaries and honeydew-producing insects share exotic ant mutualists. *Ann Bot* 111:1295–1307
- Schifani E, Peri E, Giannetti D, Colazza S, Grasso DA (2023) Ant attendance does not necessarily imply protection of aphids from their arthropod natural enemies. *Ecol Entomol*. <https://doi.org/10.1111/een.13226>
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9:671–675
- Schoereder JH, Sobrinho TG, Madureira MS et al (2010) The arboreal ant community visiting extrafloral nectaries in the Neotropical cerrado savanna. *Terr Arthropod Rev* 3:3–27
- Stadler B, Dixon AFG (2005) Ecology and evolution of aphid–ant interactions. *Annu Rev Ecol Evol Syst* 36(1):345–372
- Stefani V, Pires TL, Torezan-Silingardi HM, Del-Claro K (2015) Beneficial effects of ants and spiders on the reproductive value of *Eriothea gracilipes* (Malvaceae) in a tropical savanna. *PLoS ONE* 10:e0131843

- Styrsky JD, Eubanks MD (2010) A facultative mutualism between aphids and an invasive ant increases plant reproduction. *Ecol Entomol* 35:190–199
- Torezan-Silingardi HM (2011) Predatory behavior of *Pachodynerus brevithorax* (Hymenoptera: Vespidae, Eumeninae) on endophytic herbivore beetles in the Brazilian tropical savanna. *Sociobiology* 57:181
- Trager MD, Bhotika S, Hostetler JA et al (2010) Benefits for plants in ant-plant protective mutualisms: a meta-analysis. *PLoS ONE* 5:e14308
- Velasque M, Del-Claro K (2016) Host plant phenology may determine the abundance of an ecosystem engineering herbivore in a tropical savanna. *Ecol Entomol* 41(4):421–430
- Vilela AA, Del-Claro K (2018) Effects of different ant species on the attendance of neighbouring hemipteran colonies and the outcomes for the host plant. *J Nat Hist* 52:415–428
- Weber MG, Porturas LD, Keeler KH (2015) World list of plants with extrafloral nectaries. <http://www.extrafloralnectaries.org/>
- Züst T, Agrawal AA (2016) Mechanisms and evolution of plant resistance to aphids. *Nat Plants* 2(1):1–9
- Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common statistical problems. *Methods Ecol Evol* 1:3–14
- Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.
- Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.