

Triangles in the Plane

Felix Christian Clemen (IBS ECOPRO)

Graph Theory Seminar at Shanghai Center for Mathematical Science

This is partially joint work with József Balogh and Adrian Dumitrescu.

October 22th, 2024




Hypergraph Turán Theory


$\text{ex}(n, H)$ = The maximum number of edges in an n -vertex k -graph G which does not contain H as a copy.

The Turán density of H :
$$\pi(H) = \lim_{n \rightarrow \infty} \frac{\text{ex}(n, H)}{\binom{n}{k}}$$

H_1



K_4^3




Hypergraph Turán Theory

$\text{ex}(n, H)$ = The maximum number of edges in an n -vertex k -graph G which does not contain H as a copy.


The Turán density of H :
$$\pi(H) = \lim_{n \rightarrow \infty} \frac{\text{ex}(n, H)}{\binom{n}{k}}$$

H_1



$$\begin{aligned}\text{ex}(n, H_1) &= 0 \\ \pi(H_1) &= 0\end{aligned}$$


K_4^3




Hypergraph Turán Theory

$\text{ex}(n, H)$ = The maximum number of edges in an n -vertex k -graph G which does not contain H as a copy.

The Turán density of H :
$$\pi(H) = \lim_{n \rightarrow \infty} \frac{\text{ex}(n, H)}{\binom{n}{k}}$$



$$\begin{aligned}\text{ex}(n, H_1) &= 0 \\ \pi(H_1) &= 0\end{aligned}$$



$$\begin{aligned}\text{ex}(n, K_4^3) &=? \\ \frac{5}{9} \leq \pi(K_4^3) &\leq 0.5615\end{aligned}$$

Turán's Tetrahedron Conjecture (1961): $\pi(K_4^3) = \frac{5}{9}$ (500\$)

Hypergraph Turán Theory

Some questions one might ask:

- Given a k -graph H , determine $\pi(H)$.
- What can be said about $\{\pi(H) : H \text{ is finite } k\text{-graph}\}$?

Hypergraph Turán Theory

Some questions one might ask:

- Given a k -graph H , determine $\pi(H)$.
- What can be said about $\{\pi(H) : H \text{ is finite } k\text{-graph}\}$?

Let \mathcal{H} be a family of k -graphs.


$\text{ex}(n, \mathcal{H})$ = The maximum number of edges in an n -vertex k -graph G which does not contain any $H \in \mathcal{H}$ as a copy.

The Turán density of \mathcal{H} :

$$\pi(\mathcal{H}) = \lim_{n \rightarrow \infty} \frac{\text{ex}(n, \mathcal{H})}{\binom{n}{k}}$$


Application 1: The unit distance problem

n points in \mathbb{R}^2



Application 1: The unit distance problem

n points in \mathbb{R}^2



Question (Erdős, 1946)

What is the maximum number of times that the unit distance can occur among n points in the plane?

$$u(n) := \max_{P \subseteq \mathbb{R}^2, |P|=n} \left| \{ \{u, v\} \subset P : |u - v| = 1 \} \right|.$$

Application 1: The unit distance problem

- Erdős (1946): $n^{1+c_1/\log \log n} \leq u(n) \leq O(n^{3/2})$
- Józsa and Szemerédi (1975): $u(n) = o(n^{3/2})$
- Beck and Spencer (1984): $u(n) = O(n^{13/9})$
- Spencer, Szemerédi, Trotter (1984): $u(n) = O(n^{4/3})$

Application 1: The unit distance problem

- Erdős (1946): $n^{1+c_1/\log \log n} \leq u(n) \leq O(n^{3/2})$
- Józsa and Szemerédi (1975): $u(n) = o(n^{3/2})$
- Beck and Spencer (1984): $u(n) = O(n^{13/9})$
- Spencer, Szemerédi, Trotter (1984): $u(n) = O(n^{4/3})$

500\$ for $u(n) = O(n^{1+\varepsilon})$ for every $\varepsilon > 0$.

Application 1: The unit distance problem

- Erdős (1946): $n^{1+c_1/\log \log n} \leq u(n) \leq O(n^{3/2})$
- Józsa and Szemerédi (1975): $u(n) = o(n^{3/2})$
- Beck and Spencer (1984): $u(n) = O(n^{13/9})$
- Spencer, Szemerédi, Trotter (1984): $u(n) = O(n^{4/3})$

500\$ for $u(n) = O(n^{1+\varepsilon})$ for every $\varepsilon > 0$.

Erdős' proof of $u(n) = O(n^{3/2})$:

Let $G = (V, E)$ be the graph with $V = P$ and edges $e = xy \in E(G)$ iff $|x - y| = 1$. The graph G is $K_{3,2}$ -free.



$K_{3,2}$

$$u(n) \leq \text{ex}(n, K_{3,2}) \leq O(n^{3/2}).$$



Application 1: The unit distance problem

Combinatorial Geometry

Two circles of unit radius cannot intersect in \geq three points

Extremal Graph Theory


The unit distance graph is $K_{3,2}$ -free

UB on $u(n)$

UB on $\text{ex}(n, K_{3,2})$


Application 2: Almost congruent unit triangles

n points in \mathbb{R}^2



Application 2: Almost congruent unit triangles

n points in \mathbb{R}^2



Up to $\binom{n}{3}$ triangles.

Question (Erdős, Purdy, 1975)


What is the maximum number of triangles almost congruent to the unit triangle?

Application 2: Almost congruent unit triangles


Question (Erdős, Purdy, 1975)

What is the maximum number of triangles almost congruent to the unit triangle?

unit triangle



ε -unit triangle




Application 2: Almost congruent unit triangles


Question (Erdős, Purdy, 1975)

What is the maximum number of triangles almost congruent to the unit triangle?

unit triangle



ε -unit triangle




Definition


$g(n, \varepsilon)$ = The maximum number of ε -unit triangles in a point set $P \subseteq \mathbb{R}^2$ of size n .

$$g(n) = \min_{\varepsilon > 0} g(n, \varepsilon).$$


Almost congruent unit triangles



Almost congruent unit triangles




Almost congruent unit triangles



$$g(n) \geq \left\lfloor \frac{n}{3} \right\rfloor \cdot \left\lfloor \frac{n+1}{3} \right\rfloor \cdot \left\lfloor \frac{n+2}{3} \right\rfloor \approx \frac{n^3}{27}$$

Almost congruent unit triangles



$$g(n) \geq \left\lfloor \frac{n}{3} \right\rfloor \cdot \left\lfloor \frac{n+1}{3} \right\rfloor \cdot \left\lfloor \frac{n+2}{3} \right\rfloor \approx \frac{n^3}{27}$$

Theorem (Balogh, C., Dumitrescu, 2023+)

For every positive integer n , we have



$$g(n) = \left\lfloor \frac{n}{3} \right\rfloor \cdot \left\lfloor \frac{n+1}{3} \right\rfloor \cdot \left\lfloor \frac{n+2}{3} \right\rfloor.$$

Application 2: Almost congruent unit triangles

We say a 3-graph G is *cancellative*, if there do not exist 3 edges A, B, C with $A \triangle B \subset C$.

Application 2: Almost congruent unit triangles

We say a 3-graph G is *cancellative*, if there do not exist 3 edges A, B, C with $A \triangle B \subset C$. Equivalently, G is cancellative iff G is $\{K_4^{3-}, F_5\}$ -free.

K_4^{3-}	123, 124, 134	
F_5	123, 124, 345	

Theorem (Bollobás, 1974)


The maximum number of edges in a cancellative 3-graph on n vertices is

$$\left\lfloor \frac{n}{3} \right\rfloor \cdot \left\lfloor \frac{n+1}{3} \right\rfloor \cdot \left\lfloor \frac{n+2}{3} \right\rfloor.$$

Application 2: Almost congruent unit triangles

Theorem (Bollobás, 1974)

$$\text{ex}(n, \{K_4^{3-}, F_5\}) = \left\lfloor \frac{n}{3} \right\rfloor \cdot \left\lfloor \frac{n+1}{3} \right\rfloor \cdot \left\lfloor \frac{n+2}{3} \right\rfloor.$$




Application 2: Almost congruent unit triangles


Observation

Let P be a set of 4 points in the plane where the minimum pairwise distance is at least 1. Then $\text{diam}(P) \geq \sqrt{2}$.

convex 4-gon



convex 3-gon




Application 2: Almost congruent unit triangles


Observation

Let P be a set of 4 points in the plane where the minimum pairwise distance is at least 1. Then $\text{diam}(P) \geq \sqrt{2}$.

convex 4-gon



convex 3-gon




Application 2: Almost congruent unit triangles


Observation

Let P be a set of 4 points in the plane where the minimum pairwise distance is at least 1. Then $\text{diam}(P) \geq \sqrt{2}$.

convex 4-gon



convex 3-gon




Application 2: Almost congruent unit triangles


Observation

Let P be a set of 4 points in the plane where the minimum pairwise distance is at least 1. Then $\text{diam}(P) \geq \sqrt{2}$.

convex 4-gon



convex 3-gon




Application 2: Almost congruent unit triangles


Observation

Let a be a positive real number and P be a set of 4 points in the plane where the minimum pairwise distance is at least a . Then $\text{diam}(P) \geq \sqrt{2}a$.

convex 4-gon



convex 3-gon



Application 2: Almost congruent unit triangles

Proof of UB on $g(n)$:

Application 2: Almost congruent unit triangles

Proof of UB on $g(n)$:

- Let $\varepsilon > 0$ sufficiently small and P a planar point set of size n .

Application 2: Almost congruent unit triangles

Proof of UB on $g(n)$:

- Let $\varepsilon > 0$ sufficiently small and P a planar point set of size n .
- Construct an auxiliary 3-graph $H(P, \varepsilon)$ with vertex set P and edges triples corresponding to ε -unit triangles.

Application 2: Almost congruent unit triangles

Proof of UB on $g(n)$:

- Let $\varepsilon > 0$ sufficiently small and P a planar point set of size n .
- Construct an auxiliary 3-graph $H(P, \varepsilon)$ with vertex set P and edges triples corresponding to ε -unit triangles.
- The shadowgraph of $H(P, \varepsilon)$ does not contain a copy of K_4 , otherwise there were 4 points with pairwise distance in $(1 - \varepsilon, 1 + \varepsilon)$.

Application 2: Almost congruent unit triangles

Proof of UB on $g(n)$:

- Let $\varepsilon > 0$ sufficiently small and P a planar point set of size n .
- Construct an auxiliary 3-graph $H(P, \varepsilon)$ with vertex set P and edges triples corresponding to ε -unit triangles.
- The shadowgraph of $H(P, \varepsilon)$ does not contain a copy of K_4 , otherwise there were 4 points with pairwise distance in $(1 - \varepsilon, 1 + \varepsilon)$.
- $H(P, \varepsilon)$ is K_4^{3-} and F_5 -free, because they both contain a K_4 in the shadowgraph.

Application 2: Almost congruent unit triangles

Proof of UB on $g(n)$:

- Let $\varepsilon > 0$ sufficiently small and P a planar point set of size n .
- Construct an auxiliary 3-graph $H(P, \varepsilon)$ with vertex set P and edges triples corresponding to ε -unit triangles.
- The shadowgraph of $H(P, \varepsilon)$ does not contain a copy of K_4 , otherwise there were 4 points with pairwise distance in $(1 - \varepsilon, 1 + \varepsilon)$.
- $H(P, \varepsilon)$ is K_4^{3-} and F_5 -free, because they both contain a K_4 in the shadowgraph.

$$g(n) \leq \text{ex}(n, \{K_4^{3-}, F_5\}) = \left\lfloor \frac{n}{3} \right\rfloor \cdot \left\lfloor \frac{n+1}{3} \right\rfloor \cdot \left\lfloor \frac{n+2}{3} \right\rfloor. \quad \square$$

Application 2: Almost congruent unit triangles

Combinatorial Geometry

4 points cannot have pairwise distance close to 1

Extremal Graph Theory

Shadowgraph of auxiliary 3-graph is K_4 -free


K_4^{3-}, F_5 -free

UB on $g(n)$


UB on $\text{ex}(n, \{K_4^{3-}, F_5\})$

Application 3: Almost congruent general triangles



triangle T



ε -congruent to T



Application 3: Almost congruent general triangles



Definition

$h_c(n, T, \varepsilon)$ = The maximum number of triangles ε -congruent to T in a point set $P \subseteq \mathbb{R}^2$ of size n .

$$h_c(n, T) = \min_{\varepsilon > 0} h_c(n, T, \varepsilon).$$

Observation

If two triangles T, T' are similar to each other, then $h_c(n, T) = h_c(n, T')$

Application 3: Almost congruent general triangles

Theorem (Balogh, C., Dumitrescu, 2023+)

Let T be a triangle and n be a positive integer.

- (a) Let T be a *right triangle*. Then, $h_c(n, T) \leq \frac{n^3}{16}$, and if additionally n is divisible by 4, then $h_c(n, T) = \frac{n^3}{16}$.

Application 3: Almost congruent general triangles


Theorem (Balogh, C., Dumitrescu, 2023+)

Let T be a triangle and n be a positive integer.


- (a) Let T be a **right triangle**. Then, $h_c(n, T) \leq \frac{n^3}{16}$, and if additionally n is divisible by 4, then $h_c(n, T) = \frac{n^3}{16}$.
- (b) Let T be of type **$(120^\circ, 30^\circ, 30^\circ)$** . Then, $h_c(n, T) \leq \frac{4}{81}n^3$, and if additionally n is divisible by 9, then $h_c(n, T) = \frac{4}{81}n^3$.
- (c) Let T be of type **$(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7})$** . Then, $h_c(n, T) \leq \frac{2}{49}n^3$, and if additionally n is divisible by 7, then $h_c(n, T) = \frac{2}{49}n^3$.
- (d) Let T be of type **$(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$** . Then, $h_c(n, T) \leq \frac{n^3}{25}$, and if additionally n is divisible by 5, then $h_c(n, T) = \frac{n^3}{25}$.
- (e) Let T be **not of type (a)-(d)**. Then, $h_c(n, T) \leq \frac{n^3}{27}$, and if additionally n is divisible by 3, then $h_c(n, T) = \frac{n^3}{27}$.

Application 3: Almost congruent general triangles


right triangle




$(120^\circ, 30^\circ, 30^\circ)$




$\left(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7}\right)$



$(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$



arbitrary T



Application 3: Almost congruent general triangles

Definition

$H(P, T, \varepsilon) :=$ The 3-graph with vertex set P and edges the triples corresponding to triangles ε -congruent to T .

Combinatorial Geometry

Some forbidden configuration
on P for T

UB on $h_c(n, T)$

Extremal Graph Theory



$H(P, T, \varepsilon)$ is
 $\mathcal{F}(T)$ -free for small ε

UB on $\text{ex}(n, \mathcal{F}(T))$

Application 3: Almost congruent general triangles

Theorem (Balogh, C., Dumitrescu)

(a) Let T be a right triangle. Then, $h_c(n, T) \leq \frac{n^3}{16}$.

$F_{3,2}$	123, 145, 245, 345	
J_4	123, 124, 125, 134, 135, 145	

If T is not $(90^\circ, 60^\circ, 30^\circ)$, then

$$h_c(n, T) \leq \text{ex}(n, \{F_{3,2}, J_4\}) = \frac{n^3}{16}(1 + o(1)),$$




by Falgas-Ravry and Vaughan.



Application 3: Almost congruent general triangles

Theorem (Balogh, C., Dumitrescu)

(c) Let T be of type $\left(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7}\right)$. Then, $h_c(n, T) \leq \frac{2}{49} n^3$.

$F_{3,2}$	123, 145, 245, 345	
K_4^{3-}	123, 124, 134	
C_5^3	123, 234, 345, 145, 125	

$$h_c(n, T) \leq \text{ex}(n, \{K_4^{3-}, F_{3,2}, C_5^3\}) = \frac{2}{49} n^3 (1 + o(1)),$$

by Falgas-Ravry and Vaughan.



Application 3: A new strategy

Combinatorial Geometry

Some forbidden configuration
on P for T

Extremal Graph Theory

$H(P, T, \varepsilon)$ is
 $\mathcal{F}(T)$ -free for small ε

UB on $h_c(n, T)$

Application 3: A new strategy

Combinatorial Geometry

Some forbidden configuration
on P for T

Extremal Graph Theory

$H(P, T, \varepsilon)$ is
 $\mathcal{F}(T)$ -free for small ε



Weighted hypergraph
Turán problem

UB on $h_c(n, T)$



Application 3: Hypergraph Lagrangians

Let H be an n -vertex 3-graph. The *Lagrangian polynomial* of H is

$$\lambda_H(x_1, \dots, x_n) := \sum_{ijk \in H} x_i x_j x_k,$$

and the *Lagrangian* of H is

$$\lambda(H) := \max\{\lambda_H(x_1, \dots, x_n) : (x_1, x_2, \dots, x_n) \in \Delta_n\},$$

where $\Delta_n = \{(x_1, x_2, \dots, x_n) \in [0, 1]^n : x_1 + x_2 + \dots + x_n = 1\}$.

Application 3: Hypergraph Lagrangians

Let H be an n -vertex 3-graph. The *Lagrangian polynomial* of H is

$$\lambda_H(x_1, \dots, x_n) := \sum_{ijk \in H} x_i x_j x_k,$$

and the *Lagrangian* of H is

$$\lambda(H) := \max\{\lambda_H(x_1, \dots, x_n) : (x_1, x_2, \dots, x_n) \in \Delta_n\},$$

where $\Delta_n = \{(x_1, x_2, \dots, x_n) \in [0, 1]^n : x_1 + x_2 + \dots + x_n = 1\}$.

Example: $V(K_4^{3-}) = \{1, 2, 3, 4\}$, $E(K_4^{3-}) = \{123, 124, 134\}$

$$\begin{aligned}\lambda(K_4^{3-}) &= \max_{x_i \geq 0, x_1 + x_2 + x_3 + x_4 = 1} x_1 x_2 x_3 + x_1 x_2 x_4 + x_1 x_3 x_4 \\ &= \max_{0 \leq x \leq 1} 3x \left(\frac{1-x}{3} \right)^2 = \frac{4}{81}\end{aligned}$$

Application 3: The key lemma

$H(P, T) :=$ The 3-graph with vertex set P and edges the triples corresponding to triangles congruent to T .

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Application 3: Sketch of proof of key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Sketch of proof:

$$\frac{h_c(n, T)}{n^3} = \frac{e(H(P, T, \varepsilon))}{n^3} = \lambda_{H(P, T, \varepsilon)}\left(\frac{1}{n}, \dots, \frac{1}{n}\right) \leq \lambda(H(P, T, \varepsilon)).$$

Application 3: Sketch of proof of key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Sketch of proof:

$$\frac{h_c(n, T)}{n^3} = \frac{e(H(P, T, \varepsilon))}{n^3} = \lambda_{H(P, T, \varepsilon)}\left(\frac{1}{n}, \dots, \frac{1}{n}\right) \leq \lambda(H(P, T, \varepsilon)).$$

Application 3: Sketch of proof of key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Sketch of proof:

$$\frac{h_c(n, T)}{n^3} = \frac{e(H(P, T, \varepsilon))}{n^3} = \lambda_{H(P, T, \varepsilon)}\left(\frac{1}{n}, \dots, \frac{1}{n}\right) \leq \lambda(H(P, T, \varepsilon)).$$

- Let $\mathbf{x} \in \Delta_n$ be such that $\lambda(H(P, T, \varepsilon)) = \lambda_{H(P, T, \varepsilon)}(\mathbf{x})$ with the fewest non-zero entries.

Application 3: Sketch of proof of key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Sketch of proof:

$$\frac{h_c(n, T)}{n^3} = \frac{e(H(P, T, \varepsilon))}{n^3} = \lambda_{H(P, T, \varepsilon)}\left(\frac{1}{n}, \dots, \frac{1}{n}\right) \leq \lambda(H(P, T, \varepsilon)).$$

- Let $\mathbf{x} \in \Delta_n$ be such that $\lambda(H(P, T, \varepsilon)) = \lambda_{H(P, T, \varepsilon)}(\mathbf{x})$ with the fewest non-zero entries.
- If there are two vertices, not contained in an edge, with positive weights, we can move weights from one to the other.

Application 3: Sketch of proof of key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Sketch of proof:

$$\frac{h_c(n, T)}{n^3} = \frac{e(H(P, T, \varepsilon))}{n^3} = \lambda_{H(P, T, \varepsilon)}\left(\frac{1}{n}, \dots, \frac{1}{n}\right) \leq \lambda(H(P, T, \varepsilon)).$$

- Let $\mathbf{x} \in \Delta_n$ be such that $\lambda(H(P, T, \varepsilon)) = \lambda_{H(P, T, \varepsilon)}(\mathbf{x})$ with the fewest non-zero entries.
- If there are two vertices, not contained in an edge, with positive weights, we can move weights from one to the other.
- Compactness argument; the points with positive weight form a 3-distance set. Shinohara: At most 7 points have positive weight. \square

Application 3: Applying the key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Theorem (Balogh, C., Dumitrescu)

(d) *Let T be $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then, $h_c(n, T) \leq \frac{n^3}{25}$.*

Application 3: Applying the key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Theorem (Balogh, C., Dumitrescu)

(d) Let T be $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then, $h_c(n, T) \leq \frac{n^3}{25}$.

- Q is a 2-distance set $\rightarrow |Q| \leq 5$.

Application 3: Applying the key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Theorem (Balogh, C., Dumitrescu)

(d) Let T be $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then, $h_c(n, T) \leq \frac{n^3}{25}$.

- Q is a 2-distance set $\rightarrow |Q| \leq 5$.
- If $|Q| = 5$, then because Q is a 2-distance set, Q is a regular pentagon. Then $H(Q, T) \cong C_5^3$ and $\lambda(C_5^3) = \frac{1}{25}$.

Application 3: Applying the key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Theorem (Balogh, C., Dumitrescu)

(d) Let T be $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then, $h_c(n, T) \leq \frac{n^3}{25}$.

- Q is a 2-distance set $\rightarrow |Q| \leq 5$.
- If $|Q| = 5$, then because Q is a 2-distance set, Q is a regular pentagon. Then $H(Q, T) \cong C_5^3$ and $\lambda(C_5^3) = \frac{1}{25}$.
- $|Q| = 4$ is not possible, because there is no K_4^{3-} -free 3-graph on 4 vertices with complete shadow-graph.

Application 3: Applying the key lemma

Lemma (Balogh, C., Dumitrescu)

Let T be a triangle and n be a positive integer. Then there exists a point set Q of size $|Q| \leq 7$ such that in $H(Q, T)$ every pair of vertices is contained in an edge, and $h_c(n, T) \leq n^3 \lambda(H(Q, T))$.

Theorem (Balogh, C., Dumitrescu)

(d) Let T be $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then, $h_c(n, T) \leq \frac{n^3}{25}$.

- Q is a 2-distance set $\rightarrow |Q| \leq 5$.
- If $|Q| = 5$, then because Q is a 2-distance set, Q is a regular pentagon. Then $H(Q, T) \cong C_5^3$ and $\lambda(C_5^3) = \frac{1}{25}$.
- $|Q| = 4$ is not possible, because there is no K_4^{3-} -free 3-graph on 4 vertices with complete shadow-graph.
- If $|Q| = 3$, then $H(Q, T)$ is just a single edge and

$$\lambda(H(Q, T)) = \max_{x_i \geq 0, x_1 + x_2 + x_3 = 1} x_1 x_2 x_3 = \frac{1}{27}.$$



Application 3: Applying the key lemma

Theorem (Balogh, C., Dumitrescu)

- (e) Let T be not right angled, and not $(120^\circ, 30^\circ, 30^\circ)$, $\left(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7}\right)$, $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then,
- $$h_c(n, T) \leq \frac{n^3}{27}.$$

Application 3: Applying the key lemma

Theorem (Balogh, C., Dumitrescu)

(e) Let T be not right angled, and not $(120^\circ, 30^\circ, 30^\circ)$, $\left(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7}\right)$, $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then,
$$h_c(n, T) \leq \frac{n^3}{27}.$$

- Q is a 3-distance set, $|Q| \leq 7$ and $H(Q, T)$ is $\{K_4^{3-}, C_5^3, F_{3,2}\}$ -free

Application 3: Applying the key lemma

Theorem (Balogh, C., Dumitrescu)

(e) Let T be not right angled, and not $(120^\circ, 30^\circ, 30^\circ)$, $\left(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7}\right)$, $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then,
$$h_c(n, T) \leq \frac{n^3}{27}.$$

- Q is a 3-distance set, $|Q| \leq 7$ and $H(Q, T)$ is $\{K_4^{3-}, C_5^3, F_{3,2}\}$ -free
- If $|Q| = 3$, then $H(Q, T)$ is a single edge and $\lambda(H(Q, T)) = \frac{1}{27}$.

Application 3: Applying the key lemma

Theorem (Balogh, C., Dumitrescu)

(e) *Let T be not right angled, and not $(120^\circ, 30^\circ, 30^\circ)$, $(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7})$, $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then,*

$$h_c(n, T) \leq \frac{n^3}{27}.$$

- Q is a 3-distance set, $|Q| \leq 7$ and $H(Q, T)$ is $\{K_4^{3-}, C_5^3, F_{3,2}\}$ -free
- If $|Q| = 3$, then $H(Q, T)$ is a single edge and $\lambda(H(Q, T)) = \frac{1}{27}$.
- $|Q| = 4, 5$ is not possible, because there is no $\{K_4^{3-}, C_5^3, F_{3,2}\}$ -free $|Q|$ -vertex 3-graph with complete shadow graph.

Application 3: Applying the key lemma

Theorem (Balogh, C., Dumitrescu)









(e) *Let T be not right angled, and not $(120^\circ, 30^\circ, 30^\circ)$, $(\frac{4 \cdot 180^\circ}{7}, \frac{2 \cdot 180^\circ}{7}, \frac{180^\circ}{7})$, $(108^\circ, 36^\circ, 36^\circ)$ or $(72^\circ, 72^\circ, 36^\circ)$. Then,*

$$h_c(n, T) \leq \frac{n^3}{27}.$$

- Q is a 3-distance set, $|Q| \leq 7$ and $H(Q, T)$ is $\{K_4^{3-}, C_5^3, F_{3,2}\}$ -free
- If $|Q| = 3$, then $H(Q, T)$ is a single edge and $\lambda(H(Q, T)) = \frac{1}{27}$.
- $|Q| = 4, 5$ is not possible, because there is no $\{K_4^{3-}, C_5^3, F_{3,2}\}$ -free $|Q|$ -vertex 3-graph with complete shadow graph.
- If $|Q| = 6, 7$, we use a characterization of 3-distance sets by Shinohara.

- If $|Q| = 6, 7$, then T has 3 different sides. We use a characterization of 3-distance sets by Shinohara:

- If $|Q| = 6, 7$, then T has 3 different sides. We use a characterization of 3-distance sets by Shinohara:



Further questions

One of my favourite questions:

- Determine the maximum number of acute triangles in a planar point set of size n .

A Question for graduate students:

- Similar questions but $P \subseteq \mathbb{R}^d$ for $d \geq 3$.

Thank you!

Thank you for your attention!

