

## Our goal

We need to determine if an argument like the one below is a good one. It's deductive. So, it's good if it is both valid and sound. Our focus at the moment is validity:

Bill: "God must exist."

Jill: "How do you know."

Bill: "Well if the Bible is accurate, then God exists."

Jill: "Why should I believe the Bible?"

Bill: "Well, if God exists, then it has got to be accurate!"

## Validity

An argument is valid just if it is impossible for the conclusion to be false and **all** of the premises to be true.

## Invalidity

An argument is invalid just if it is possible for the conclusion to be false and *all* of the premises to be true.

## Truth Tables

Truth tables allow us determine if it is possible for the conclusion to be false and, at the same time, the premises to be true.

Strategy: construct a large truth table with smaller truth tables as their parts. One part is a truth-table for the conclusion. There is also one truth-table for each of the premises. After constructing the large truth-table, we will see if any line has a false conclusion and true premises.

## Example

Bill: "God must exist."

Jill: "How do you know."

Bill: "Well if the Bible is accurate, then God exists."

Jill: "Why should I believe the Bible?"

Bill: "Well, if God exists, then it has got to be accurate!"

'P': God exists

'Q': The Bible is accurate

- P1: If God exists, then the Bible is accurate. ( $p \rightarrow q$ )
- P2: If the Bible is accurate, then God exists. ( $q \rightarrow p$ )
- C: God exists. ( $p$ )