Problem 1

Theorem 1 (All pairs have different sums). Prove or disprove: It is possible to label each square of an 8 x 8 checkerboard with an integer, so that no two squares are labeled with the same integer and no two adjacent squares (up, down, left, right) have labels that differ by 8 or more.

Proof (long form).

- 1. Assume you have filled some chessboard with integers to satisfy the theorem.
- 1.1. Construct a set S by doing the following...
- 1.2. Add the square with the smallest integer not in S into S, stopping when either every row has at least one square in S or when every column has at least one square in S.
- 1.3. Without loss of generality, we can assume that every row will have at least one square in S (if it's columns, just rotate the chessboard).
- 1.4. Call T the set of chessboard squares adjacent to those in S but not in S.
- 1.5. Consider the largest numbered square in S, s, where the number in s is num(s).
- 1.6. Before adding s to S, there were no squares in S in this row.
- 1.7. Likewise, no row was full (if one had been, we would have already stopped adding to S since there was a square in every column).
- 1.8. Therefore, after the addition of s into S, no row will be completely contained in S.
- 1.9. Conversely, every row will have at least one square in S.
- 1.10. Therefore, there must be at least one square in T in every row.
- 1.11. That is, $|T| \geq 8$.
- 1.12. Since there are at least 8 squares in T and the numbers in the squares in T must be unique integers > num(s), the lower bound on the largest numbered square in T will be num(s) + 8.
- 1.13. However, by the definition of the theorem, there is no square in S that the largest square in T can be adjacent to.
- 1.14. Contradiction.
- 2. By block 1 you cannot fill a chessboard with integers to satisfy the theorem.
- 3. That is, the theorem is false.