Math 120 Optimization: Homework 5

Stanley Cohen (scohe001)

Solve the following linear programming problems using the matrix form (i.e. the tableau) of simplex method.

Problem 1:

maximize
$$2x_1 + x_2$$

subject to $x_1 \le 2$
 $x_1 + x_2 \le 3$
 $x_1 + 2x_2 \le 5$
 $x_1, x_2 \ge 0$.

Putting this into standard form will make it look like:

minimize
$$-2x_1 - x_2$$

subject to $x_1 + s_1 = 2$
 $x_1 + x_2 + s_2 = 3$
 $x_1 + 2x_2 + s_3 = 5$
 $x_1, x_2, s_1, s_2, s_3 \ge 0$.

Given this LP, the initial tableau will look like:

$$\begin{bmatrix}
1 & 0 & 1 & 0 & 0 & 2 \\
1 & 1 & 0 & 1 & 0 & 3 \\
1 & 2 & 0 & 0 & 1 & 5 \\
-2 & -1 & 0 & 0 & 0 & 0
\end{bmatrix}$$

We'll column 1 look like column 3 to get:

$$\begin{bmatrix}
1 & 0 & 1 & 0 & 0 & 2 \\
0 & 1 & -1 & 1 & 0 & 1 \\
0 & 2 & -1 & 0 & 1 & 3 \\
0 & -1 & 2 & 0 & 0 & 4
\end{bmatrix}$$

Next we'll make column 2 look like column 4 to get:

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 2 \\ 0 & 1 & -1 & 1 & 0 & 1 \\ 0 & 0 & 1 & -2 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 & 5 \end{bmatrix}$$

So we get an optimal value of 5 when $(x_1, x_2) = (2, 1)$.

Problem 2:

maximize
$$6x_1 + 4x_2 + 7x_3 + 5x_4$$

subject to $x_1 + 2x_2 + x_3 + 2x_4 \le 20$
 $6x_1 + 5x_2 + 3x_3 + 2x_4 \le 100$
 $3x_1 + 4x_2 + 9x_3 + 12x_4 \le 75$
 $x_1, x_2, x_3, x_4 \ge 0$.

Puttin this into standard form will make it look like:

minimize
$$-6x_1 - 4x_2 - 7x_3 - 5x_4$$

subject to $x_1 + 2x_2 + x_3 + 2x_4 + s_1 = 20$
 $6x_1 + 5x_2 + 3x_3 + 2x_4 + s_2 = 100$
 $3x_1 + 4x_2 + 9x_3 + 12x_4 + s_3 = 75$
 $x_1, x_2, x_3, x_4, s_1, s_2, s_3 \ge 0$.

Given this LP, the inital tableau will look like:

$$\begin{bmatrix} 1 & 2 & 1 & 2 & 1 & 0 & 0 & 20 \\ 6 & 5 & 3 & 2 & 0 & 1 & 0 & 100 \\ 3 & 4 & 9 & 12 & 0 & 0 & 1 & 75 \\ -6 & -4 & -7 & -5 & 0 & 0 & 0 & 0 \end{bmatrix}$$

First we'll make column 3 look like column 7:

$$\begin{bmatrix} 2/3 & 14/9 & 0 & 2/3 & 1 & 0 & -1/9 & 35/3 \\ 5 & 11/3 & 0 & -2 & 0 & 1 & -1/3 & 75 \\ 1/3 & 4/9 & 1 & 4/3 & 0 & 0 & 1/9 & 25/3 \\ -11/3 & -8/9 & 0 & 13/3 & 0 & 0 & 7/9 & 175/3 \end{bmatrix}$$

Next we'll make column 1 look like column 6:

$$\begin{bmatrix} 0 & 16/15 & 0 & 14/15 & 1 & -2/15 & -1/15 & 5/3 \\ 1 & 11/15 & 0 & -2/5 & 0 & 1/5 & -1/15 & 25 \\ 0 & 1/5 & 1 & 22/15 & 0 & -1/15 & 2/15 & 10/3 \\ 0 & 9/5 & 0 & 11/5 & 0 & 11/15 & 8/15 & 340/3 \end{bmatrix}$$

So we get an optimal value of 340/3 when $(x_1, x_2, x_3, x_4) = (15, 0, 10/3, 0)$.

Problem 3:

minimize
$$x_1 + x_2 - x_3$$

subject to $2x_1 + x_2 + x_3 \le 14$
 $4x_1 + 2x_2 + 3x_3 \le 28$
 $2x_1 + 5x_2 + 5x_3 \le 30$
 $x_1, x_2, x_3 \ge 0$.

Puttin this into standard form will make it look like:

minimize
$$x_1 + x_2 - x_3$$

subject to $2x_1 + x_2 + x_3 + s_1 = 14$
 $4x_1 + 2x_2 + 3x_3 + s_2 = 28$
 $2x_1 + 5x_2 + 5x_3 + s_3 = 30$
 $x_1, x_2, x_3, s_1, s_2, s_3 \ge 0$.

Given this LP, the inital tableau will look like:

$$\begin{bmatrix} 2 & 1 & 1 & 1 & 0 & 0 & | & 14 \\ 4 & 2 & 3 & 0 & 1 & 0 & | & 28 \\ 2 & 5 & 5 & 0 & 0 & 1 & | & 30 \\ 1 & 1 & -1 & 0 & 0 & 0 & | & 0 \end{bmatrix}$$

First we'll make column 3 look like column 6:

$$\begin{bmatrix} 8/5 & 0 & 0 & 1 & 0 & -1/5 & 8 \\ 14/5 & -1 & 0 & 0 & 1 & -3/5 & 4 \\ 2/5 & 1 & 1 & 0 & 0 & 1/5 & 6 \\ 7/5 & 2 & 0 & 0 & 0 & 1/5 & 6 \end{bmatrix}$$

So we get an optimal value of -6 when $(x_1, x_2, x_3) = (0, 0, 6)$.

Problem 4:

Puttin this into standard form will make it look like:

Given this LP, the inital tableau will look like:

$$\begin{bmatrix} 3 & 1 & 1 & 4 & 1 & 0 & 0 & | & 12 \\ 1 & -3 & 2 & 3 & 0 & 1 & 0 & | & 7 \\ 2 & 1 & 3 & -1 & 0 & 0 & 1 & | & 10 \\ -2 & -4 & -3 & -1 & 0 & 0 & 0 & | & 0 \end{bmatrix}$$