




INTRODUCTON

- Numerical simulations of decadal climate are done at resolutions far coarser than the natural scale of precipitation. To even have a chance of understanding future precipitation extremes we must reconcile the relation between the statistics of broad-scale precipitation and high resolution observations.
- To this end The Department of Energy's ARM Climate Research Facility operates a network of 5 and 3 cm scanning radar systems.
- Fixed sites are at the Azores, Barrow on the North Slope of Alaska and a multi-scale heterogeneous network on the Southern Great Plains of Oklahoma.



ACHIEVING INSIGHT WITH THE COMMUNITY: THE PYTHON ARM RADAR TOOLKIT

- Weather radars are not a new invention, first academic mention in Bent et al. (1943).
- Massive advances in computing and radar software has not kept up.
- They Python ARM Radar Toolkit, Py-ART is a data model driven architecture for interactively and offline processing of active remote sensing data. Open source and, using GitHub, community based.
- Part of a larger growing international community of codes, see Heistermann et al. (2014)
- Twenty four forks, eight active contributors from multiple agencies and nations. Broad user base.



LINKS

The Python ARM Radar Toolkit :




This is an open source poster, for notebooks, code and tex:



RAW DATA TO QUANTITATIVE PRECIPITATION ESTIMATES

- Raw collected radar data in engineering units is unsuitable for comparison with model data.
- Shorter wavelength radar have a higher attenuation cross section. However Signal to noise in phase information much higher and calibration insensitive.
- Measured phase is a mix of propagation phase, phase shift on backscatter and artifacts:
 $\phi_{dp}^{total}(r) = \phi_{dp}^{prop}(r) + \delta(r) + E(r)$
- When calculating Specific Differential Phase, K_{dp} only the propagation component should be considered, $K_{dp} = \frac{\partial \phi_{dp}^{prop}}{\partial r}$.
- Method of Giangrande et al. (2013) used to extract ϕ_{dp}^{prop} and a 20 point sobel filter
 $K_{dp} = \phi_{dp}^{max} * I_{20} = \sum_{M=1}^{19} \phi_{dp}(r-M)f(M)$ where $f(M)$ is a linear ramp through zero.




- Specific attenuation is calculated using K_{dp} and Z_e using a method after Gu et al. (2011) and is used as an estimator for rainfall using a method after Ryzhkov et al. (2014).
- In Giangrande et al. (2014) it was shown that specific attenuation at short wavelengths performed as well as retrievals at S-band wavelengths.

MAPPING: OBJECTIVE ANALYSIS

- Radar data is on θ and ϕ coordinates, there is a need to estimate on different coordinates systems (Cartesian, Sigma, pressure).
- Py-ART tags each gate with an estimate of its central coordinate and inserts these into a cloud.
- For propagation insensitive variables (not radial velocity or ϕ_{dp}), gates can be drawn from multiple radars to be estimated onto a single grid.
- In Py-ART the act of gridding takes in a n -tuple of radar objects and returns a grid object which can be saved to a CF-Radial complaint file.

```
mesh_mapped_x = pyart.map.grid_from_radars((xmw_radar,xsw_radar,xse_radar),
grid_shape=(35, 401, 401),
grid_limits=((0, 17000), (-50000, 40000), (-60000, 40000)),
grid_origin = (36.57861, -97.363611),
fields=['corrected_reflectivity', 'rain_rate_A', 'reflectivity'],
refl_field='corrected_reflectivity',
max_refl=100.)
```



ADVECTIVE INTERPOLATION

- Simply accumulating precipitation retrievals by integrating can create a "chain of pearls effect" due to the lack of spatial coherency between successive radar scans.
- This can create false information and skew any interpretation of the scale of rainfall.
- The solution is to calculate the image advection and then generate sub-temporal-scale images and average:

$$G(t + \Delta t, z, y, x) = \left(1 - \frac{t + \Delta t - t_1}{t_2 - t_1}\right) G_1(t_1, z, y + v\Delta t, x + u\Delta t) + \frac{t + \Delta t - t_1}{t_2 - t_1} G_2(t_2, z, y - v\Delta t, x - u\Delta t)$$


- Setting $t_1 = 0$

$$G(t + \Delta t, z, y, x) = \left(1 - \frac{t + \Delta t}{t_2}\right) G_1(t_1, z, y + v\Delta t, x + u\Delta t) + \frac{t + \Delta t}{t_2} G_2(t_2, z, y - v\Delta t, x - u\Delta t)$$

- In a nice novel twist (we Py-ART developers love applying math to radar) the advection can be determined from the cross correlation phase between the two images, $r = \mathcal{F}^{-1}\{C\}$.
- Borrowing liberally from Wikipedia:

$$\begin{aligned} \mathbf{G}_{t1} &= \mathcal{F}\{R_{t1}\}, \mathbf{G}_{t2} = \mathcal{F}\{R_{t2}\} \\ C &= \frac{\mathbf{G}_{t1} \circ \mathbf{G}_{t2}^*}{|\mathbf{G}_{t1} \circ \mathbf{G}_{t2}^*|} \\ r &= \mathcal{F}^{-1}\{C\} \\ \Delta x, \Delta y &= \operatorname{argmax}\{r\} \end{aligned}$$

- where \mathcal{F} is the Fourier transform, $*$ is the complex conjugate and \circ is the element wise multiplication.



REFERENCES

- Bent, A. E., Massachusetts Institute of Technology., and Radiation Laboratory., 1943: *Radar echoes from atmospheric phenomena*. Radiation Laboratory, Massachusetts Institute of Technology, [Cambridge, Mass.].
- Giangrande, S. E., S. Collis, A. K. Theisen, and A. Tokay, 2014: Precipitation estimation from the ARM distributed radar network during the MC3e campaign. *J. Appl. Meteor. Climatol.*, doi:10.1175/JAMC-D-13-0321.1. URL <http://journals.ametsoc.org/doi/abs/10.1175/JAMC-D-13-0321.1>
- Giangrande, S. E., R. McGraw, and L. Lei, 2013: An application of linear programming to polarimetric radar differential phase processing. *Journal of Atmospheric and Oceanic Technology*, **30**, 1716–1729, doi:10.1175/JTECH-D-12-00147.1. URL <http://journals.ametsoc.org/doi/abs/10.1175/JTECH-D-12-00147.1>
- Gu, J.-Y., A. Ryzhkov, P. Zhang, P. Neille, M. Knight, B. Wolf, and D.-I. Lee, 2011: Polarimetric attenuation correction in heavy rain at c band. *J. Appl. Meteor. Climatol.*, **50**, 39–58, doi:10.1175/2010JAMC2258.1. URL <http://journals.ametsoc.org/doi/abs/10.1175/2010JAMC2258.1>
- Heistermann, M., S. Collis, M. J. Dixon, S. Giangrande, J. J. Helmus, B. Kelley, J. Koistinen, D. B. Michelson, M. Peura, T. Pfaff, and D. B. Wolff, 2014: The emergence of open source software for the weather radar community. *Bull. Amer. Meteor. Soc.*, doi:10.1175/BAMS-D-13-00240.1. URL <http://journals.ametsoc.org/doi/abs/10.1175/BAMS-D-13-00240.1>
- Ryzhkov, A., M. Diederich, P. Zhang, and C. Simmer, 2014: Potential utilization of specific attenuation for rainfall estimation, mitigation of partial beam blockage, and radar networking. *J. Atmos. Oceanic Technol.*, **31**, 599–619, doi:10.1175/JTECH-D-13-00038.1. URL <http://journals.ametsoc.org/doi/abs/10.1175/JTECH-D-13-00038.1>