

MATH4404 Theorems

Conal O'Leary

October 2021

The theorem about the existence of a completion of a normed space.

Theorem 1. If E is a normed linear space, then there exists a complete normed space \tilde{E} such that

- i) $E \subset \tilde{E}$
- ii) $\|x\|_E = \|x\|_{\tilde{E}}$
- iii) E is dense in \tilde{E}

Proof. Construct \tilde{E} as equivalence classes of Cauchy Sequences $(x_1, x_2, \dots) \sim (y_1, y_2, \dots)$ if $(x_1, y_1, x_2, y_2, \dots)$ is also Cauchy. We embed each $x \in E$ as $(x, x, x, \dots) \in \tilde{E}$. This is certainly a Cauchy sequence and as such, well defined.

Now we will show that our operations $(+, \cdot)$ are well defined. Take two sequences $(x_n)_{n=1}^\infty$ and $(y_n)_{n=1}^\infty$ in \tilde{E} such that $(x_n)_{n=1}^\infty \in [x']$ and $(y_n)_{n=1}^\infty \in [y']$. Our operations is defined as below,

$$(x_1, x_2, \dots) + (y_1, y_2, \dots) = (x_1 + y_1, x_2 + y_2, \dots)$$

We wish to show that this sequence is an element of $[x' + y']$. Consider the sequence,

$$(z_n)_{n=1}^\infty = (x_1 + y_1, x' + y', x_2 + y_2, x' + y', \dots)$$

We will show this is Cauchy.

Given $\varepsilon > 0$, there exists large enough $N \in \mathbb{N}$ such that (x_n) and (y_n) behave as expected. Now for all $n > m$, there are two cases,

$$\begin{aligned} \|z_n - z_m\| &= \|x_n + y_n - x_m - y_m\| \\ &\leq \|x_n - x_m\| + \|y_n - y_m\| \\ &< \varepsilon \end{aligned}$$

or

$$\begin{aligned} \|z_n - z_m\| &= \|x_n + y_n - x' - y'\| \\ &\leq \|x_n - x'\| + \|y_n - y'\| \\ &< \varepsilon \end{aligned}$$

So we have $(z_n)_{n=1}^\infty \in [x' + y']$.

Multiplication is defined as expected. The norm in \tilde{E} is defined as follows,

$$\|[(x_1, x_2, \dots)]\|_{\tilde{E}} = \lim_{n \rightarrow \infty} \|x_n\|_E$$

We will now show that the properties of the norm hold. It is trivial that $\|\cdot\|_{\tilde{E} \geq 0}$ since $\|\cdot\|_E \geq 0$. We will now show that $x = 0 \iff \|x\|_{\tilde{E}} = 0$

\implies

Let $(x_n)_{n=1}^\infty$ be an element of $[0]$, so we know for every $\varepsilon > 0$ and large enough n , $\|x_n - 0\| < \varepsilon$. Continuing,

$$\begin{aligned} \|[(x_1, x_2, \dots)]\|_{\tilde{E}} &= \lim_{n \rightarrow \infty} \|x_n\|_E \\ &< \varepsilon \end{aligned}$$

This holds for every $\varepsilon > 0$ So we have $\|[(x_1, x_2, \dots)]\| = 0$.

\impliedby

Assume $\|[(x_1, x_2, \dots)]\| = 0$, we wish to show $(x_1, x_2, \dots) \in [0]$. By the continuity of the norm (Reverse Triangle Inequality),

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \|x_n\| \\ &= \|\lim_{n \rightarrow \infty} x_n\| \end{aligned}$$

We have that E is a normed space so it holds that $\|x_n\|_E = 0 \implies x_n = 0$. So it is quite clear that $(x_1, 0, x_2, 0, \dots)$ is Cauchy. Thus, $(x_n)_{n=1}^\infty \in [0]$.

Now we will prove the Triangle Inequality, let $(x_n)_{n=1}^\infty$ and $(y_n)_{n=1}^\infty$ in \tilde{E} . Wish to show $\|[(x_1, x_2, \dots)] + [(y_1, y_2, \dots)]\| \leq \|[(x_1, x_2, \dots)]\| + \|[(y_1, y_2, \dots)]\|$

$$\begin{aligned} \|[(x_1, x_2, \dots)] + [(y_1, y_2, \dots)]\| &= \lim_{n \rightarrow \infty} \|x_n + y_n\| \\ &\leq \lim_{n \rightarrow \infty} \|x_n\| + \|y_n\| \\ &= \lim_{n \rightarrow \infty} \|x_n\| + \lim_{n \rightarrow \infty} \|y_n\| \end{aligned}$$

Since $(\|x_n\|)_{n=1}^\infty$ and $(\|y_n\|)_{n=1}^\infty$ are Cauchy \mathbb{R} and \mathbb{R} is complete. So $\|\cdot\|_{\tilde{E}}$ is a valid norm. We will now show that \tilde{E} is indeed complete.

Take some Cauchy sequence $(\tilde{x}_n)_{n=1}^\infty \subset \tilde{E}$, will show there exists an $\tilde{x} \in \tilde{E}$ such that $\lim_{n \rightarrow \infty} \tilde{x}_n = \tilde{x}$. Since E is dense in \tilde{E} , for every $n \in \mathbb{N}$ there exists $x_n \in E$ such that

$$\|(x_n, x_n, \dots) - \tilde{x}_n\| < \frac{1}{n}$$

$$(x_1, x_1, x_1, \dots) - \tilde{x}_2 < 1$$

$$(x_2, x_2, x_2, \dots) - \tilde{x}_2 < \frac{1}{2}$$

Define $\tilde{x} = [(x_1, x_2, x_3, \dots)] = [(x_n)_{n=1}^\infty]$. We now need to verify that $(x_n)_{n=1}^\infty$ is indeed Cauchy.

$$\begin{aligned} \|x_k - x_m\|_E &= \|(x_k, x_k, \dots) - (x_m, x_m, \dots)\|_{\tilde{E}} \\ &\leq \|(x_k, x_k, \dots) - \tilde{x}_k\| + \|\tilde{x}_k - \tilde{x}_m\|_{\tilde{E}} + \|\tilde{x}_m - (x_m, x_m, \dots)\|_{\tilde{E}} \\ &\leq \frac{1}{k} + \frac{1}{m} + \|\tilde{x}_k - \tilde{x}_m\|_{\tilde{E}} \end{aligned}$$

Since $(\tilde{x}_n)_{n=1}^\infty$ is Cauchy this goes to 0 as $k, m \rightarrow \infty$. Thus $(x_n)_{n=1}^\infty$ is Cauchy.

$$\begin{aligned} \|\tilde{x}_n - \tilde{x}\|_{\tilde{E}} &= \|\tilde{x}_n - (x_n, x_n, \dots)\|_{\tilde{E}} + \|(x_n, x_n, \dots) - \tilde{x}\|_{\tilde{E}} \\ &\leq \frac{1}{n} + \lim_{m \rightarrow \infty} \|x_n - x_m\|_E \end{aligned}$$

Since $(x_n)_{n=1}^\infty$ we have that this goes to 0 as $n, m \rightarrow \infty$, and so $(\tilde{x}_n)_{n=1}^\infty$ converges to \tilde{x} . \square

The Bunyakovsky–Cauchy–Schwarz inequality.

Theorem 2. Let H be an inner product space, for all $x, y \in H$,

$$|(x, y)|^2 \leq (x, x)(y, y)$$

Proof. If $x = 0$ or $y = 0$ use linearity to prove. Assume $y \neq 0$. Let $\lambda = \frac{(x, y)}{(y, y)}$. Note that

$$\begin{aligned} (x - \lambda y, y) &= (x, y) - \lambda(y, y) \\ &= (x, y) - \frac{(x, y)}{(y, y)}(y, y) \\ &= 0 \end{aligned}$$

So,

$$\begin{aligned} 0 &\leq (x - \lambda y, x - \lambda y) \\ &= (x - \lambda y, x) - \overline{\lambda}(x - \lambda y, y) \\ &= (x - \lambda y, x) \\ &= (x, x) - \lambda(y, x) \\ &= (x, x) - \frac{|(x, y)|^2}{(y, y)} \end{aligned}$$

So, $|(x, y)|^2 \leq (x, x)(y, y)$. \square

The theorem about the equivalence of all the norms in \mathbb{R}^n .

Theorem 3. Any two norms in \mathbb{R}^n are equivalent. If $\|\cdot\|_1$ and $\|\cdot\|_2$ are norms in \mathbb{R}^n then there exists $c_1, c_2 > 0$ such that

$$c_1\|x\|_1 \leq \|x\|_2 \leq c_2\|x\|_1$$

Proof. We may assume that $\|\cdot\|_1$ is the standard Euclidian norm in \mathbb{R}^n . We know that the unit sphere (w.r.t the standard norm) $S_1(0)$ is compact in \mathbb{R}^n since it is closed and bounded (Heine-Borel). We will now prove that the function $x \mapsto \|x\|_2$ is a continuous function. Indeed, consider the sequence $(x_n)_{n=1}^\infty$ which converges to x , we wish to show that $\|x_n\|_2$ converges to $\|x\|_2$.

$$||x_n| - |x|| \leq \|x_n - x\|_2 \rightarrow 0$$

So indeed this function is continuous. Therefore by EVT it attains its maximum and minimum on $S_1(0)$. Therefore

$$\begin{aligned} \|x\|_2 &= \|x\|_1 \times \left\| \frac{x}{\|x\|_1} \right\|_2 \\ &\leq \|x\|_1 \times \sup_{x \in S_1(0)} \|x\|_2 \end{aligned}$$

Analogously for the other direction, inf non-zero since $x \in S_1(0)$. \square

A functional (operator) is continuous at one point if and only if it is continuous everywhere.

Theorem 4. (Same for both) Let E_1, E_2 be normed vector spaces, if A , a linear operator between E_1 and E_2 is continuous at one point, then A is continuous everywhere.

Assume A is continuous at the point $x_0 \in E_1$. Given $x \neq x_0$ we wish to prove A is continuous at x . Take a sequence $(x_n)_{n=1}^\infty \subset E$ such that $\lim_{n \rightarrow \infty} x_n = x$. We will show $\lim_{n \rightarrow \infty} A(x_n) = A(x)$.

$$\begin{aligned} \lim_{n \rightarrow \infty} A(x_n + x_0 - x_0 + x - x) &= \lim_{n \rightarrow \infty} A((x_n - x) + x_0) + A(x - x_0) \\ &\rightarrow A(x_0) + A(x) - A(x_0) \\ &= A(x) \end{aligned}$$

So $A(x_n) \rightarrow A(x)$, so A is continuous.

A functional (operator) is continuous if and only if it is bounded.

Theorem 5. (Same for both) Let E be a normed vector space and l be a linear functional on E . l is continuous if and only if it is bounded.

Proof. \implies Assume l is bounded, we wish to show that l is continuous at one point, and thus everywhere. Consider a sequence $(x_n)_{n=1}^\infty \subset E$ such that $\lim_{n \rightarrow \infty} x_n = 0$.

$$|l(x_n)| \leq c \|x_n\| \rightarrow 0 \quad n \rightarrow \infty$$

So l is continuous at 0 and thus everywhere.

\Leftarrow For the sake of contradiction, assume l is continuous but unbounded. There exists a sequence $(x_n)_{n=1}^\infty \subset E$ such that

$$|l(x_n)| \geq n \|x_n\|$$

Define $\hat{x}_n = \frac{x_n}{n \|x_n\|}$, we have that $\|\hat{x}_n\| = \frac{1}{n} \rightarrow 0$ as $n \rightarrow \infty$. So by continuity, $l(\hat{x}_n)$ should go to 0. However,

$$|l(\hat{x}_n)| = \frac{|l(x_n)|}{n \|x_n\|} \geq 1$$

Here lies our contradiction. □

The theorem about the completeness of the dual space.

Theorem 6. Let E be a normed vector space, E' , it's dual, is complete.

Proof. Take a Cauchy sequence $(l_n)_{n=1}^\infty \subset E'$. Let us construct its limit l . For $x \in E$, we have

$$|l_n(x) - l_m(x)| \leq \|l_n - l_m\| \cdot \|x\| \rightarrow 0$$

As $n, m \rightarrow \infty$. So let us define,

$$l(x) = \lim_{n \rightarrow \infty} l_n(x)$$

We will now show linearity and boundedness of l .

$$\begin{aligned} l(\lambda x + \gamma y) &= \lim_{n \rightarrow \infty} (l_n(\lambda x + \gamma y)) \\ &= \lim_{n \rightarrow \infty} (\lambda l_n(x) + \gamma l_n(y)) \\ &= \lambda l(x) + \gamma l(y) \end{aligned}$$

$$\begin{aligned}
||l|| &= \sup_{||x|| \leq 1} ||l(x)|| \\
&= \sup_{||x|| \leq 1} ||\lim_{n \rightarrow \infty} l_n(x)|| \\
&\leq \sup_{||x|| \leq 1} \lim_{n \rightarrow \infty} ||l_n|| \cdot ||x|| \\
&\leq C \cdot \sup_{||x|| \leq 1} ||x|| \\
&\leq C
\end{aligned}$$

For some $C > 0$ as $(||l_n||)_{n=1}^\infty$ is Cauchy, it is bounded. It remains to show $||l_n - l|| \rightarrow 0$. Since $(l_n)_{n=1}^\infty$ is Cauchy, for large enough n and m ,

$$|l_n - l_m| < \varepsilon ||x||$$

Now pass the limit $m \rightarrow \infty$

$$\begin{aligned}
\lim_{m \rightarrow \infty} |l_n - l_m| &= |l_n - l| \\
&\leq \varepsilon ||x||
\end{aligned}$$

So finally,

$$||l_n - l|| \rightarrow 0$$

□

The Hahn–Banach theorem.

Theorem 7. Let E be a normed space and G a linear subset. For every $l \in G'$, there exists some $L \in E'$ such that $L|_G = l$ and $||L|| = ||l||$.

Lemma 1 (Dense subset). Let E be a normed space, $G \subset E$ a dense subset and $l \in G'$. Then there exists a unique $L \in E'$ such that $L|_G = l$ and $||L|| = ||l||$.

Proof. Take $x \in E$. By density there exists a sequence $(g_n)_{n=1}^\infty \subset G$ such that $\lim_{n \rightarrow \infty} g_n = x$. Define

$$L(x) = \lim_{n \rightarrow \infty} l(g_n)$$

Why is this well defined?

$$|l(g_n) - l(g_m)| \leq ||l|| \cdot ||g_n - g_m|| \rightarrow 0$$

as $n, m \rightarrow \infty$. Thus $(l(g_n))_{n=1}^\infty$ converges. Why is $L(x)$ independent of the sequence $(g_n)_{n=1}^\infty$. Indeed, consider $(g'_n)_{n=1}^\infty \subset G$ converging to x .

$$|l(g_n) - l(g'_n)| \leq ||l|| \cdot (||g_n - x|| + ||g'_n - x||) \rightarrow 0$$

as $n \rightarrow \infty$. $L|_G = l$ since for every $x \in G$, take $g_n = x$ for every $n \in \mathbb{N}$. Let us show that the norm is preserved.

$$\begin{aligned}
|L(x)| &= \left| \lim_{n \rightarrow \infty} l(g_n) \right| \\
&\leq \lim_{n \rightarrow \infty} ||l|| \cdot ||g_n|| \\
&= ||l|| \cdot \lim_{n \rightarrow \infty} ||g_n|| \\
&= ||l|| \cdot ||x||, \quad \text{by continuity of the norm}
\end{aligned}$$

Thus $||L|| \leq ||l||$ and so $||L|| = ||l||$. Now we will show uniqueness, consider $L_1|_G = L_2|_G = l$, by continuity,

$$L_1(x) = \lim_{n \rightarrow \infty} l(g_n) = L_2(x)$$

□

Lemma 2 (Extension of a functional by one dimension). Let E be a real normed space and $G \subset E$ a linear subspace. Choose $y \notin G$ and define $F = \text{span}(G \cup \{y\})$. For any linear bounded functional l on G . There is a bounded linear functional $L \in F'$ such that $L|_G = l$ and $\|L\| = \|l\|$.

Proof. Every $x \in F$ can be written as

$$x = g + \lambda y$$

where $g \in G$ and $\lambda \in \mathbb{R}$ in a unique way. Define $L : F \rightarrow \mathbb{R}$ by,

$$L(x) = l(y) + \lambda \cdot c$$

for some $c \in \mathbb{R}$. Let us now show that we can pick such a $c \in \mathbb{R}$ so $\|L\| = \|l\|$, it is sufficient to show that $\|L\| \leq \|l\|$, that is $\|L\|$ has not increased any larger than $\|l\|$.

The inequality $\|L\| \leq \|l\|$ is equivalent to

$$\begin{aligned} |L(g + \lambda y)| &= |l(y) + \lambda \cdot c| \\ &\leq \|l\| \cdot |g + \lambda y| \quad (*) \end{aligned}$$

Sufficient to prove this for $\lambda > 0$, since $g + \lambda y = -(g + (-\lambda)y)$. We can rewrite $(*)$ as follows,

$$-\|l\| \cdot |g + y| - l(y) \leq \lambda c \leq \|l\| \cdot |g + y| - l(y)$$

Now let $h = \lambda^{-1}g$ and divide the inequality by λ

$$-\|l\| \cdot |h + y| - l(h) \leq c \leq \|l\| \cdot |h + y| - l(h)$$

To find such a c , notice that for all $h_1, h_2 \in G$.

$$\begin{aligned} l(h_2) - l(h_1) &\leq \|l\| \cdot \|h_2 - h_1 + y - y\| \\ &\leq \|l\| \cdot \|(h_2 + y) + -(h_1 + y)\| \\ &\leq \|l\| \cdot |h_1 + y| + \|l\| \cdot |h_2 + y| \end{aligned}$$

Therefore,

$$-\|l\| \cdot |h_1 + y| - l(h_1) \leq \|l\| \cdot |h_2 + y| - l(h_2)$$

This implies that

$$\sup_{h_1 \in G} (-\|l\| \cdot |h_1 + y| - l(h_1)) \leq \inf_{h_2 \in G} (\|l\| \cdot |h_2 + y| - l(h_2))$$

Choosing c in between this sup and inf will obtain $(*)$. □

Lemma 3 (Real and separable). Let E be a real, linear, separable normed space and $G \subset E$ a linear subset. For every $l \in G'$ there exists an $L \in E'$ such that $L|_G = l$ and $\|L\| = \|l\|$.

Proof. Let

$$A = \{x_1, x_2, x_3, \dots\}$$

be a dense subset of E . Choose x_{n_1} to be the first element of A that does not lie in G . Define $G_1 = \text{span}(G \cup \{x_{n_1}\})$. Extend l to $l_1 : G_1 \rightarrow \mathbb{R}$ preserving the norm by Lemma 2. Repeating this procedure we obtain the subspaces,

$$G \subset G_1 \subset G_2 \subset \dots$$

and similarly, functionals l, l_1, l_2, \dots all with the same norm.

Define

$$M = \bigcup_{i=1}^{\infty} G_n$$

This is a linear set and is dense in E since it contains all of A . Now define $L_0 : M \rightarrow \mathbb{R}$ as follows, if $x \in M$, then $x \in G_n$ for some $n \in \mathbb{N}$, so define

$$L_0(x) = l_n(x)$$

This clearly also preserves the norm since $\|l\| = \|l_n\|$ for every $n \in \mathbb{N}$. By continuity we can extend L_0 to all of E . \square

Lemma 4 (Complex and separable). Let E be a complex, linear, separable normed space and $G \subset E$ a linear subset. For every $l \in G'$ there exists an $L \in E'$ such that $L|_G = l$ and $\|L\| = \|l\|$.

Proof. We can consider E as a vector space over the reals, denote it $E_{\mathbb{R}}$. Take $k \in E'$ and let $m = \Re(k)$ and $n = \Im(k)$, both of these lie in $E_{\mathbb{R}}$. Let us show they are bounded

$$|m(x)| \leq |m(x) + in(x)| = |k(x)| \leq \|k\| \cdot \|x\|$$

Similar for $|n(x)|$. Thus $\|m\|, \|n\| \leq \|k\|$. Observe

$$\begin{aligned} m(ix) + in(ix) &= k(ix) \\ &= ik(x) \\ &= i(m(x) + in(x)) \\ &= -n(x) + im(x) \end{aligned}$$

So $m(ix) = -n(x)$ and $n(ix) = m(x)$.

We have that l is defined on $G \subset E$. We can extend l from $G_{\mathbb{R}}$ to $E_{\mathbb{R}}$ and moreover $\Re(l) = m \in G'_{\mathbb{R}}$ can be extended to $M \in E'_{\mathbb{R}}$ by Lemma 3.

Define

$$L(x) = M(x) - iM(ix)$$

Clearly $L|_G = l$ and L is linear, indeed,

$$\begin{aligned} L((\alpha + i\beta)x) &= \alpha L(x) + \beta L(ix) \\ &= \alpha L(x) + \beta(M(ix) - iM(ix)) \\ &= \alpha L(x) + \beta i(M(x) - iM(-x)) \end{aligned}$$

Now we will show $\|L\| = \|l\|$, indeed take θ such that

$$L(x) = |L(x)|e^{i\theta}$$

Now,

$$\begin{aligned} |L(x)| &= L(x)e^{-i\theta} \\ &= L(xe^{-i\theta}) \\ &= M(xe^{-i\theta}) \\ &\leq |M(xe^{-i\theta})| \\ &\leq \|M\| \cdot \|xe^{-i\theta}\| \\ &= \|M\| \cdot |e^{-i\theta}| \cdot \|x\| \\ &= \|M\| \cdot \|x\| \\ &\leq \|l\| \cdot \|x\| \end{aligned}$$

So $\|L\| \leq \|l\|$. \square

Proof. (The field over E is \mathbb{R}) Let l_P be an extension of l to the subspace $P \subset E$ containing G . At least one such an extension exists by Lemma 2.

Let X be the set of all such extensions which preserve the norm. Define $l_p \leq l_q$ if $P \subset Q$ and $l_p = l_q|_P$. This is a partial ordering on X .

Let $Y = \{l_{P_\alpha} \mid \alpha \in A\}$ be a chain in X . This chain has an upper bound, set

$$\tilde{P} = \bigcup_{\alpha \in A} P_\alpha$$

The define \tilde{l} on \tilde{P} by,

$$\tilde{l}(x) = l_\alpha(x)$$

if $x \in P_\alpha$. Clearly well defined and $||\tilde{l}|| = ||l||$. We can then extend \tilde{l} to the closure of \tilde{P} by continuity, then \tilde{l} is an upper bound for Y .

By Zorn's Lemma, X has a maximal element, denoted L . This is defined on all of E since if it weren't we could extend it and it wouldn't be maximal. By construction \tilde{l} is a norm-preserving extension of l .

If E is complex, same argument as used in the separable case. □

The theorem about a monomorphism between a normed space and its second dual.

Theorem 8. The map $\Phi : E \rightarrow E''$ given by $x \mapsto \phi_x$ is an isometric mono-morphism.

Proof. Let us first show linearity,

$$\begin{aligned} \Phi(\alpha x + \beta y)(l) &= \phi_{\alpha x + \beta y}(l) \\ &= l(\alpha x + \beta y) \\ &= \alpha l(x) + \beta l(y) \\ &= \alpha \phi_x(l) + \beta \phi_y(l) \\ &= \alpha \Phi(x)(l) + \beta \Phi(y)(l) \end{aligned}$$

Now we will show the isomorphism. Take $x \neq y$, we will show that $\phi_{x-y} \neq 0$. Indeed, by Hahn-Banach, there exists some $l \in E'$ such that $l(x - y) = ||x - y|| \neq 0$. Then,

$$\phi_{x-y}(l) = l(x - y) \neq 0$$

Now we will show the norm is preserved. Indeed

$$\begin{aligned} |\phi_x(l)| &= |l(x)| \\ &\leq ||l|| \cdot ||x|| \end{aligned}$$

So $||\phi_x|| \leq ||x||$ Now we will show $||x|| \leq ||\phi_x||$. If $x = 0$ this is trivial. Take $x \neq 0$. By Hahn Banach, there exists some $l \in E'$ such that $||l|| = 1$ and $l(x) = ||x||$.

$$\begin{aligned} ||x|| &= l(x) \\ &= \phi_x(l) \\ &\leq ||\phi_x|| \cdot ||l|| \\ &= ||\phi_x|| \end{aligned}$$

□

The Banach–Steinhaus theorem.

Theorem 9. Let E be a Banach space. Consider a sequence $(l_n)_{n=1}^\infty \subset E'$. Assume that for every $x \in E$, there exists $c_x > 0$ such that

$$||l_n(x)|| \leq c_x$$

for all $n \in \mathbb{N}$. The conclusion is that there exists $c > 0$ such that

$$||l_n|| \leq c$$

for all $n \in \mathbb{N}$.

Lemma 1 (Nested Ball Lemma). Consider a sequence of balls,

$$\overline{B_{r_1}(x_1)} \supset \overline{B_{r_1}(x_2)} \supset \overline{B_{r_3}(x_3)} \supset \cdots$$

Such that $r_i \rightarrow 0$ as $i \rightarrow \infty$. Then

$$\bigcap_{i=1}^{\infty} \overline{B_{r_i}(x_i)} \neq \emptyset$$

Proof. To begin $(x_i)_{i=1}^\infty$ is a Cauchy sequence. Denote

$$x^* = \lim_{i \rightarrow \infty} x_i$$

Given $m \in \mathbb{N}$, $x_k \in \overline{B_{r_m}(x_m)}$ for all $k \geq m$. By closedness we know the limit point $x^* \in \overline{B_{r_m}(x_m)}$. Thus x^* is in all of the balls and the intersection is non-empty. \square

Lemma 2 (Baire Category Theorem). Let $(C_n) \subset E$ be a sequence of closed sets such that

$$\bigcup_{n=1}^{\infty} C_n = E$$

Then there exists $n \in \mathbb{N}$, $x \in E$ and $r > 0$ such that $B_r(x) \subset C_n$.

Proof. Assume the contrary, choose $x \in E \setminus C_1$. Since C_1 is closed, the complement must be open. Thus there must exist some $r_1 > 0$ such that $\overline{B_{r_1}(x)} \subset E \setminus C_1$. Now by our assumption, $B_{r_1}(x) \cap (E \setminus C_2) \neq \emptyset$. Take $x_2 \in B_{r_1}(x) \cap (E \setminus C_2)$. There exists some $r_2 < \frac{r_1}{2}$ such that $\overline{B_{r_2}(x_2)} \subset B_{r_1}(x) \cap (E \setminus C_2)$.

Now there exists some $x_3 \in B_{r_2}(x_2) \cap (E \setminus C_3)$. Choose $r_3 < \frac{r_2}{2}$ such that $\overline{B_{r_3}(x_3)} \subset B_{r_2}(x_2) \cap (E \setminus C_3)$. Continuing this procedure we obtain,

$$\overline{B_{r_1}(x_1)} \supset \overline{B_{r_2}(x_2)} \supset \cdots$$

With $r_n \rightarrow 0$ as $n \rightarrow \infty$. Clearly

$$\bigcap_{n=1}^{\infty} \overline{B_{r_n}(x_n)} \supset \{x\}$$

for some $x \in E$. By construction $x \in E \setminus C_n$ for all $n \in \mathbb{N}$. This means that

$$\bigcup_{n=1}^{\infty} C_n \neq E$$

Contradiction. \square

Lemma 3. There exists $x \in E$ and $r > 0$ such that the set

$$S_{r,a} = \{l_n(x) \mid x \in \overline{B_r(a)}, n \in \mathbb{N}\}$$

is bounded.

Proof. Denote

$$C_m = \{x \in E \mid |l_n(x)| \leq m, \forall n \in \mathbb{N}\}$$

We have that C_m is closed. Indeed if $(x_k)_{k=1}^\infty \subset C_m$ converges to some $x \in E$. Then for every $n \in \mathbb{N}$

$$|l_n(x)| = \left| \lim_{k \rightarrow \infty} l_n(x_k) \right| \leq m$$

By assumption of the theorem,

$$E = \bigcup_{m=1}^{\infty} C_m$$

Thus, by Lemma 2, at least one C_m contains a ball. □

Proof. Recall that

$$\|l_n\| = \sup_{x \in B_1(0)} |l_n(x)|$$

It suffices to show that $|l_n(x)| \leq c$ for all $x \in \overline{B_1(0)}$ and $n \in \mathbb{N}$.

Given $x \in \overline{B_1(0)}$, define $x' = rx + a$ where r and a are from Lemma 3. Thus $x = \frac{1}{r}(x' - a)$ and

$$\begin{aligned} |l_n(x)| &= \left| l_n\left(\frac{1}{r}(x' - a)\right) \right| \\ &\leq \frac{1}{r}(|l_n(x')| + |l_n(a)|) \end{aligned}$$

By Lemma 3 the two summands are bounded by some $c' > 0$ and so

$$|l_n(x)| \leq \frac{2c'}{r}$$

and is bounded on $B_1(0)$ uniformly in n . □

The theorem about the weak compactness of the unit sphere in the dual space.

Theorem 10. If E is separable then the unit sphere in E' is weakly compact, i.e. $(l_n)_{n=1}^\infty \subset S^1(0)$ has a weakly convergent subsequence.

Theorem "Previous Theorem". Let E be a Banach space. Assume that M is dense in E . The sequence $(l_n)_{n=1}^\infty \subset E'$ converges weakly to $l \in E'$ iff

- a) $l_n(x) \rightarrow l(x)$ on M .
- (b) $\|l_n\| \leq c$ for some c independent of M

Proof. Let $M = \{x_1, x_2, x_3, \dots\}$ be dense in E . Then,

$$|l_n(x_1)| \leq \|l\| \cdot \|x_1\| = \|x_1\|$$

So there exists a subsequence $(l_{k_1})_{k=1}^\infty$ such that

$$(l_{k_1}(x_1))_{k=1}^\infty$$

converges. By the same token, there is a subsequence $(l_{k_2})_{k=1}^\infty \subset (l_{k_1})_{k=1}^\infty$ such that

$$(l_{k_2}(x_2))_{k=1}^\infty$$

converges. Continuing in this manner to obtain a subsequence $(l_{k_q})_k = 1$ such that

$$(l_{k_q}(x_q))_{k=1}^\infty$$

converges for every q . The sequence,

$$(l_{k_k}(x_k))_{k=1}^\infty$$

converges for every $k \in \mathbb{N}$. Define

$$l(x_i) = \lim_{k \rightarrow \infty} l_{k_k}(x_i)$$

for $i \in \mathbb{N}$. This is quite clearly linear, further more,

$$\|l\| \leq \sup \|l_{k_k}\| = 1$$

We can then extend l to all of E by continuity. The previous theorem implies that $l_{k_k} \rightarrow l$ weakly. \square

The theorem about the existence and uniqueness of the orthogonal projection in a Hilbert space.

Theorem 11. Let $G \subset H$ be a subspace. Given $x \in H$ there exists unique $y \in G$ such that $(x - y) \perp G$, i.e. $(x - y) \in G^\perp$.

Proof. If $x \in G$ then $y = x$. Assume $x \notin G$. Then let

$$d := \text{dist}(x, G) = \inf_{y \in G} \|x - y\| > 0$$

Choose $(y_n)_{n=1}^\infty \subset G$ s.t $\|x - y_n\| \rightarrow d$ as $n \rightarrow \infty$. We will show that $(y_n)_{n=1}^\infty$ is a Cauchy sequence. Take $h \in G$ such that $h \neq 0$ and $\eta \in \mathbb{C}$. Then,

$$\begin{aligned} d^2 &\leq \|x - (y_n + \eta h)\|^2 \\ &= (x - (y_n + \eta h), x - (y_n + \eta h)) \\ &= \|x - y_n\|^2 + |\eta|^2 \|h\|^2 - \bar{\eta}(x - y_n, h) - \eta(h, x - y_n) \end{aligned}$$

Now let

$$\eta = \|h\|^{-2}(x - y_n, h)$$

This simplifies the above to

$$d^2 \leq \|x - y_n\|^2 - \frac{|(x - y_n, h)|^2}{\|h\|^2}$$

That is,

$$\|h\|^2 \|x - y_n\|^2 - |(x - y_n, h)|^2 - d^2 \|h\|^2 \geq 0$$

or

$$|(x - y_n, h)|^2 \leq \|h\|^2 (d_n^2 - d^2) \quad (*)$$

Where $d_n = \|x - y_n\|$. Thus,

$$|(x - y_n, h)| \leq \|h\| \sqrt{d_n^2 - d^2}$$

This holds for $h \in G$. Set $h = y_n - y_m$, for $n, m \in \mathbb{N}$. Then we have

$$\begin{aligned} |(y_n - y_m \pm x, h)| &\leq |(x - y_n, y)| + |(x - y_m, h)| \\ &\leq \|h\| (\sqrt{d_n^2 - d} + \sqrt{d_m^2 - d}) \end{aligned}$$

So

$$\|y_n - y_m\|^2 \leq \|y_n - y_m\| (\sqrt{d_n^2 - d} + \sqrt{d_m^2 - d}) \rightarrow 0$$

as $n, m \rightarrow \infty$. Thus $(y_n)_{n=1}^\infty$ is Cauchy. Denote $y = \lim_{n \rightarrow \infty} y_n$, certainly $y \in G$. Passing the limit through to $(*)$ we find that $(x - y, h) = 0$ for all $h \in G$. Thus $y = \text{proj}_G(x)$. Why is y unique? If $(x - y) \perp G$ and $(x - y') \perp G$ then

$$(y - y' \pm x, h) = (y - x, h) + (x - y', h) = 0$$

for all $h \in G$ so set $h = y - y'$, we find

$$\|y - y'\|^2 = 0$$

So $y = y'$. \square

The Riesz representation theorem.

Theorem 12. Let H be a Hilbert space, given $y \in H$ the formula

$$x \mapsto (x, y) \quad (*)$$

defines a functional in H' . Conversely for every $l \in H'$, there exists some $y \in H$ such that

$$l(x) = (x, y)$$

for all $x \in H$. Moreover $\|l\| = \|y\|$.

Proof. $(*)$, we will denote this as l from here on is clearly linear, why is it bounded? By Cauchy-Schwarz we have

$$|(x, y)| \leq \|x\| \cdot \|y\|$$

so

$$\|l\| \leq \|y\|$$

Indeed, this is an equality, since $|(y, y)| = \|y\|^2$.

Conversely, take $l \in H'$. Denote $G = \ker(l)$. G is a subspace and $H = G \oplus G^\perp$. Moreover $\dim(G^\perp)$ has dimension 1. Fix $c \in G^\perp$ with $\|c\| = 1$. For every $x \in H$ we have $x = g + \lambda c$ where $g \in \ker(l)$ and $\lambda \in \mathbb{C}$. Thus,

$$\begin{aligned} l(x) &= l(g + \lambda c) \\ &= l(g) + \lambda l(c) \\ &= \lambda l(c) \end{aligned}$$

and

$$\begin{aligned} (x, c) &= (g + \lambda c, c) \\ &= (g, c) + \lambda (c, c) \\ &= \lambda \end{aligned}$$

We conclude that

$$l(x) = (x, c)l(c) = (x, \overline{l(c)}c)$$

Now take $y = \overline{l(c)}c$ to get $(*)$. Now why is y unique? If $l(x) = (x, y) = (x, y')$. Then $(x, y - y') = l(x) - l(x) = 0$ for all $x \in H$ so choose $x = y - y'$ and $\|y - y'\|^2 = 0$ so $y = y'$. \square

The theorem about the invertibility of $\mathbb{1} - A$ when $\|A\|$ is small.

Theorem 13. Let E be a Banach Space. Let $A \in \mathcal{L}(E)$, satisfy $\|A\| = q < 1$. Then $\mathbb{1} - A$ is invertible and

$$(\mathbb{1} - A)^{-1} = \mathbb{1} + A + A^2 + A^3 + \dots$$

Proof. Let

$$S_n = \mathbb{1} + A + A^2 + \dots + A^n$$

We claim that S_n is Cauchy, we have

$$\begin{aligned} \|S_{n+p} - S_n\| &= \|A^{n+1}\| + \|A^{n+2}\| + \dots + \|A^{n+p}\| \\ &\leq q^{n+1} + \dots + q^{n+p} \\ &\leq q^n \frac{1}{1 - q} \\ &\rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$. Since E is Banach so is $\mathcal{L}(E)$, so $(S_n)_{n=1}^\infty$ converges to S in norm. We need to show $S(\mathbb{1} - A) = (\mathbb{1} - A)S = \mathbb{1}$. We will first show that $(\mathbb{1} - A)S_n \rightarrow (\mathbb{1} - A)S$. Indeed

$$\|(\mathbb{1} - A)S_n - (\mathbb{1} - A)S\| \leq \|(\mathbb{1} - A)\| \cdot \|S_n - S\| \rightarrow 0$$

It suffices to show that $(\mathbb{1} - A)S_n \rightarrow \mathbb{1}$. We have that

$$\begin{aligned} (\mathbb{1} - A)S_n &= S_n(\mathbb{1} - A) \\ &= \mathbb{1} + A + \cdots + A^n - A - A^2 - \cdots - A^n - A^{n+1} \\ &= \mathbb{1} - A^{n+1} \end{aligned}$$

So

$$\begin{aligned} \|(\mathbb{1} - A)S_n - \mathbb{1}\| &= \|S_n(\mathbb{1} - A) - \mathbb{1}\| \\ &= \|\mathbb{1} - A^{n+1} - \mathbb{1}\| \\ &= \|A^{n+1}\| \\ &\leq \|A\|^{n+1} \\ &\rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$. □

The open mapping theorem and the bounded inverse theorem.

Theorem 14.1 (Open Mapping Theorem). Let E_1 and E_2 be Banach spaces. The operator $A \in \mathcal{L}(E_1, E_2)$ is open iff it is surjective.

Lemma 1. There exists some $y_0 \in E_1$, $r_0 > 0$ and $\alpha \in (0, 1)$ such that

$$B_{\alpha r_0}(Ay_0) \subset \overline{A(B_{r_0}(y_0))}$$

Proof. The sets $\overline{B_n(0)}$ are closed and their union over $n \in \mathbb{N}$ is E_2 . By the Baire Category Theorem. At least one of these sets contains a ball, i.e. there exists $r > 0, n \in \mathbb{N}$ and $x \in E_2$ s.t.

$$B_r(x) \subset \overline{A(B_n(0))}$$

Choose $y_0 \in E_1$ such that $x = Ay_0$ and choose $r_0 > 0$ such that $B_{r_0}(y_0) \supset B_n(0)$. Thus,

$$B_r(Ay_0) \subset \overline{A(B_n(0))} \subset \overline{A(B_{r_0}(y_0))}$$

It suffices to set $\alpha = \frac{r}{r_0}$. □

Lemma 2. For every point $y \in E_1$ and $r > 0$

$$B_{\alpha r}(Ay) \subset \overline{A(B_r(y))}$$

Proof. Take $z \in B_{\alpha r}(Ay)$ then,

$$A(y_0) + \frac{r_0}{r}(z - Ay) \in B_{\alpha r_0}(Ay_0)$$

So,

$$A(y_0) + \frac{r_0}{r}(z - Ay) \in \overline{A(B_{r_0}(y_0))}$$

By Lemma 1. Hence there exists some sequence $(z_n)_{n=1}^\infty \subset B_{r_0}(y_0)$ such that

$$Ay_0 + \frac{r_0}{r}(z - Ay) - Az_n \rightarrow 0$$

Rewriting this we get,

$$z - A(y + \frac{r}{r_0}(z_n - y_0)) \rightarrow 0$$

and we have

$$A(y + \frac{r}{r_0}(z_n - y_0)) \in B_r(y)$$

So $z \in \overline{A(B_r(y))}$

□

Lemma 3. For every $y \in E_1$ and every $r > 0$

$$B_{\frac{\alpha r}{2}}(Ay) \subset A(B_r(y))$$

Proof. Take $z \in B_{\frac{\alpha}{2}r}(Ay)$ and $\varepsilon < 1$. By Lemma 2 there exists some $y_1 \in B_{\frac{r}{2}}(y)$ such that

$$\|z - Ay_1\| < \alpha\varepsilon$$

so, $z \in B_{\alpha\varepsilon}(Ay_1) \subset \overline{A(B_\varepsilon(y_1))}$. Again there is a $y_2 \in B_\varepsilon(y_1)$ such that

$$\|z - Ay_2\| < \alpha\varepsilon^2$$

, then $z \in B_{\alpha\varepsilon^2}(Ay_2) \subset \overline{A(B_{\varepsilon^2}(y_2))}$. Continue this process to obtain a sequence $(y_n)_{n=1}^\infty$ such that

$$\|z - Ay_n\| < \alpha\varepsilon^n$$

and

$$\|y_n - y_{n+1}\| < \varepsilon^n$$

So $(y_n)_{n=1}^\infty$ converges to some $y^* \in E_1$ and $Ay^* = z$. Now we will show that y^* is indeed in $B_r(y)$.

$$\begin{aligned} \|y - y^*\| &\leq \|y \pm y_n \pm y_{n-1} \pm \cdots - y^*\| \\ &\leq \|y - y_n\| + \sum_{k=2}^n \|y_k - y_{k-1}\| + \|y_1 - y^*\| \\ &\leq 0 + \sum_{k=2}^n \varepsilon^k + \frac{r}{2} \\ &\leq \frac{\varepsilon}{1-\varepsilon} + \frac{r}{2} \end{aligned}$$

We can choose ε small enough such that this is less than r .

□

Proof. \implies

Assume A is an open mapping, we will show A is surjective. Indeed take $B_1(0) \subset E_1$, we know that $A(B_1(0))$ is an open set in E_2 so there exists $\varepsilon > 0$ such that

$$B_\varepsilon(0) \subset A(B_1(0))$$

We can write every $y \in E_2$ as an element of this ball by setting $\tilde{y} = \frac{\varepsilon}{2\|y\|}y$. Thus there exists $x \in B_1(0)$ such that

$$\begin{aligned} A(x) &= \tilde{y} \\ A(x) &= \frac{\varepsilon}{2\|y\|}y \\ A\left(\frac{2\|y\|}{\varepsilon}x\right) &= y \end{aligned}$$

So we have that A is surjective.

\Leftarrow

Assume that A is surjective. Take an open set $V \subset E_1$, we need to show $A(V)$ is open. Take $y \in A(V)$, let x be such that $Ax = y$. Since V is open, there exists some $r > 0$ such that

$$B_r(x) \subset V$$

By Lemma 3 we have

$$B_{\frac{r}{2}}(Ax) \subset A(B_r(x)) \subset AV$$

This holds for all y and x satisfying the above, thus AV is open. \square

Theorem 14.2 (Bounded Inverse Theorem). Let $A \in \mathcal{L}(E_1, E_2)$ be a bijection. Then A is invertible. That is, $A^{-1} : E_2 \rightarrow E_1$.

Proof. For A^{-1} the pre-image of an open set is open. \square

The closed graph theorem.

Theorem 15. If E_1 and E_2 are Banach spaces, then $A : E_1 \rightarrow E_2$ is a linear operator and $\Gamma(A)$ is closed. Then A is bounded.

Proof. Observe that $\Gamma(A)$ is a Banach space with the norm

$$\|(x, Ax)\|_{\Gamma(A)} = \|x\|_{E_1} + \|Ax\|_{E_2}$$

Now define $\pi_1 : \Gamma(A) \rightarrow E_1$ and $\pi_2 : \Gamma(A) \rightarrow E_2$ by

$$\pi_1((x, Ax)) = x$$

and

$$\pi_2((x, Ax)) = Ax$$

Clearly π_1 and π_2 are bounded. Moreover π_1 is invertible by the Bounded Inverse Theorem. This implies that

$$A = \pi_2 \pi_1^{-1}$$

is bounded. \square

The theorem about the decomposition of an operator into its real and imaginary parts.

Theorem 16. Let H be a Hilbert space. For every operator $A \in \mathcal{L}(H)$ there exists $\Re(A), \Im(A)$ both in $\mathcal{L}(H)$ such that they are self adjoint and $A = \Re(A) + \Im(A)$.

Proof. Let $\Re(A) = \frac{A+A^*}{2}$ and $\Im(A) = \frac{A-A^*}{2i}$.

$$\begin{aligned} (\Re(A))^* &= \frac{A^* + (A^*)^*}{2} = \frac{A^* + A}{2} = \Re(A) \\ (\Im(A))^* &= \frac{A^* - (A^*)^*}{-2i} = \frac{A - A^*}{2i} = \Im(A) \end{aligned}$$

\square

The Hellinger-Toeplitz Theorem

Theorem 17. Let H be a Hilbert space. Assume $A : H \rightarrow H$ is a linear and $(Ax, y) = (x, Ay)$ for all $x, y \in H$. Then A is bounded, i.e. $A \in \mathcal{L}(H)$.

Proof. Let us prove that $\Gamma(A)$ is closed. Take a sequence (x_n, Ax_n) converging to $(x, y) \in H \times H$. We wish to show that $y = Ax$. Given $z \in H$, we can see that

$$\begin{aligned}(z, y) &= \lim_{n \rightarrow \infty} (z, Ax_n) \\ &= \lim_{n \rightarrow \infty} (Az, x_n) \\ &= (Az, y) \\ &= (z, Ay)\end{aligned}$$

Thus $(z, y - Ax) = 0$ for all $z \in H$, take $z = y - Ax$ and $\|y - Ax\|^2 = 0$ so $y = Ax$. □

Every sesquilinear form is generated by an operator.

Theorem 18. Let H be a Hilbert space. Assume b is a sesquilinear form on H such that

$$|b(x, y)| \leq c\|x\| \cdot \|y\|$$

for some $c > 0$. Then there exists some $A \in \mathcal{L}(H)$ such that $b(x, y) = (Ax, y)$.

Proof. Fix $x \in H$ and set $l(y) = \overline{b(x, y)}$. This is a bounded linear functional on H .

By Riesz Representation Theorem, there exists $a_x \in H$ such that

$$\overline{b(x, y)} = l(y) = (y, a_x)$$

Now set $Ax = a_x$. We will show that A is linear and bounded.

$$\begin{aligned}(A(\alpha x_1 + \beta x_2), y) &= b(\alpha x_1 + \beta x_2, y) \\ &= b(\alpha x_1, y) + b(\beta x_2, y) \\ &= \alpha b(x_1, y) + \beta b(x_2, y) \\ &= \alpha (Ax_1, y) + \beta (Ax_2, y)\end{aligned}$$

$$\|Ax\|^2 = b(x, Ax) \leq c\|x\| \cdot \|Ax\|$$

So $\|Ax\| \leq c\|x\|$ □

Every self adjoint operator that squares itself is a projector

Theorem 19. Let $A \in \mathcal{L}(H)$ be a self-adjoint operator with $A^2 = A$. Then $A = P_G$ for some $G \subset H$.

Proof. Define $G = \ker(A - \mathbb{1})$. This is certainly a subspace of H . Since $A^2 = A$ for all $x \in H$ we have $Ax \in G$. Given $g \in G$,

$$\begin{aligned}(Ax, g) &= (x, Ag) \\ &= (x, g) \\ &= (x, P_G g) \\ &= (P_G x, g)\end{aligned}$$

So we have $(Ax - P_G x, g) = 0$ for all $g \in G$. Let $g = Ax - P_G x$ and we arrive at $\|Ax - P_G x\|^2 = 0$ so $Ax = P_G x$. □

The theorem about the inverse of a unitary operator

Theorem 20. Let H be a Hilbert space and $A \in \mathcal{L}(H)$ a unitary operator. Then $A^* = A^{-1}$.

Proof. We know that A is invertible since

$$\|Ax\| = \|x\|$$

since A is unitary. Now given $x, y \in H$

$$\begin{aligned} (x, A^*y) &= (Ax, y) \\ &= (Ax, AA^{-1}y) \\ &= (x, A^{-1}y) \end{aligned}$$

So $A^*y = A^{-1}y$ for all $y \in H$. □

The Hilbert-Schmidt norm is basis independent

Theorem 21. If $(e_i)_{i=1}^\infty$ and $(f_i)_{i=1}^\infty$ are both orthonormal bases of H , then

$$\sum_{i=1}^\infty \|Ae_i\|^2 = \sum_{i=1}^\infty \|Af_i\|^2$$

Proof.

$$\begin{aligned} \sum_{i=1}^\infty \|Ae_i\|^2 &= \sum_{i=1, j=1}^\infty |(Ae_i, f_j)|^2 \\ &= \sum_{i=1, j=1}^\infty |(e_i, A^*f_j)|^2 \\ &= \sum_{i=1, j=1}^\infty |(A^*f_j, e_i)|^2 \\ &= \sum_{i=1}^\infty \|A^*f_i\|^2 \\ &= \sum_{k=1, j=1}^\infty |(A^*f_j, f_k)|^2 \\ &= \sum_{k=1, j=1}^\infty |(Af_k, f_j)|^2 \\ &= \sum_{k=1}^\infty \|Af_k\|^2 \end{aligned}$$

□

The theorem about the closedness and boundedness of the spectrum.

Theorem 22.1. Let E be a normed space and consider an operator $A \in \mathcal{L}(E)$. Then

$$\text{spec}(A) \subset \overline{B_{\|A\|}(0)}$$

Proof. Assume that $|z| > \|A\|$, we will show that z is a regular point of A . That is, $(A - z\mathbb{1})$ is invertible. Indeed, $A - z\mathbb{1} = -z(\mathbb{1} - \frac{A}{z})$ and $\frac{\|A\|}{|z|} < 1$ so $(A - z\mathbb{1})$ is invertible and z is a regular point of A . □

Theorem 22.2. The spectrum of an operator A is closed.

Proof. Let z_0 be a regular point of A , if $z \in \mathbb{C}$ then

$$\begin{aligned} A - z\mathbb{1} &= A + z_0\mathbb{1} - z_0\mathbb{1} - z\mathbb{1} \\ &= A - z_0\mathbb{1} + (z_0 - z)\mathbb{1} \end{aligned}$$

Then choose z small enough so that $\|(z_0 - z)\mathbb{1}\| < \|(A - z_0\mathbb{1})\|^{-1}$. Then the sum is invertible. \square

The spectrum is non-empty.

Theorem 23. Let A be an operator, $\text{spec}(A) \neq \emptyset$.

Lemma 1. The function $z \mapsto \|R_z\|$ is bounded on $\overline{B_{2\|A\|}(0)}$. Moreover,

$$\lim_{|z| \rightarrow \infty} \|R_z\| \rightarrow 0$$

Proof. Outside of $\overline{B_{2\|A\|}(0)}$,

$$\begin{aligned} \|R_z\| &= \|(-z)^{-1}(\mathbb{1} - A\frac{1}{z})^{-1}\| \\ &= |z|^{-1} \|(\mathbb{1} - A\frac{1}{z})\|^{-1} \\ &\leq \frac{1}{|z|} \cdot \frac{1}{1 - \frac{\|A\|}{|z|}} \\ &\leq \frac{1}{z} \cdot \frac{1}{1 - \frac{\|A\|}{2\|A\|}} \\ &= \frac{2}{z} \\ &\leq \frac{1}{\|A\|} \end{aligned}$$

\square

Lemma 2. Fix $\alpha \in E$ and $l \in E'$. Define $f_{x,l} : \mathbb{C} \setminus \text{spec}(A) \rightarrow \mathbb{C}$ by

$$f_{x,l}(z) = l(R_z(x))$$

Then $f_{x,l}$ is analytic.

Proof. For $z \in \mathbb{C} \setminus \text{spec}(A)$,

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{1}{h} (f_{x,l}(z+h) - f_{x,l}(z)) &= \lim_{h \rightarrow 0} \frac{1}{h} l((R_{z+h} - R_z)(x)) \\ &= l(\lim_{h \rightarrow 0} \frac{1}{h} (R_{z+h} - R_z)(x)) \\ &= l(\lim_{h \rightarrow 0} \frac{1}{h} (z+h-z)(R_{z+h}R_z)(x)) \\ &= l(R_z^2(x)) \end{aligned}$$

\square

Proof. Assume $\text{spec}(A) = \emptyset$. Then $f_{x,l}$ is analytic on \mathbb{C} . Moreover, it is bounded. Indeed $\|R_z\|$ is bounded on $\mathbb{C} \setminus \overline{B_{2\|A\|}(0)}$ by Lemma 1 and on $\overline{B_{2\|A\|}(0)}$ by continuity. Thus

$$\sup_{z \in \mathbb{C}} \|R_z\| = d < \infty$$

Therefore,

$$|f_{x,l}(z)| = |l(R_z(x))| \leq \|l\| \cdot \|x\| \cdot d$$

By Liouville's Theorem, $f_{x,l}$ is constant. Since $\lim_{|z| \rightarrow \infty} \|R_z\| = 0$, $f_{x,l}$ is identically 0. This means

$$l(R_z x) = 0$$

for all l and so $R_z x = 0$ for every x and so $R_z = 0$. This is impossible since R_z is invertible, here lies our contradiction. \square