

HW 11 - MATH411

Danesh Sivakumar

April 16, 2022

Problem 1 (Exercise 5, Page 481)

Let \mathbf{I} be a generalized rectangle in \mathbb{R}^n and suppose that the function $f: \mathbf{I} \rightarrow \mathbb{R}$ is integrable. Assume that $f(\mathbf{x}) \geq 0$ if \mathbf{x} is a point in \mathbf{I} with a rational component. Prove that $\int_{\mathbf{I}} f \geq 0$.

Proof. Because f is integrable, there exists an Archimedean sequence of partitions \mathbf{P}_k of \mathbf{I} . Because the rationals are dense in \mathbb{R} , it follows that for each rectangle \mathbf{J} , $M(f, \mathbf{J}) \geq 0$. Thus, it follows that the upper Darboux sum $U(f, \mathbf{P}_k) \geq 0$. The fact that the function is integrable means that we can take the limit as k tends to ∞ , resulting in

$$\int_{\mathbf{I}} f = \lim_{k \rightarrow \infty} U(f, \mathbf{P}_k) \geq 0$$

□

Problem 2 (Exercise 10, Page 482)

Let \mathbf{I} be a generalized rectangle in \mathbb{R}^2 and suppose that the bounded function $f: \mathbf{I} \rightarrow \mathbb{R}$ has the value 0 on the interior of \mathbf{I} . Show that $f: \mathbf{I} \rightarrow \mathbb{R}$ is integrable and that $\int_{\mathbf{I}} f = 0$. Is the same result true for a generalized rectangle \mathbf{I} in \mathbb{R}^n ?

Proof. We prove the general case. Take the regular partition \mathbf{P}_k of the generalized rectangle into k intervals in each component. Because f is bounded, take $M > 0$ such that $|f| < M$. Observe that all rectangles except two in each dimension will not share an edge with the boundary of \mathbf{I} , meaning that there are $(k-2)^n$ rectangles that do not share any edge with the boundary. Thus, the number of rectangles that have nonzero values is equal to $k^n - (k-2)^n$. Thus it follows that

$$0 \leq |U(f, \mathbf{P}_k) - L(f, \mathbf{P}_k)| \leq \sum_{\mathbf{J} \in \mathbf{P}_k} |M_{\mathbf{J}} - m_{\mathbf{J}}| \text{vol} \mathbf{J} \leq \frac{2M(k^n - (k-2)^n)}{k^n}$$
$$\frac{2M(k^n - (k-2)^n)}{k^n} = 2M \left[1 - \left(1 - \frac{2}{k}\right)^n \right] \rightarrow 0 \quad \text{as } k \rightarrow \infty$$

Also note that

$$|U(f, \mathbf{P}_k)| \leq \sum_{\mathbf{J} \in \mathbf{P}_k} |M_{\mathbf{J}}| \text{vol} \mathbf{J} \leq M \frac{k^n - (k-2)^n}{k^n} \rightarrow 0 \quad \text{as } k \rightarrow \infty$$

which means that by triangle inequality

$$|U(f, \mathbf{P}_k) - L(f, \mathbf{P}_k)| \leq |U(f, \mathbf{P}_k)| + |L(f, \mathbf{P}_k)| \rightarrow 0 \quad \text{as } k \rightarrow \infty$$

meaning that

$$|L(f, \mathbf{P}_k)| \rightarrow 0 \quad \text{as } k \rightarrow \infty$$

Because the limit of the upper and lower Darboux sums are equal, it follows from the definition of the Riemann integral that f is integrable, and

$$\int_{\mathbf{I}} f = \lim_{k \rightarrow \infty} U(f, \mathbf{P}_k) = 0$$

□

Problem 3 (Exercise 7, Page 482)

For the generalized rectangle $\mathbf{I} = [0, 1] \times [0, 1]$ in the plane \mathbb{R}^2 , define

$$f(x, y) = \begin{cases} 5 & \text{if } (x, y) \text{ is in } \mathbf{I} \text{ and } x > 1/2 \\ 1 & \text{if } (x, y) \text{ is in } \mathbf{I} \text{ and } x \leq 1/2 \end{cases}$$

Use the Archimedes-Riemann Theorem to show that the function $f: \mathbf{I} \rightarrow \mathbb{R}$ is integrable.

Proof. Let $P_k = [(0, 1/2 - 1/3k, 1/2 + 1/3k, 1] \times [0, 1]$. We have that $U(f, P_k) = (1/2 - 1/3k) + 5(2/3k) + 5(1/2 - 1/3k)$ and $L(f, P_k) = (1/2 - 1/3k) + (2/3k) + 5(1/2 - 1/3k)$. Thus $U(f, P_k) - L(f, P_k) = 1/2k \rightarrow 0$, thus by the Archimedes Riemann theorem f is integrable. □

Problem 4 (Exercise 11, Page 482)

For the rectangle $\mathbf{I} = [0, 1] \times [0, 1]$ in the plane \mathbb{R}^2 , define the function $f: \mathbf{I} \rightarrow \mathbb{R}$ by

$$f(x, y) = xy \quad \text{for } (x, y) \text{ in } \mathbf{I}$$

Use the Archimedes-Riemann Theorem to evaluate $\int_{\mathbf{I}} f$.

Proof. Let \mathbf{P}_k be the regular partition of $[0, 1] \times [0, 1]$. Each rectangle in the partition has area $1/k^2$ and for indices i and j between 1 and k , we set

$$\mathbf{J} = \left[\frac{i-1}{k}, \frac{i}{k} \right] \times \left[\frac{j-1}{k}, \frac{j}{k} \right]$$

then

$$m(f, \mathbf{J}) = \frac{(i-1)(j-1)}{k^2} \quad \text{and} \quad M(f, \mathbf{J}) = \frac{ij}{k^2}.$$

It follows that

$$U(f, \mathbf{P}_k) = \sum_{\mathbf{J} \in \mathbf{P}_k} M(f, \mathbf{J}) \text{vol } \mathbf{J} = \sum_{1 \leq i, j \leq k} \frac{ij}{k^4} = \frac{1}{k^4} \sum_{i=1}^k i \left[\sum_{j=1}^k j \right] = \frac{1}{k^4} \left[\frac{k(k+1)}{2} \right]^2$$

Similarly, we note that

$$L(f, \mathbf{P}_k) = \frac{1}{k^4} \left[\frac{k(k-1)}{2} \right]^2.$$

We observe that

$$\lim_{k \rightarrow \infty} U(f, \mathbf{P}_k) = 1/4 \quad \text{and} \quad \lim_{k \rightarrow \infty} L(f, \mathbf{P}_k) = 1/4.$$

Thus we deduce that $\{\mathbf{P}_k\}$ is an Archimedean sequence and it follows from the Archimedes-Riemann Theorem that f is integrable and

$$\int_{\mathbf{I}} f = 1/4.$$

□

Problem 5 (Exercise 8, Page 489)

Let $\{\mathbf{u}_k\}$ be a convergent sequence in \mathbb{R}^n . Show that the set $\{\mathbf{u}_k \mid k \in \mathbb{N}\}$ has Jordan content 0.

Proof. Suppose that $\{\mathbf{u}_k\}$ converges to \mathbf{u} . Let $\varepsilon > 0$ be arbitrary and take $r > 0$ such that $r < \left(\frac{\varepsilon}{2}\right)^n$. By the convergence of $\{\mathbf{u}_k\}$, there exists K such that for all $k \geq K$ we have that $\mathbf{u}_k \in B_{r/2}(\mathbf{u})$. We can cover this open ball with the rectangle

$$R_{K+1} = \left[u_1 - \frac{r}{2}, u_1 + \frac{r}{2}\right] \times \cdots \times \left[u_n - \frac{r}{2}, u_n + \frac{r}{2}\right]$$

where $u = (u_1, \dots, u_n)$ in components. It follows that $B_{r/2}(\mathbf{u}) \subset R_{K+1}$, so that this rectangle covers all of the elements after index K . For the finitely many elements with index $1 \leq i \leq K$, define R_i to be the rectangle of side length $\frac{r}{K^{\frac{1}{n}}}$ with each u_i in the center of R_i . Now let \mathcal{F} be the collection of rectangles $\{R_1, \dots, R_{K+1}\}$. This covers the set $\{\mathbf{u}_k \mid k \in \mathbb{N}\}$ and the sum of volumes is

$$\sum_{i=1}^{K+1} \text{vol } R_i = K \cdot \left(\frac{r}{K^{\frac{1}{n}}}\right)^n + r^n = 2r^n < \varepsilon$$

□

Problem 6 (Exercise 10, Page 489)

Let \mathbf{I} be a generalized rectangle in \mathbb{R}^n and suppose that the function $f: \mathbf{I} \rightarrow \mathbb{R}$ is continuous. Assume that $f(\mathbf{x}) \geq 0$ for all points \mathbf{x} in \mathbf{I} . Prove that if $\int_{\mathbf{I}} f = 0$, then $f(\mathbf{x}) = 0$ for all points \mathbf{x} in \mathbf{I} . Is continuity necessary for this to hold?

Proof. Suppose not; that is, there exists a point \mathbf{x}_0 such that $f(\mathbf{x}_0) > 0$. By the continuity of f , there exists an open ball that contains a rectangle \mathbf{J} on which $f(\mathbf{x}) > 0$ for all $\mathbf{x} \in \mathbf{J}$. Because \mathbf{J} is a rectangle, it is compact and thus attains its minimum value $m > 0$. The continuity of f implies that it is integrable, so that there exists a sequence of Archimedean partitions such that

$$\begin{aligned} \int_{\mathbf{I}} f &= \lim_{k \rightarrow \infty} L(f, \mathbf{P}_k) = \lim_{k \rightarrow \infty} \sum_{\mathbf{R} \in \mathbf{P}_k} m_{\mathbf{R}} \text{vol} \mathbf{R} \geq \lim_{k \rightarrow \infty} \sum_{\mathbf{R} \in \mathbf{J}} m_{\mathbf{R}} \text{vol} \mathbf{R} \\ &\geq \lim_{k \rightarrow \infty} \sum_{\mathbf{R} \in \mathbf{J}} m \text{vol} \mathbf{R} = \lim_{k \rightarrow \infty} m \text{vol} \mathbf{J} = m \text{vol} \mathbf{J} > 0 \end{aligned}$$

But this contradicts the assumption that the integral of f is 0.

The condition that f be continuous is necessary; consider a function $f: \mathbf{I} \rightarrow \mathbb{R}$ such that $f(\mathbf{x}) = 0$ for every $\mathbf{x} \in \mathbf{I}$ except for a single \mathbf{x}_0 at which $f(\mathbf{x}_0) = 1$. This function is integrable because the single point at which the function is discontinuous has Jordan content 0, and the integral is equal to 0. \square