

Newton-Raphson method (II)

Wednesday, September 18, 2024 10:30 AM

Class 10: September 18, 2024

Recall: Last class we introduced perhaps the most famous rootfinding method: the **Newton-Raphson** method. Given a twice continuously differentiable function f , we assumed we have access to evaluate $f(x)$ and $f'(x)$ at a given guess x_0 for our root. We find the equation of the tangent line (best linear approximation at x_0) $l(x) = f(x_0) + f'(x_0)(x - x_0)$, and set it equal to zero. This gives us the Newton step:

$$x_n = x_{n-1} - f(x_{n-1}) / f'(x_{n-1})$$

The main result on the Newton-Raphson method (and why it is so popular) is the following theorem, which we proved using the fact that Newton-Raphson is a Fixed Point Iteration for the function

$$g_{\text{NR}}(x) = x - f(x) / f'(x)$$

Theorem 1: (Convergence of Newton) If f in $C^2([a,b])$ and r a "single root" in $[a,b]$ (such that $f'(r)$ is not zero), then there exists a $\delta > 0$ such that if $|x_0 - r| < \delta$, then starting Newton at x_0 converges quadratically* to r .

This theorem is easily proven by showing that, given the assumptions, $g'_{\text{NR}}(r) = 0$.

What does this theorem tell us? In so many words: if we start Newton "close enough" to a single root, it converges quadratically. We saw examples of this quadratic convergence: once it gets going, you get to machine precision *very* fast; digits of precision double every step (e.g. 1,2,4,8,16).

However, there are some issues with Newton: for one, this gives us no guarantees if we start away from this "basin of quadratic convergence" (which we generally have no way to know) and if r is a multiple root, Newton only converges linearly. Today we will continue this discussion and give more examples.

WARM UP: NEWTON-RAPHSON

PSEUDOCODE:

INPUTS - $f(x)$, $f'(x)$, x_0 , TOL , n_{\max}

OUTPUTS - r

Initialize $n = 0$

While ($n \leq n_{\max}$ AND $|x_n - x_{n-1}| \geq TOL$)

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)};$$

$$n = n + 1$$

Return r

LAST TIME WE ASKED: WHAT IF $f(r) = 0$, $f'(r) = 0$?

• ROOT MULTIPLICITY: $f(r) = f'(r) = \dots = f^{(m-1)}(r) = 0$, $f^{(m)}(r) \neq 0$

$\hookrightarrow \exists h(x)$ s.t. $h(r) \neq 0$, $f(x) = (x-r)^m h(x)$.

$$g_{\text{NR}}(x) = x - \frac{f(x)}{f'(x)} = x - \frac{(x-r)^m h(x)}{(x-r)^{m-1} [m h(x) + (x-r) h'(x)]}$$

$$g'_{\text{NR}}(x) = 1 - \frac{(h(x) + (x-r) h'(x)) [m h(x) + (x-r) h'(x)] - ((x-r) h(x))^2}{[m h(x) + (x-r) h'(x)]^2}$$

$$\lim_{x \rightarrow r} g'_{\text{NR}}(x) = 1 - \frac{m h(r)}{m^2 h(r)} = 1 - \frac{1}{m} \begin{matrix} \xrightarrow{m=1} 0 \\ \xrightarrow{m>1} \in [\frac{1}{2}, 1) \end{matrix}$$

$$\lim_{x \rightarrow r} g_{NR}(x) = 1 - \frac{f(x)}{m^2 h(r)} = \underline{1 - 1/m} \rightarrow m > 1 \in (\frac{1}{2}, 1)$$

FIXES:

□ Given $f(x)$ w/ multiple root at r .

$$\hookrightarrow f(x) = (x-r)^m h(r),$$

$$f'(x) = (x-r)^{m-1} [\quad]$$

$$\boxed{f(x)/f'(x)} = \frac{h(r)(x-r)}{1}$$

Apply Newton to f/f' . (requires $(f/f')'$)

⊙ $g_{NR}(x) = x - f(x)/f'(x)$

$\tilde{g}_{NR}(x) = x - \alpha f(x)/f'(x)$ what is α
s.t. $\tilde{g}_{NR}'(r) = 0?$

"Quasi Newton" \rightarrow Secant method.

- Historically what is $f'(x)$?
- Currently — $f'(x)$ can be expensive,
or hard to compute accurately.

$$\frac{f(x+h) - f(x)}{h} \approx f'(x)$$

\rightarrow Quasi Newton:

- Want to evaluate $f'(x)$ 1 or 0 times.
- Want superlinear convergence close to r .

1ST IDEA: "Lazy Newton" (Chord Iteration)

$$1 \ 1 \setminus \dots$$

$$x_n = x_{n-1} - \left(\frac{1}{f'(x_0)} \right) f(x_{n-1}).$$

⑥ f' is only evaluated once ✓

⑦ Convergence \rightarrow Linear (when it converges).

Secant method

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

x_0, x_1 close to each other and close to r ,

$$l_{\text{sec}}(x) = f(x_1) + \underbrace{\left[\frac{f(x_1) - f(x_0)}{x_1 - x_0} \right]}_{\approx f'(x_1)} (x - x_1)$$

$$x_2 = x_1 - \underbrace{\frac{1}{m_{\text{SEC}}}}_{\left(\frac{x_1 - x_0}{f(x_1) - f(x_0)} \right)} \cdot (f(x_1))$$