



Protein serves as the building block nutrient throughout the meal prep process, providing the nutrients essential to build and maintaining lean muscle while ensuring proper metabolic functioning. In this video, Chris discusses how he categorizes his protein sources throughout prep while discussing when and why he recommends specific protein sources over others.

Chris's Key Takeaways

Fattier meats, such as salmon and steak, contain greater micronutrient levels compared to other meats.

Wild Atlantic salmon will have higher Omega-3s and will contain slightly less dietary fat compared to farmed salmon.

Minimize traveling with fish due to the possibility of smell.

The fat content of sirloin or flank steak will vary, ranging from 3-4g of fat per 3oz cut.

Flank steak will have higher levels of fat marbling, whereas with sirloin, most of the fat resides on the edges of the cut.

Vacuum-sealed ground bison (1lb packages) usually have a 90/10 fat content, ranging between 9-11g of fat per 4oz.

Vacuum-sealed ground meats have a refrigerator shelf life when unopened for up to 2 weeks.

Ground meats will taste better when eaten fresh versus prepped for a later meal.

Steak is easily traveled with, resulting in minimal changes to the flavor after storing and consuming for multiple days.

Salmon has a higher fat content compared to sirloin and flank steak.

Planning a meal with a fatty protein source at the end of the day will help with satiety due to the increased time required for digestion.

Cod and chicken breast are leaner protein sources commonly used throughout prep.

The prevalence of higher mercury levels in fish often increases in the fattier, larger predatory fish.

Skipjack tuna is a medium-to-small-sized tuna that is known for having lower levels of mercury.

Cook and consume cod fresh or within 1-2 days.

Supplementing other meat proteins with cod can speed up weight loss due to its extremely low-fat content.

When purchasing chicken breast from a grocery store, choose air-chilled chicken for optimal flavor and minimal water retention.