





Common Hip Injuries in Bodybuilding



Injury Type	General Information	Symptoms	Relevant Risk Factors
Femoracetabular Impingement (FAI)  PINCER IMPINGEMENT CAM IMPINGEMENT <i>Adapted from Pepermpron/Shutterstock.com</i>	There are two types of FAI: cam and pincer; both types create abnormal contact with the acetabulum-femur during movement (particularly hip flexion and internal rotation), producing pain secondary to labrum irritation and loss of motion.	The C-sign pain description (see the Getting Technical callout box following this table); central groin and/or lateral hip pain with/without clicking; anterior groin pain is worse with prolonged sitting, standing, and walking; pain with squatting.	Loss of hip internal rotation; increased body mass; decreased hip flexor and gluteal strength.
Hip Bursitis  <i>Adapted from Pepermpron/Shutterstock.com</i>	Swelling and irritation of the bursa; bursas are numerous throughout the body and assist in providing frictionless interfaces between connective tissue structures and with additional stress; may become inflamed and cause pain with movement.	Pain with movement that may be present in the anterior thigh (iliopsoas bursitis) with recruitment of the hip flexors or lateral hip (trochanteric bursitis).	Overuse; hip flexor weakness; hip flexor length test (Thomas Test).

Common Hip Injuries in Bodybuilding



Injury Type	General Information	Symptoms	Relevant Risk Factors
Hamstring Strain  Adapted from Pepermprom/Shutterstock.com	Occurs when the hamstrings are placed under excessive stress (typically from eccentric-induced muscle actions) that result in variable amounts of tearing of the muscle fibers of the hamstring musculature.	Depends on the severity of the muscle tear; pain in the deep posterior thigh with/without presence of bruising and associated muscle weakness in hip extension and knee flexion.	Eccentric hamstring muscle weakness; weakness in hip extension with the knee bent; limitations in hip flexion with knee extension.
Piriformis Syndrome  Adapted from Pepermprom/Shutterstock.com	Thought to occur with irritation of the sciatic nerve as it passes through or near the piriformis muscle producing local and/or radiating pain down the affected leg.	Pain in the gluteal region with or without radiating pain down the leg; pain with or reproduction of pain with sitting and squatting; reproduction of pain with the piriformis stretch.	Gluteus maximus/gluteus medius weakness is thought to predispose the piriformis muscle to work harder to produce similar joint actions, leading to overuse.