

Impairments, Considerations, and Corrections

Consideration		Determination	Correction
Motor Learning (Cueing & Feedback)		- First, determine what are/were the performance client's impairment findings from the assessment process - Is there any pain? If so, stop the lift and consider referring out - Based on this information, determine if the performance client can get into the correct position(s) for the lift they are performing (i.e., maintain knee and spine alignment throughout motion)	- Make sure the performance client is in proper kinetic chain alignment - Make sure the performance client is aware of the impairment(s) that are occurring - Provide specific cues and feedback to help the performance client maintain correct posture and alignment. If this does not resolve the impairment, it may be more of a mobility or stability/strength issue.
Neuromuscular Control	Mobility (ROM)	- Is there any pain? If so, stop the lift and consider referring out - Determine if the impairment happens at a specific point in the movement or from the beginning. In other words, is the performance client able to maintain correct posture and alignment to a certain point before the impairment(s) appear	- Make sure the exercise selected is appropriate for the performance client's current level of mobility - Makes sure the performance client has the proper kinetic chain alignment relative to the lift they are performing (i.e., feet wide enough properly turned, proper grip, elbows properly positioned, etc.) If this does not resolve the issue, it may be a stability/strength issue.
	Stability/Strength (Load)	- Is there any pain? If so, stop the lift and consider referring out - Determine if the performance client was able to perform the lift at lighter loads with minimal to no impairment(s) and if the impairment occurs later in the set or after they are fatigued	- Make sure the load is appropriate for the performance client's current level of stability/ strength - Increased loads require greater stability & strength to perform and can often make the performance client move into greater ROM, which they may not yet have - May need to change the number of reps or sets depending on when impairment occurs If this does not resolve the issue, it may be a combination of these elements.