



case_study_report


Connie Wu, Jason McEachin, Joe Choo, Scott Heng

10/10/2020

```
#pairs plot to show correlation between response variables  
train_response <- data.frame(train[,c("R_moment_1", "R_moment_2", "R_moment_3", "R_moment_4")])  
pairs(train_response)
```



```
#correlation between predictor variables (in case we need to include interaction effects)  
train_predictors <- data.frame(train[,c("Re", "Fr", "St")])  
pairs(train_predictors)
```



```
#linear regression with factored Re and Fr
```

```
train1 <- train
train1$Re <- as.factor(train$Re)
train1$Fr <- as.factor(train$Fr)
lm1_m1 <- lm(R_moment_1 ~ St + Fr + Re, data=train1)
lm2_m1 <- lm(R_moment_2 ~ St + Fr + Re, data=train1)
lm3_m1 <- lm(R_moment_3 ~ St + Fr + Re, data=train1)
lm4_m1 <- lm(R_moment_4 ~ St + Fr + Re, data=train1)
summary(lm1_m1)
```

```
##
## Call:
## lm(formula = R_moment_1 ~ St + Fr + Re, data = train1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.038834 -0.008614  0.001702  0.009854  0.039423
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  0.106488   0.003971  26.818 < 2e-16 ***
## St           0.012213   0.002078   5.877 8.42e-08 ***
## Fr0.3        -0.007623   0.004245  -1.796  0.07618 .
## FrInf        -0.010210   0.003787  -2.696  0.00849 **
## Re224        -0.108091   0.003682 -29.353 < 2e-16 ***
## Re398        -0.111553   0.004632 -24.081 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.01529 on 83 degrees of freedom
## Multiple R-squared:  0.9293, Adjusted R-squared:  0.9251
```

F-statistic: 218.2 on 5 and 83 DF, p-value: < 2.2e-16

Introduction

Turbulence is highly versatile motion that is often times difficult to predict and understanding fluid motion and its effect on natural problems poses a great challenge. Interpreting turbulence data is an incredibly important task in the engineering from understanding the cosmos to the cosmic cycle.

Parametric modeling is effective when we want to compactly represent features as model parameters. Unlike certain “black box” machine learning techniques, it offers a higher level of interpretability, which is especially useful for a practical setting. Rather than simply outputting a classification result, a parametric model allows us to investigate in more detail which aspects of turbulence differ between high and low particle cluster volumes.

In our project, we use linear and (...) and apply it to interpret the difference in model parameters in order to achieve the the following objectives:

Methodology

Results

Conclusion