



Xoring Ninja

by [amititkgp](#)

Problem

Submissions

Leaderboard

Discussions

Editorial

An **XOR** operation on a list is defined here as the *xor* (\oplus) of all its elements (e.g.: $XOR(\{A, B, C\}) = A \oplus B \oplus C$).

The *XorSum* of set S is defined here as the sum of the *XORs* of all S 's non-empty subsets. If we refer to the set of S 's non-empty subsets as S' , this can be expressed as:

$$XorSum(S) = \sum_{i=1}^{2^n-1} XOR(S'_i) = XOR(S'_1) + XOR(S'_2) + \dots + XOR(S'_{2^n-2}) + XOR(S'_{2^n-1})$$

For example: Given set $S = \{n_1, n_2, n_3\}$

- The set of possible non-empty subsets is: $S' = \{\{n_1\}, \{n_2\}, \{n_3\}, \{n_1, n_2\}, \{n_1, n_3\}, \{n_2, n_3\}, \{n_1, n_2, n_3\}\}$
- The *XorSum* of these non-empty subsets is then calculated as follows:
 $XorSum(S) = n_1 + n_2 + n_3 + (n_1 \oplus n_2) + (n_1 \oplus n_3) + (n_2 \oplus n_3) + (n_1 \oplus n_2 \oplus n_3)$

Given a list of n space-separated integers, determine and print $XorSum \% (10^9 + 7)$.

Note: The cardinality of `powerset(n)` is 2^n , so the set of non-empty subsets of set S of size n contains $2^n - 1$ subsets.

Input Format

The first line contains an integer, T , denoting the number of test cases.

Each test case consists of two lines; the first is an integer, n , describing the size of the set, and the second contains n space-separated integers (a_1, a_2, \dots, a_n) describing the set.

Constraints

$$1 \leq T \leq 5$$

$$1 \leq n \leq 10^5$$

$$0 \leq a_i \leq 10^9, i \in [1, n]$$

Output Format

For each test case, print its $XorSum \% (10^9 + 7)$ on a new line; the i^{th} line should contain the output for the i^{th} test case.

Sample Input

```
1
3
1 2 3
```

Sample Output

```
12
```

Explanation

The input set, $S = \{1, 2, 3\}$, has 7 possible non-empty subsets: $S' = \{\{1\}, \{2\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 3\}, \{1, 2, 3\}\}$.

We then determine the *XOR* of each subset in S' :

$$XOR(\{1\}) = 1$$

$XOR(\{2\}) = 2$
 $XOR(\{3\}) = 3$
 $XOR(\{1, 2\}) = 1 \oplus 2 = 3$
 $XOR(\{2, 3\}) = 2 \oplus 3 = 1$
 $XOR(\{1, 3\}) = 1 \oplus 3 = 2$
 $XOR(\{1, 2, 3\}) = 1 \oplus 2 \oplus 3 = 0$

Then sum the results of the XOR of each individual subset in S' , resulting in $XorSum = 12$. We print 12, because $12 \% (10^9 + 7) = 12$.

Submissions: 2018

Max Score: 55

Difficulty: Difficult

 Download PDF

 Suggest Edits

Choose a translation

English

[Collapse](#)

Current Buffer (saved locally, editable)  

C



```
1 #include <stdio.h>
2 #include <string.h>
3 #include <math.h>
4 #include <stdlib.h>
5
6 int main() {
7
8     /* Enter your code here. Read input from STDIN. Print output to STDOUT */
9     return 0;
10 }
11
```

Line: 1 Col: 1

 Upload Code as File

☐ Test against custom input

Run Code

Submit Code

Copyright © 2016 HackerRank. All Rights Reserved

Join us on IRC at [#hackerrank](#) on freenode for hugs or bugs.

[Contest Calendar](#) | [Blog](#) | [Scoring](#) | [Environment](#) | [FAQ](#) | [About Us](#) | [Support](#) | [Careers](#) | [Privacy Policy](#) | [Request a Feature](#)