

# Dummit & Foote Ch. 4.3: Groups Acting on Themselves by Conjugation — The Class Equation

Scott Donaldson

Feb. 2024

Let  $G$  be a group.

## 1. (2/22/24)

Suppose  $G$  has a left action on a set  $A$ , denoted by  $g \cdot a$  for all  $g \in G$  and  $a \in A$ . Denote the corresponding right action on  $A$  by  $a \cdot g$ . Prove that the (equivalence) relations  $\sim$  and  $\sim'$  defined by

$$a \sim b \text{ if and only if } a = g \cdot b \text{ for some } g \in G$$

and

$$a \sim' b \text{ if and only if } a = b \cdot g \text{ for some } g \in G$$

are the same relation (i.e.,  $a \sim b$  if and only if  $a \sim' b$ ).

*Proof.* To show that  $a \sim b$  implies  $a \sim' b$ , we must show that, given a  $g \in G$  with  $a = g \cdot b$ , there exists an  $h \in G$  such that  $a = b \cdot h$ . By definition, the corresponding right action of a left action is specified to be  $g \cdot x = x \cdot g^{-1}$  for all  $g \in G$ ,  $x \in A$ . Letting  $h = g^{-1}$ , we have found an element where  $a = g \cdot b = b \cdot h$ , and so  $a \sim' b$ .

The proof for  $a \sim' b$  implies  $a \sim b$  is identical, letting  $h = g^{-1}$  but with  $h$  acting on the left.  $\square$

## 2. (2/22/24)

Find all conjugacy classes and their sizes in the following groups:

(a)  $D_8$ :

$$\{1\}_1 \quad \{r^2\}_1 \quad \{r, r^3\}_2 \quad \{s, sr^2\}_2 \quad \{sr, sr^3\}_2$$

(b)  $Q_8$ :

$$\{1\}_1 \quad \{-1\}_1 \quad \{\pm i\}_2 \quad \{\pm j\}_2 \quad \{\pm k\}_2$$

(c)  $A_4$ :

$$\begin{aligned} \{1\}_1 \quad & \{(1\ 2\ 3), (1\ 3\ 4), (1\ 4\ 2), (2\ 4\ 3)\}_4 \quad \{(1\ 3\ 2), (1\ 2\ 4), (1\ 4\ 3), (2\ 3\ 4)\}_4 \\ & \{(1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)\}_3 \end{aligned}$$

### 3. (2/22/24)

Find all the conjugacy classes and their sizes in the following groups:

(a)  $Z_2 \times S_3$ :

$$\begin{aligned} & \{(0, 1)\}_1 \quad \{(1, 1)\}_1 \quad \{(0, (1\ 2)), (0, (1\ 3)), (0, (2\ 3))\}_3 \\ & \{(1, (1\ 2)), (1, (1\ 3)), (1, (2\ 3))\}_3 \quad \{(0, (1\ 2\ 3)), (0, (1\ 3\ 2))\}_2 \\ & \{(1, (1\ 2\ 3)), (1, (1\ 3\ 2))\}_2 \end{aligned}$$

(b)  $S_3 \times S_3$ :

$$\begin{aligned} & \{(1, 1)\}_1 \quad \{(1, 2\text{-cycle})\}_3 \quad \{(2\text{-cycle}, 1)\}_3 \quad \{(1, 3\text{-cycle})\}_2 \quad \{(3\text{-cycle}, 1)\}_2 \\ & \{(2\text{-cycle}, 2\text{-cycle})\}_9 \quad \{(2\text{-cycle}, 3\text{-cycle})\}_6 \quad \{(3\text{-cycle}, 2\text{-cycle})\}_6 \\ & \{(3\text{-cycle}, 3\text{-cycle})\}_4 \end{aligned}$$

(c)  $Z_3 \times A_4$  (using representatives from the conjugacy classes of  $A_4$  above):

$$\begin{aligned} & \{(0, 1)\}_1 \quad \{(1, 1)\}_1 \quad \{(2, 1)\}_1 \\ & \{(0, \overline{(1\ 2\ 3)})\}_4 \quad \{(1, \overline{(1\ 2\ 3)})\}_4 \quad \{(2, \overline{(1\ 2\ 3)})\}_4 \\ & \{(0, \overline{(1\ 3\ 2)})\}_4 \quad \{(1, \overline{(1\ 3\ 2)})\}_4 \quad \{(2, \overline{(1\ 3\ 2)})\}_4 \\ & \{(0, \overline{(1\ 2)(3\ 4)})\}_3 \quad \{(1, \overline{(1\ 2)(3\ 4)})\}_3 \quad \{(2, \overline{(1\ 2)(3\ 4)})\}_3 \end{aligned}$$

### 4. (2/22/24)

Prove that if  $S \subseteq G$  and  $g \in G$  then  $gN_g(S)g^{-1} = N_G(gSg^{-1})$  and  $gC_g(S)g^{-1} = C_G(gSg^{-1})$ .

*Proof.* Let  $x \in N_G(S)$ . So  $xsx^{-1} \in S$  for all  $s \in S$ . Then

$$\begin{aligned} gxsx^{-1}g^{-1} & \in gSg^{-1} \\ gxx^{-1}gsg^{-1}gx^{-1}g^{-1} & \in gSg^{-1} \\ (gxx^{-1})gsg^{-1}(gx^{-1}g^{-1}) & \in gSg^{-1} \\ (gxx^{-1})gsg^{-1}(gxx^{-1})^{-1} & \in gSg^{-1}, \end{aligned}$$

which implies that  $gxg^{-1} \in N_G(gSg^{-1})$ , and so  $gN_G(S)g^{-1} \subseteq N_G(gSg^{-1})$ .

Conversely, let  $x \in N_G(gSg^{-1})$ . So  $xsg^{-1}x^{-1} \in gSg^{-1}$  for all  $s \in S$ . Then

$$\begin{aligned} xsg^{-1}x^{-1} &\in gSg^{-1} \\ g^{-1}xsg^{-1}x^{-1} &\in Sg^{-1} \\ g^{-1}xsg^{-1}x^{-1}g &\in S \\ (g^{-1}xg)s(g^{-1}xg)^{-1} &\in S \\ g^{-1}xg &\in N_G(S) \\ x &\in gN_G(S)g^{-1}, \end{aligned}$$

which shows that  $N_G(gSg^{-1}) \subseteq gN_G(S)g^{-1}$ . This proves that  $N_G(gSg^{-1}) = gN_G(S)g^{-1}$ .

Next, let  $x \in C_G(S)$ . So  $xs = sx$  for all  $s \in S$ . Then

$$\begin{aligned} xs &= sx \\ gsg^{-1} &= gsg^{-1} \\ gsg^{-1}gxg^{-1} &= gsg^{-1}gxg^{-1} \\ (gsg^{-1})(gxg^{-1}) &= (gsg^{-1})(gxg^{-1}), \end{aligned}$$

and so  $gxg^{-1} \in C_G(gSg^{-1})$ , which implies that  $gC_G(S)g^{-1} \subseteq C_G(gSg^{-1})$ . Finally, let  $x \in C_G(gSg^{-1})$ . So  $x(gsg^{-1}) = (gsg^{-1})x$  for all  $x \in S$ . Then

$$\begin{aligned} xsg^{-1} &= gsg^{-1}x \\ g^{-1}xsg^{-1} &= sg^{-1}x \\ g^{-1}xgs &= sg^{-1}xg \\ (g^{-1}xg)s &= s(g^{-1}xg), \end{aligned}$$

which implies that  $g^{-1}xg \in C_G(S)$ , so  $x \in gC_G(S)g^{-1}$ . It follows that  $C_G(gSg^{-1}) \subseteq gC_G(S)g^{-1}$ , and therefore  $gC_G(S)g^{-1} = C_G(gSg^{-1})$ .  $\square$

## 10. (2/28/24)

Let  $\sigma$  be the 5-cycle (1 2 3 4 5) in  $S_5$ . In each of (a) to (c) find an explicit element  $\tau \in S_5$  which accomplishes the specified conjugation:

- (a)  $\tau\sigma\tau^{-1} = \sigma^2 = (1\ 3\ 5\ 2\ 4)$ . Let  $\tau = (2\ 3\ 5\ 4)$ . Then  $\tau\sigma\tau^{-1} = (\tau(1)\tau(2)\tau(3)\tau(4)\tau(5)) = (1\ 3\ 5\ 2\ 4) = \sigma^2$ .
- (b)  $\tau\sigma\tau^{-1} = \sigma^{-1} = (1\ 5\ 4\ 3\ 2)$ . Let  $\tau = (2\ 5)(3\ 4)$ .
- (c)  $\tau\sigma\tau^{-1} = \sigma^{-2} = (1\ 4\ 2\ 5\ 3)$ . Let  $\tau = (2\ 4\ 5\ 3)$ .

## 11. (2/28/24)

In each of (a) - (d) determine whether  $\sigma_1$  and  $\sigma_2$  are conjugate. If they are, give an explicit permutation  $\tau$  such that  $\tau\sigma_1\tau^{-1} = \sigma_2$ .

- (a)  $\sigma_1 = (12)(345)$  and  $\sigma_2 = (123)(45)$ . Both have cycle type 1, 1, 3 and so they are conjugate. Let  $\tau = (14253)$ . Then  $\tau\sigma_1\tau^{-1} = \sigma_2$ .
- (b)  $\sigma_1 = (15)(372)(106811)$  and  $\sigma_2 = (37510)(49)(13112)$ . In  $S_{13}$ , both have cycle type 1, 1, 1, 1, 2, 3, 4 and so they are conjugate. Let  $\tau = (14)(211103)(5967138)$ . Then  $\tau\sigma_1\tau^{-1} = \sigma_2$ .
- (c)  $\sigma_1 = (15)(372)(106811)$  and  $\sigma_2 = \sigma_1^3 = (15)(101186)$ . They do not have the same cycle type ( $\sigma_1$  contains a 3-cycle that  $\sigma_2$  does not), and so they are not conjugate.
- (d)  $\sigma_1 = (13)(246)$  and  $\sigma_2 = (35)(24)(56) = (24)(356)$ . Let  $\tau = (12345)$ . Then  $\tau\sigma_1\tau^{-1} = \sigma_2$ .

## 13. (2/28/24)

Find all finite groups which have exactly two conjugacy classes.

*Proof.* Let  $G$  be a non-trivial finite group. Since the conjugacy class of 1 is  $\{1\}$ , if  $G$  has exactly two conjugacy classes, then every other element in  $G$  must have the same conjugacy class, namely  $G - \{1\}$ .

From Proposition 6, for any  $g \in G$ , the number of conjugates of  $g$  (i.e. the cardinality of the conjugacy class of  $g$ ) is the index of the centralizer of  $g$ ,  $|G : C_G(g)|$ . Therefore the size of the conjugacy class of  $g$  must divide the order of  $G$ .

Let  $|G| = n$ . Then the size of the conjugacy class of  $g$  is  $|G - \{1\}| = n - 1$ . This is only possible when  $|G| = 2$ , and so  $G$  must be the unique group of order two.  $\square$