

Dummit & Foote Ch. 2.3: Cyclic Groups and Cyclic Subgroups

Scott Donaldson

Jun. 2023

1. (6/18/23)

Find all subgroups of $Z_{45} = \langle x \rangle$, giving a generator for each. Describe the containments between these subgroups.

Proof. The subgroups of $Z_{45} = \langle x \rangle$ are those cyclic groups generated by x^n , where n divides 45. These are:

- $\langle 1 \rangle = \{1\}$, the trivial subgroup
- $\langle x^{15} \rangle = \{1, x^{15}, x^{30}\} \cong \mathbb{Z}/3\mathbb{Z}$
- $\langle x^9 \rangle = \{1, x^9, x^{18}, x^{27}, x^{36}\} \cong \mathbb{Z}/5\mathbb{Z}$
- $\langle x^5 \rangle = \{1, x^5, x^{10}, x^{15}, x^{20}, x^{25}, x^{30}, x^{35}, x^{40}\} \cong \mathbb{Z}/9\mathbb{Z}$
- $\langle x^3 \rangle = \{1, x^3, x^6, \dots, x^{39}, x^{42}\} \cong \mathbb{Z}/15\mathbb{Z}$
- $\langle x \rangle = Z_{45}$ itself

Among these subgroups, we have $\langle 1 \rangle$ contained within every other subgroup, as well as $\langle x^{15} \rangle \leq \langle x^5 \rangle$, $\langle x^{15} \rangle \leq \langle x^3 \rangle$, and $\langle x^9 \rangle \leq \langle x^3 \rangle$. \square

2. (6/19/23)

If x is an element of the finite group G and $|x| = |G|$, prove that $G = \langle x \rangle$. Give an explicit example to show that this result need not be true if G is an infinite group.

Proof. Let $|x| = |G| = n < \infty$. By definition, G is closed, so it contains all powers of $x : 1, x, x^2, \dots, x^{n-1}$. These are exactly n elements, so G contains no other elements. It is therefore generated by x , that is, $G = \langle x \rangle$.

However, if G is an infinite group and $x \in G$ with $|x| = \infty$, then this is not necessarily the case. For example, if $G = \mathbb{Z}$ and $x = 2$, then x generates all even integers in \mathbb{Z} , but does not generate the element 5. \square

3. (6/19/23)

Find all generators for $\mathbb{Z}/48\mathbb{Z}$.

Proof. From Proposition 6., the generators for $\mathbb{Z}/48\mathbb{Z}$ are those positive integers $n < 48$ for which n is relatively prime to 48. These are: 1, 5, 7, 11, 13, 17, 19, 23, 25, 29, 31, 35, 37, 41, 43, and 47. \square

4. (6/19/23)

Find all generators for $\mathbb{Z}/202\mathbb{Z}$.

Proof. As above, the generators for $\mathbb{Z}/202\mathbb{Z}$ are those positive integers $n < 202$ for which n is relatively prime to 202. The integer 202 only has two divisors greater than 1, namely 2 and 101. Therefore the generators of $\mathbb{Z}/202\mathbb{Z}$ are every odd positive integer less than 202 except for 101. \square

5. (6/19/23)

Find the number of generators for $\mathbb{Z}/49000\mathbb{Z}$.

Proof. We are concerned with the number of integers n between 0 and 48999 for which n is relatively prime to 49000. It will be helpful to write 49000 uniquely as the product of primes: $2^3 \cdot 5^3 \cdot 7^2$.

Let us first consider the generators for $\mathbb{Z}/49000\mathbb{Z}$ between 0 and 69, that is, all the numbers that are relatively prime to 49000 between 0 and 69: 1, 3, 9, 11, 13, 17, 19, 23, 27, 29, 31, 33, 37, 39, 41, 43, 47, 51, 53, 57, 59, 61, 67, and 69. There are 24 such generators.

Next, we show that, for any $n \in \{0, \dots, 48999\}$, the greatest common divisor of n and 49000 is equal to the greatest common divisor of $n \bmod 70$ and 49000. This is because 70 is equal to the product of the bases of the prime factors of 49000: $70 = 2 \cdot 5 \cdot 7$. So for any n , we have $n = m + 70k = m + (2 \cdot 5 \cdot 7)k$, where $m \in \{0, \dots, 69\}$ and $k \geq 0$. Suppose that m is *not* in the list of the above generators (that is, that the greatest common divisor of m and 49000 is greater than 1). Then either 2, 5, or 7 divides m (otherwise m would be relatively prime to 49000). Without loss of generality, suppose that 2 divides m , and write $m = 2p$. We can then rewrite n as:

$$n = m + (2 \cdot 5 \cdot 7)k = 2p + (2 \cdot 5 \cdot 7)k = 2(p + (5 \cdot 7)k),$$

that is, 2 divides n , so it is not relatively prime to 49000 (similarly, if 5 or 7 divide m , then 5 or 7 also divide n , respectively). It follows that the generators for $\mathbb{Z}/49000\mathbb{Z}$ between 0 and 69 repeat (mod 70) over the rest of 49000. Since $49000/70 = 700$, there are thus $700 \cdot 24 = 16800$ generators for $\mathbb{Z}/49000\mathbb{Z}$. \square

6. (6/20/23)

In $\mathbb{Z}/48\mathbb{Z}$ write out all elements of $\langle \bar{a} \rangle$ for every \bar{a} . Find all inclusions between subgroups in $\mathbb{Z}/48\mathbb{Z}$.

- Subgroup of order 48: $\langle \bar{1} \rangle = \langle \bar{5} \rangle = \langle \bar{7} \rangle = \langle \bar{11} \rangle = \langle \bar{13} \rangle = \langle \bar{17} \rangle = \langle \bar{19} \rangle = \langle \bar{23} \rangle = \langle \bar{25} \rangle = \langle \bar{29} \rangle = \langle \bar{31} \rangle = \langle \bar{35} \rangle = \langle \bar{37} \rangle = \langle \bar{41} \rangle = \langle \bar{43} \rangle = \langle \bar{47} \rangle$.
- Subgroup of order 24: $\langle \bar{2} \rangle = \langle \bar{10} \rangle = \langle \bar{14} \rangle = \langle \bar{22} \rangle = \langle \bar{26} \rangle = \langle \bar{34} \rangle = \langle \bar{38} \rangle = \langle \bar{46} \rangle$.
- Subgroup of order 16: $\langle \bar{3} \rangle = \langle \bar{9} \rangle = \langle \bar{15} \rangle = \langle \bar{21} \rangle = \langle \bar{27} \rangle = \langle \bar{33} \rangle = \langle \bar{39} \rangle = \langle \bar{45} \rangle$.
- Subgroup of order 12: $\langle \bar{4} \rangle = \langle \bar{20} \rangle = \langle \bar{28} \rangle = \langle \bar{44} \rangle$.
- Subgroup of order 8: $\langle \bar{6} \rangle = \langle \bar{18} \rangle = \langle \bar{30} \rangle = \langle \bar{42} \rangle$.
- Subgroup of order 6: $\langle \bar{8} \rangle = \langle \bar{40} \rangle$.
- Subgroup of order 4: $\langle \bar{12} \rangle = \langle \bar{36} \rangle$.
- Subgroup of order 3: $\langle \bar{16} \rangle = \langle \bar{32} \rangle$.
- Subgroup of order 2: $\langle \bar{24} \rangle$.
- Subgroup of order 1, the trivial subgroup: $\{0\}$.

Among these subgroups, all contain the trivial subgroup. The subgroups of order 2 and 3 are distinct, but both are contained in the subgroup of order 6. The subgroup of order 2 is also contained in the subgroup of order 4. The subgroups of order 4 and 6 are both contained in the subgroup of order 12. The subgroup of order 4 is also contained in the subgroup of order 8. The subgroups of order 8 and 12 are both contained in the subgroup of order 24. The subgroup of order 8 is also contained in the subgroup of order 16.

7. (6/22/23)

Let $Z_{48} = \langle x \rangle$ and use the isomorphism $\mathbb{Z}/48\mathbb{Z} \cong Z_{48}$ given by $\bar{1} \mapsto x$ to list all subgroups of Z_{48} as computed in the preceding exercise.

- Subgroup of order 48: $\{1, x, x^2, \dots, x^{47}\}$.
- Subgroup of order 24: $\{1, x^2, x^4, \dots, x^{46}\}$.
- Subgroup of order 16: $\{1, x^3, x^6, \dots, x^{45}\}$.
- Subgroup of order 12: $\{1, x^4, x^8, \dots, x^{44}\}$.
- Subgroup of order 8: $\{1, x^6, x^{12}, x^{18}, x^{24}, x^{30}, x^{36}, x^{42}\}$.

- Subgroup of order 6: $\{1, x^8, x^{16}, x^{24}, x^{32}, x^{40}\}$.
- Subgroup of order 4: $\{1, x^{12}, x^{24}, x^{36}\}$.
- Subgroup of order 3: $\{1, x^{16}, x^{32}\}$.
- Subgroup of order 2: $\{1, x^{24}\}$.
- Subgroup of order 1, the trivial subgroup: $\{1\}$.

8. (6/23/23)

Let $Z_{48} = \langle x \rangle$. For which integers a does the map φ_a defined by $\varphi_a : \bar{1} \mapsto x^a$ extend to an *isomorphism* from $\mathbb{Z}/48\mathbb{Z}$ onto Z_{48} ?

Proof. We will show that φ_a is an isomorphism from $\mathbb{Z}/48\mathbb{Z}$ onto Z_{48} if and only if $a \in \mathbb{Z}$ is relatively prime to 48.

First, let $m, n \in \mathbb{Z}/48\mathbb{Z}$. Then $\varphi_a(m)\varphi_a(n) = (x^a)^m(x^a)^n = (x^a)^{m+n} = \varphi_a(m+n)$. So φ_a is a homomorphism.

Next, φ_a is one-to-one. Let $\varphi_a(n) = \varphi_a(m)$ for $m, n \in \mathbb{Z}/48\mathbb{Z}$. Then $(x^a)^m = (x^a)^n \Rightarrow x^{am} = x^{an}$, and so $am = an \pmod{48}$. Since a is relatively prime to 48, we must therefore have $m = n$, and it follows that φ_a is injective. (Note, however, that if $k > 1$ divides both a and 48, then $am = an$ does not imply that $m = n$, and φ_a is therefore not injective. For example, if $a = 14$, then $\varphi_a(7) = (x^{14})^7 = x^{98} = x^2$ and $\varphi_a(31) = (x^{14})^{31} = x^{434} = x^2$).

Finally, φ_a is onto. Let $x^b \in Z_{48}$. Suppose there exists some $n \in \mathbb{Z}/48\mathbb{Z}$ such that $\varphi_a(n) = x^b$, that is, $(x^a)^n = x^b$. Then we must have $an = b \pmod{48}$. Since a is relatively prime to 48, any integer between 0 and 47 can be written as an for some $n \in \mathbb{Z}/48\mathbb{Z}$, and so φ_a is onto.

Thus for a relatively prime to 48, $\varphi_a : \bar{1} \mapsto x^a$ is an isomorphism from $\mathbb{Z}/48\mathbb{Z}$ onto Z_{48} . \square

9. (7/2/23)

Let $Z_{36} = \langle x \rangle$. For which integers a does the map φ_a defined by $\varphi_a : \bar{1} \mapsto x^a$ extend to a *well defined homomorphism* from $\mathbb{Z}/48\mathbb{Z}$ onto Z_{36} ? Can φ_a ever be a surjective homomorphism?

Proof. We will show that $\varphi_a : \mathbb{Z}/48\mathbb{Z} \rightarrow Z_{36}$ is a well defined homomorphism if and only if a is a multiple of 3.

For φ_a to be a homomorphism, we must have $\varphi_a(b)\varphi_a(c) = \varphi_a(b+c)$ for all $b, c \in \mathbb{Z}/48\mathbb{Z}$. Now $\varphi_a(b)\varphi_a(c) = (x^a)^b(x^a)^c = (x^a)^{b+c} = x^{a(b+c)}$ and $\varphi_a(b+c) = (x^a)^{b+c} = x^{a(b+c)}$. Superficially these appear identical already. However, note that in $\varphi_a(b)\varphi_a(c)$ we compute $ab + ac \pmod{36}$, while in $\varphi_a(b+c)$ we first take $b+c \pmod{48}$ before then computing $a(b+c)$. That is, a must satisfy

$$a(b+c \pmod{48}) \pmod{36} = a(b+c) \pmod{36}$$

for all $b, c \in \mathbb{Z}/48\mathbb{Z}$. If $b + c < 48$, then the two are equal for all $a \in \mathbb{Z}$. So suppose that $b + c \geq 48$. Then $b + c \bmod 48 = b + c - 48$, so we must have

$$\begin{aligned} a(b + c - 48) \bmod 36 &= a(b + c) \bmod 36 \\ ab + ac - 48a \bmod 36 &= ab + ac \bmod 36 \\ -48a \bmod 36 &= 0 \bmod 36 \\ -48a &\cong 36 \Rightarrow 48a \cong 36, \end{aligned}$$

that is, a is some integer which, when multiplied by 48, results in a multiple of 36. Writing 48 as the product of its prime factors gives $2^4 \cdot 3$, while $36 = 2^2 \cdot 3^2$. Note that 36 has one more factor of 3, and so when a is a multiple of 3, $48a$ will be a multiple of 36. Only these values satisfy the exponents in the equation above, and thus φ_a is a homomorphism if and only if a is a multiple of 3.

It is not possible for φ_a to be a surjective homomorphism. Because a must be a multiple of 3, we have $\varphi_a(1) = x^a = x^{3n} = (x^3)^n$ for some $n \in \mathbb{Z}$. In turn, φ_a generates only the values $\varphi_a(2) = (x^6)^n, \varphi_a(3) = (x^9)^n, \dots$, that is, it only generates powers of x^3 in Z_{36} . By counterexample, there is no value in $\mathbb{Z}/48\mathbb{Z}$ whose image under φ_a is x , and so φ_a cannot be surjective. \square