


# Dummit & Foote Ch. 2.5: The Lattice of Subgroups of a Group

Scott Donaldson

Aug. 2023

## 1. (8/11/23)

Let  $H$  and  $K$  be subgroups of  $G$ . Exhibit all possible sublattices which show only  $G$ ,  $1$ ,  $H$ ,  $K$ , and their joins and intersections. What distinguishes the different drawings?



The left two lattices show the group structure when either  $H \leq K$  or  $K \leq H$  (they omit any subgroups of the smaller of the two, as well as any containing subgroups between the larger and  $G$ ).

The next lattice shows the group structure when  $H$  and  $K$  are not comparable, their intersection consists only of the identity, and their join is all of  $G$ . The final three lattices show the cases where  $H \cap K$  is a subgroup not equal to the identity, where  $\langle H, K \rangle$  is a subgroup not equal to  $G$ , and where both of these occur.

## 2. (8/11/23)

In each of (a) to (d) list all subgroups of  $D_{16}$  that satisfy the given condition.

- (a) Subgroups that are contained in  $\langle sr^2, r^4 \rangle$   
 $\{1\}, \langle sr^6 \rangle, \langle sr^2 \rangle, \langle r^4 \rangle, \langle sr^2, r^4 \rangle$

- (b) Subgroups that are contained in  $\langle sr^7, r^4 \rangle$   
 $\{1\}, \langle sr^3 \rangle, \langle sr^7 \rangle, \langle r^4 \rangle, \langle sr^7, r^4 \rangle$
- (c) Subgroups that contain  $\langle r^4 \rangle$   
 $\langle r^4 \rangle, \langle sr^2, r^4 \rangle, \langle s, r^4 \rangle, \langle r^2 \rangle, \langle sr^3, r^4 \rangle, \langle sr^5, r^4 \rangle, \langle s, r^2 \rangle, \langle r \rangle, \langle sr, r^2 \rangle, D_{16}$
- (d) Subgroups that contain  $\langle s \rangle$   
 $\langle s \rangle, \langle s, r^4 \rangle, \langle s, r^2 \rangle, \langle D_{16} \rangle$

### 3. (8/11/23)

Show that the subgroup  $\langle s, r^2 \rangle$  of  $D_8$  is isomorphic to  $V_4$ .

*Proof.* The subgroup  $\langle s, r^2 \rangle$  of  $D_8$  contains the elements  $\{1, s, r^2, sr^2\}$ . There is no element in this group of order 4. From Ch. 1.1, Exercise 36, there is only one unique group of order 4 with no element of order 4, the Klein group  $V_4$ . Thus  $\langle s, r^2 \rangle$  is isomorphic to  $V_4$ .  $\square$

### 4. (8/14/23)

Use the given lattice to find all pairs of elements that generate  $D_8$ .

*Proof.* Since  $D_8$  is generated by  $\langle s, r \rangle$ , it suffices to find pairs of elements that generate  $s$  and  $r$ . These pairs of elements are:

- $\langle s, r \rangle$  (trivial)
- $\langle s, r^3 \rangle$  ( $r = (r^3)^3$ )
- $\langle s, sr \rangle$  ( $r = s \cdot sr$ )
- $\langle s, sr^3 \rangle$  ( $r = s \cdot (sr^3)^3$ )
- $\langle sr, r \rangle$  ( $s = r \cdot sr$ )
- $\langle sr, r^2 \rangle$  ( $r^3 = r^2 \cdot sr, r = (r^3)^3, s = r \cdot sr$ )
- $\langle sr, r^3 \rangle$  ( $r = (r^3)^3, s = r \cdot sr$ )
- $\langle sr^2, r \rangle$  ( $s = sr^2 \cdot r^2$ )
- $\langle sr^2, r^3 \rangle$  ( $r = (r^3)^3, s = sr^2 \cdot r^2$ )
- $\langle sr^2, sr^3 \rangle$  ( $r = sr^2 \cdot sr^3, s = sr^2 \cdot r^2$ )
- $\langle sr^3, r \rangle$  ( $s = sr^3 \cdot r$ )
- $\langle sr^3, r^3 \rangle$  ( $s = r^3 \cdot sr^3, r = s \cdot sr^3$ )

$\square$

## 5. (8/14/23)

Use the given lattice to find all elements  $x \in D_{16}$  such that  $D_{16} = \langle x, s \rangle$ .

*Proof.* The element  $x \in D_{16}$  generates  $D_{16}$  together with  $s$  if  $r$  can be expressed as a product of  $s$  and  $x$ :

- $x = r$  (trivial)
- $x = r^3$  ( $r = (r^3)^3$ )
- $x = r^5$  ( $r = (r^5)^5$ )
- $x = r^7$  ( $r = (r^7)^7$ )
- $x = sr$  ( $r = s \cdot sr$ )
- $x = sr^3$  ( $r^3 = s \cdot sr^3$ ,  $r = (r^3)^3$ )
- $x = sr^5$  ( $r^5 = s \cdot sr^5$ ,  $r = (r^5)^5$ )
- $x = sr^7$  ( $r^7 = s \cdot sr^7$ ,  $r = (r^7)^7$ )

□

## 6. (8/14/23)

Find the centralizers of every element in the following groups:

(a)  $D_8$

- 1:  $D_8$
- $r, r^2, r^3$ :  $\langle r \rangle$
- $s, sr^2$ :  $\langle s, r^2 \rangle$
- $sr, sr^3$ :  $\langle sr, r^2 \rangle$

(b)  $Q_8$

- 1, -1:  $Q_8$
- $i, -i$ :  $\langle i \rangle$
- $j, -j$ :  $\langle j \rangle$
- $k, -k$ :  $\langle k \rangle$

(c)  $S_3$

- (1):  $S_3$
- (1, 2):  $\langle (1, 2) \rangle$
- (1, 3):  $\langle (1, 3) \rangle$
- (2, 3):  $\langle (2, 3) \rangle$
- (1, 2, 3), (1, 3, 2):  $\langle (1, 2, 3) \rangle$

(d)  $D_{16}$

- 1:  $D_{16}$
- $r, r^2, \dots, r^7$ :  $\langle r \rangle$
- $s, sr^4$ :  $\langle s, r^4 \rangle$
- $sr, sr^5$ :  $\langle sr, r^4 \rangle$
- $sr^2, sr^6$ :  $\langle sr^2, r^4 \rangle$
- $sr^3, sr^7$ :  $\langle sr^3, r^4 \rangle$

## 7. (8/14/23)

Find the center of  $D_{16}$ .

*Proof.* From the preceding exercise, the only elements that are in the centralizer of every element of  $D_{16}$  are  $\{1, r^4\} = \langle r^4 \rangle$ .  $\square$

## 8. (8/14/23)

In each of the following groups find the normalizer of each subgroup:

- (a)  $S_3$ : The subgroups (other than (1) and all of  $S_3$ ) are the three cyclic groups generated by each of the 2-cycles, and the group consisting of  $\{(1), (1, 2, 3), (1, 3, 2)\}$ . In the case of  $\langle (1, 2) \rangle$ , notice that:

$$(1, 3)(1, 2)(1, 3)^{-1} = (1, 3)(1, 2)(1, 3) = (2, 3) \notin \langle (1, 2) \rangle,$$


which implies that  $(1, 3) \notin N_{S_3}(\langle (1, 2) \rangle)$ . By extension, no 2-cycle is in the normalizer of another 2-cycle. There is no subgroup of  $S_3$  that contains a 2-cycle, a 3-cycle, but does *not* contain a different 2-cycle. Therefore each cyclic subgroup of  $S_3$  is its own normalizer.

Now for the subgroup  $\{(1), (1, 2, 3), (1, 3, 2)\}$ , we have  $(1, 2)(1, 2, 3)(1, 2) = (1, 3, 2)$ , which is included in the subgroup. It follows that the normalizer of this subgroup is all of  $S_3$ .

- (b)  $Q_8$ : The elements 1 and  $-1$  commute with all elements of  $Q_8$ , so the normalizer of  $\langle -1 \rangle$  is all of  $Q_8$ . Consider the normalizer of  $\langle i \rangle$ . Now  $j \cdot i \cdot j^{-1} = j \cdot i \cdot -j = -k \cdot -j = i$ , so  $j \in N_{Q_8}(\langle i \rangle)$ . Then the normalizer of  $i$  contains at least 5 elements, so it must be all of  $Q_8$ . By extension, every subgroup of  $Q_8$  is its own normalizer.

## 9. (8/14/23)

Draw the lattices of subgroups of the following groups:



## 10. (8/15/23)

Classify groups of order 4 by proving that if  $|G| = 4$  then  $G \cong Z_4$  or  $G \cong V_4$ .

*Proof.* From Ch. 1.1, Exercise 36, if  $G$  is a group with 4 elements and no element of order 4, then we must have  $G = \langle a, b \mid a^2 = b^2 = 1, ab = ba \rangle \cong V_4$ .

If  $G$  does have an element of order 4, then we have the cyclic group  $G = \langle a \mid a^4 = 1 \rangle \cong Z_4$ . All cyclic groups of equal order are isomorphic.


Therefore a group of order 4 must be isomorphic to either the cyclic group of order 4 or the Klein 4-group.  $\square$

## 11. (8/15/23)

Consider the group of order 16 with the following presentation:


$$QD_{16} = \langle \sigma, \tau \mid \sigma^8 = \tau^2 = 1, \sigma\tau = \tau\sigma^3 \rangle$$

(called the *quasidihedral* or *semidihedral* group of order 16). This group has three subgroups of order 8:  $\langle \tau, \sigma^2 \rangle \cong D_8$ ,  $\langle \sigma \rangle \cong Z_8$  and  $\langle \sigma^2, \sigma\tau \rangle \cong Q_8$  and every proper subgroup is contained in one of these three subgroups. Fill in the lattice of all subgroups of the quasidihedral group, exhibiting each subgroup with at most two generators.



## 12. (8/16/23)

The group  $A = Z_2 \times Z_4 = \langle a, b \mid a^2 = b^4 = 1, ab = ba \rangle$  has order 8 and has three subgroups of order 4:  $\langle a, b^2 \rangle \cong V_4$ ,  $\langle b \rangle \cong Z_4$  and  $\langle ab \rangle \cong Z_4$  and every proper subgroup is contained in one of these three. Draw the lattice of all subgroups of  $A$ , giving each subgroup in terms of at most two generators.



## 13. (8/16/23)

The group  $G = Z_2 \times Z_8 = \langle x, y \mid x^2 = y^8 = 1, xy = yx \rangle$  has order 16 and has three proper subgroups of order 8:  $\langle x, y^2 \rangle \cong Z_2 \times Z_4$ ,  $\langle y \rangle \cong Z_8$  and  $\langle xy \rangle \cong Z_8$  and every proper subgroup is contained in one of these three. Draw the lattice of all subgroups of  $G$ , giving each subgroup in terms of at most two generators.

