

Dummit & Foote Ch. 1.6: Homomorphisms and Isomorphisms

Scott Donaldson

Mar. - Apr. 2023

1. (3/25/23)

Let $\varphi : G \rightarrow H$ be a homomorphism.

- (a) Prove that $\varphi(x^n) = \varphi(x)^n$ for all $n \in \mathbb{Z}^+$.

Proof. By induction. When $n = 1$, $\varphi(x^1) = \varphi(x) = \varphi(x)^1$.

Suppose for some n , $\varphi(x^n) = \varphi(x)^n$. Then $\varphi(x^{n+1}) = \varphi(x^n x)$. By definition, because φ is a homomorphism from G to H , $\varphi(ab) = \varphi(a)\varphi(b)$ for all $a, b \in G$. So $\varphi(x^n x) = \varphi(x^n)\varphi(x)$. By the induction hypothesis, $\varphi(x^n) = \varphi(x)^n$, so this equals $\varphi(x)^{n+1}$.

Therefore $\varphi(x^n) = \varphi(x)^n$ for all $n \in \mathbb{Z}^+$. \square

- (b) Do part (a) for $n = -1$ and deduce that $\varphi(x^n) = \varphi(x)^n$ for all $n \in \mathbb{Z}$.

This proof diverges slightly from the directions but arrives at the same result.

Note that, for all $x \in G$, $\varphi(x) = \varphi(1 \cdot x) = \varphi(1)\varphi(x)$. Therefore $\varphi(1) = 1$ (in H). Now $1 = \varphi(1) = \varphi(x^n \cdot x^{-n}) = \varphi(x^n)\varphi(x^{-n})$. From part a), this equals $\varphi(x)^n \varphi(x^{-n})$. Left-multiplying both sides by $\varphi(x)^{-n}$, we obtain $\varphi(x^{-n}) = \varphi(x)^{-n}$, as desired.

2. (3/26/23)

If $\varphi : G \rightarrow H$ is an isomorphism, prove that $|\varphi(x)| = |x|$ for all $x \in G$. Deduce that any two isomorphic groups have the same number of elements of order n for each $n \in \mathbb{Z}^+$.

Proof. Let $\varphi : G \rightarrow H$ be an isomorphism and let $x \in G$. If $|x|$ is finite, then (from 1.a) $\varphi(x^n) = \varphi(x)^n$ and (from 1.b) $\varphi(1) = \varphi(x^n) = \varphi(x)^n = 1 \in H$. The order of the element $\varphi(x)^n \in H$ is therefore at most n . Because φ is an

isomorphism, there is only one element whose image is 1, and that is $\varphi(1) = 1$. Therefore for no $m < n$ do we have $\varphi(x)^m = 1$, and so the $|\varphi(x)| = n$.

Next, suppose that x has infinite order in G . Then $x^n \neq 1$ for all $n > 0$. Because φ is an isomorphism, we know that only $\varphi(1) = 1 \in H$. Therefore $\varphi(x^n) = \varphi(x)^n \neq 1$ for all $n > 0$. Therefore $|\varphi(x)| = \infty$.

This result is not necessarily true if φ is a homomorphism. For example, φ could send every element of G to the identity in H . (This is a homomorphism: $\varphi(x)\varphi(y) = 1 \cdot 1 = 1$ and $\varphi(x)\varphi(y) = \varphi(xy) = 1$.) Then for all $x \in G$, $|\varphi(x)| = 1$, regardless of the order of x . \square

3. (3/27/23)

If $\varphi : G \rightarrow H$ is an isomorphism, prove that G is abelian if and only if H is abelian. If φ is a homomorphism, what additional conditions on φ (if any) are sufficient to ensure that if G is abelian, then so is H ?

Proof. First, let G be an abelian group and $\varphi : G \rightarrow H$ be an isomorphism. Given arbitrary distinct elements of H , because φ is surjective, there are two distinct elements in G whose images are these elements in H . Let $\varphi(x), \varphi(y) \in H$ be distinct elements and $x, y \in G$. Then $\varphi(xy) = \varphi(x)\varphi(y)$. Also, because x and y commute, $\varphi(xy) = \varphi(yx) = \varphi(y)\varphi(x)$. Therefore $\varphi(x)\varphi(y) = \varphi(y)\varphi(x)$, so H is an abelian group.

Next, let H be an abelian group. Again let $\varphi(x), \varphi(y) \in H$ and $x, y \in G$. Then $\varphi(x)\varphi(y) = \varphi(xy)$. Also, $\varphi(x)\varphi(y) = \varphi(y)\varphi(x) = \varphi(yx)$. So $\varphi(xy) = \varphi(yx)$. Because φ is one-to-one, this implies that $xy = yx$, and so G is an abelian group.

If φ is a homomorphism, then G being an abelian group does not imply that H is abelian. For example, H could be a non-abelian group and φ could send every element of G to the identity in H .

A sufficient condition for a homomorphism $\varphi : G \rightarrow H$ to ensure that if G is abelian, then so is H , is that φ is surjective. Then for all $h \in H$, $h = \varphi(x)$ for some $x \in G$ (possibly more than one x). Let $h_1, h_2 \in H$ with $h_1 = \varphi(x_1) = \varphi(x_2) = \dots$ and $h_2 = \varphi(y_1) = \varphi(y_2) = \dots$ and with $x_i, y_j \in G$. φ is a homomorphism, so for any i, j , $\varphi(x_i y_j) = \varphi(x_i)\varphi(y_j) = h_1 h_2$. Also, because G is abelian, $\varphi(x_i y_j) = \varphi(y_j x_i) = \varphi(y_j)\varphi(x_i) = h_2 h_1$. Therefore $h_1 h_2 = h_2 h_1$, so H is abelian. \square

4. (3/27/23)

Prove that the multiplicative groups $\mathbb{R} - \{0\}$ and $\mathbb{C} - \{0\}$ are not isomorphic.

Proof. For any $x \in \mathbb{R} - \{0\}$, $x \neq \pm 1$, x has infinite order. The proof of this is as follows: Let $x \in \mathbb{R} - \{0, \pm 1\}$. If the absolute value of x is greater than 1, then the absolute value of x^n is greater than 1 for all n , and by induction x has infinite order. If the absolute value of x is less than 1, then the absolute value

of x^n is less than 1 for all n , and by induction x has infinite order. So 1 and -1 are the only elements of $\mathbb{R} - \{0\}$ with finite order.

In $\mathbb{C} - \{0\}$, i and $-i$ have order 4. From 2., isomorphic groups have the same number of elements of order n for each $n \in \mathbb{Z}^+$. However, $\mathbb{R} - \{0\}$ has no elements of order 4, and $\mathbb{C} - \{0\}$ has at least 2. Therefore they are not isomorphic. \square

5. (3/27/23)

Prove that the additive groups \mathbb{R} and \mathbb{Q} are not isomorphic.

Proof. Given that \mathbb{R} and \mathbb{Q} do not have the same cardinality (\mathbb{R} is uncountable while \mathbb{Q} is countably infinite), there is no map $\varphi : \mathbb{Q} \rightarrow \mathbb{R}$ that is surjective. An isomorphism is a bijection that is necessarily surjective, and so the two groups are not isomorphic.

Alternatively, consider the homomorphism $\varphi : \mathbb{Q} \rightarrow \mathbb{R}$ defined by $\varphi(q) = q$. Such a map is injective but not surjective: There is no $q \in \mathbb{Q}$ with $\varphi(q) = \sqrt{2} \in \mathbb{R}$. If we attempt to make φ surjective by assigning $\varphi(q_1) = \sqrt{2}$ for some q_1 , then q_1 now has no preimage in \mathbb{Q} , and so we must find a q_2 and assign $\varphi(q_2) = q_1$. However, now q_2 has no preimage. This process continues *ad infinitum*, and φ is forever not surjective. Therefore \mathbb{R} and \mathbb{Q} are not isomorphic. \square

6. (3/27/23)

Prove that the additive groups \mathbb{Z} and \mathbb{Q} are not isomorphic.

Proof. Consider a homomorphism $\varphi : \mathbb{Z} \rightarrow \mathbb{Q}$. For all $n \in \mathbb{Z}$, $\varphi(0) = \varphi(n + (-n)) = \varphi(n) + \varphi(-n)$. From 1.b), $\varphi(0) = 0$, so φ preserves inverses: $\varphi(-n) = -\varphi(n)$. That is, $\varphi(n) = q$ implies that $\varphi(-n) = -q$.

We also claim that, if $\varphi(1) = k$, then φ assigns all integers to their product with k in \mathbb{Q} . Since φ preserves inverses, we only have to show this for $n \in \mathbb{Z}^+$, by induction (base case given): Suppose that $\varphi(n) = kn$ for some $n \in \mathbb{Q}^+$. Then $\varphi(n+1) = \varphi(n) + \varphi(1) = kn + k = k(n+1)$, as desired. Therefore φ assigns all integers to their product with k in \mathbb{Q} .

But now it is impossible for φ to be surjective, because only integer multiples of k have preimages in \mathbb{Z} . For example, $k/2 \in \mathbb{Q}$ has no preimage. Therefore \mathbb{Z} and \mathbb{Q} are not isomorphic. \square

7. (3/27/23)

Prove that D_8 and Q_8 are not isomorphic.

Proof. $s, sr, sr^2, sr^3 \in D_8$ all have order 2. However, in Q_8 , only -1 has order 2. From 2., isomorphic groups must have the same number of elements of each order. Therefore D_8 and Q_8 are not isomorphic. \square

8. (3/28/23)

Prove that if $n \neq m$, S_n and S_m are not isomorphic.

Proof. Without loss of generality, let $n > m$. From Chapter 1.3, the order of a symmetric group S_n is $n!$. Then S_n contains $n!$ elements, and S_m contains $m!$ elements. It is trivial to show that $n > m \Rightarrow n! > m!$. Since the two groups do not have the same cardinality, there is no bijection between them. Thus S_n and S_m are not isomorphic. \square

9. (3/28/23)

Prove that D_{24} and S_4 are not isomorphic.

Proof. D_{24} has 24 elements, and S_4 has 24 elements. They are both non-abelian. In order to prove that they are not isomorphic, then, let us consider the orders of each group's respective elements.

D_{24} has 13 elements of order 2: $\{sr^i \mid i \in \{0, \dots, 11\}\}$ and r^6 .

The elements of order 2 in S_4 are those permutations with cycle decompositions that are disjoint 2-cycles:

$\{(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4), (1, 2)(3, 4), (1, 3)(2, 4), (1, 4)(2, 3)\}$. So there are 9 elements of order 2 in S_4 .

Since D_{24} and S_4 do not have the same number of elements of order 2, they are not isomorphic. \square

10. (3/31/23)

Fill in the details of the proof that the symmetric groups S_Δ and S_Ω are isomorphic if $|\Delta| = |\Omega|$ as follows: Let $\theta : \Delta \rightarrow \Omega$ be a bijection. Define

$$\varphi : S_\Delta \rightarrow S_\Omega \text{ by } \varphi(\sigma) = \theta \circ \sigma \circ \theta^{-1} \text{ for all } \sigma \in S_\Delta$$

and prove the following:

- (a) φ is well-defined, that is, if σ is a permutation of Δ then $\theta \circ \sigma \circ \theta^{-1}$ is a permutation of Ω .

To show that φ is well-defined, we need to show that it assigns a given permutation of Δ to a unique permutation of Ω .

An arbitrary permutation σ is a bijection from Δ to itself. It is represented with a cycle decomposition that shows how it assigns a given element of Δ to another element. For σ and a given element s_1 , we can say that σ assigns s_1 to $s_2 \in \Delta$.

Since Δ and Ω have the same cardinality, there exists a bijection θ between them, and we can say that θ assigns distinct $s_1, s_2 \in \Delta$ to distinct $t_1, t_2 \in \Omega$, respectively.

Now let us consider what happens when we apply φ to σ . By definition, $\varphi(\sigma) = \theta \circ \sigma \circ \theta^{-1}$. θ^{-1} is a bijection: $\Omega \rightarrow \Delta$, σ is a bijection: $\Delta \rightarrow \Delta$, and θ is a bijection: $\Delta \rightarrow \Omega$. Applying the compositions, we see that $\varphi(\sigma)$ is a map from $\Omega \rightarrow \Omega$ (not yet proven to be a bijection).

t_1 is an arbitrary element of Ω with preimage $s_1 \in \Delta$. Then:

$$\varphi(\sigma)(t_1) = \theta(\sigma(\theta^{-1}(t_1))) = \theta(\sigma(s_1)) = \theta(s_2) = t_2,$$

that is, $\varphi(\sigma)$ is a permutation of Ω that uniquely assigns t_1 to t_2 . Therefore φ is well-defined.

(b) φ is a bijection from S_Δ onto S_Ω .

We have shown that φ is a well-defined map from S_Δ onto S_Ω . However, it remains to be shown that φ is a bijection.

To show that φ is invertible, define a map $\gamma : S_\Omega \rightarrow S_\Delta$, with $\gamma(\tau) = \theta^{-1} \circ \tau \circ \theta$ for $\tau \in \Omega$. The proof above suffices to show that γ is well-defined.

Consider what happens when we take $\gamma(\varphi(\sigma))$:

$$\gamma(\varphi(\sigma)) = \gamma(\theta \circ \sigma \circ \theta^{-1}) = \theta^{-1} \circ (\theta \circ \sigma \circ \theta^{-1}) \circ \theta = (\theta^{-1} \theta) \circ \sigma \circ (\theta^{-1} \theta) = \sigma.$$

That is, $\gamma(\varphi(\sigma)) = \sigma$ for all $\sigma \in S_\Delta$. Therefore $\gamma = \varphi^{-1}$. Since φ has a well-defined inverse, it is a bijection from S_Δ onto S_Ω .

(c) φ is a homomorphism, that is, $\varphi(\sigma \circ \tau) = \varphi(\sigma) \circ \varphi(\tau)$.

We apply the function compositions:

$$\begin{aligned} \varphi(\sigma \circ \tau) &= \\ (\theta \circ \sigma \circ \theta^{-1}) \circ (\theta \circ \tau \circ \theta^{-1}) &= \theta \circ \sigma \circ (\theta^{-1} \circ \theta) \circ \tau \circ \theta^{-1} = \\ &= \theta \circ \sigma \circ \tau \circ \theta^{-1} = \varphi(\sigma) \circ \varphi(\tau). \end{aligned}$$

Thus φ is a homomorphism, and since it is also a bijection, the groups S_Δ and S_Ω are isomorphic.

11. (4/1/23)

Let A and B be groups. Prove that $A \times B \cong B \times A$.

Proof. Consider the map $\varphi : A \times B \rightarrow B \times A$ defined by $\varphi(a, b) = (b, a)$. φ is injective, since $\varphi(a_1, b_1) = \varphi(a_2, b_2) \Rightarrow (b_1, a_1) = (b_2, a_2) \Rightarrow a_1 = a_2$ and $b_1 = b_2$. φ is surjective, since for every $(b, a) \in B \times A$, there exists by definition $(a, b) \in A \times B$ with $\varphi(a, b) = (b, a)$. Therefore φ is a bijection from $A \times B \rightarrow B \times A$.

φ is also a homomorphism: Let $(a_1, b_1), (a_2, b_2) \in A \times B$. Then:

$$\begin{aligned}\varphi((a_1, b_1)(a_2, b_2)) &= \varphi(a_1 a_2, b_1 b_2) = \\ &= (b_1 b_2, a_1 a_2) = (b_1, a_1)(b_2, a_2) = \varphi(a_1, b_1)\varphi(a_2, b_2).\end{aligned}$$

Since φ is a bijective homomorphism, it is an isomorphism, and so $A \times B \cong B \times A$. \square

12. (4/5/23)

Let A, B , and C be groups and let $G = A \times B$ and $H = B \times C$. Prove that $G \times C \cong A \times H$.

Proof. Let $\varphi : G \times C \rightarrow A \times H$ defined by $\varphi \circ ((a, b), c) = (a, (b, c))$. We will show that φ is a bijective homomorphism, that is, an isomorphism, and thus that $G \times C \cong A \times H$.

To show that φ is injective, let $((a_1, b_1), c_1)$ and $((a_2, b_2), c_2) \in G \times C$, and suppose that applying φ to both gives the same element $(a, (b, c)) \in A \times H$. Then, by definition of φ , $a_1 = a$ and $a_2 = a$, so $a_1 = a_2$. The same logic shows that $b_1 = b_2$ and $c_1 = c_2$. Thus the two elements in G are in fact the same element, and therefore φ is injective.

To show that φ is surjective, let $(a, (b, c)) \in A \times H$. Then, by definition of φ , there exists $a, b, c \in A, B, C$, respectively, such that for $((a, b), c) \in G \times C$, $\varphi \circ ((a, b), c) = (a, (b, c))$. Therefore φ is surjective, and so it is a bijection.

Finally, let $((a_1, b_1), c_1)$ and $((a_2, b_2), c_2) \in G \times C$. Then:

$$\begin{aligned}\varphi \circ (((a_1, b_1), c_1)((a_2, b_2), c_2)) &= \varphi \circ ((a_1 a_2, b_1 b_2), c_1 c_2) = (a_1 a_2, (b_1 b_2, c_1 c_2)) = \\ &= (a_1, (b_1, c_1))(a_2, (b_2, c_2)) = \varphi \circ ((a_1, b_1), c_1)\varphi \circ ((a_2, b_2), c_2).\end{aligned}$$

Thus φ is an isomorphism from $G \times C \rightarrow A \times H$, and so $G \times C \cong A \times H$. \square

13. (4/5/23)

Let G and H be groups and let $\varphi : G \rightarrow H$ be a homomorphism. Prove that the image of φ , $\varphi(G)$, is a subgroup of H . Prove that if φ is injective then $G \cong \varphi(G)$.

Proof. To prove that $\varphi(G)$ is a subgroup of H , we must show that it is closed under the binary operation of H and that it is closed under inverses (the other group properties follow from these). By definition, for $a, b \in G$, $\varphi(a), \varphi(b) \in H$. Since φ is a homomorphism, their product, $\varphi(a)\varphi(b) = \varphi(ab)$, is also an element of H . Thus $\varphi(G)$ is closed under the binary operation of H .

To show that $\varphi(G)$ is closed under inverses, let $a \in G$. From 1.b), $\varphi(1) = 1$. Also, $\varphi(1) = \varphi(aa^{-1}) = \varphi(a)\varphi(a^{-1})$. Therefore $\varphi(a^{-1}) = \varphi(a)^{-1}$, and so $\varphi(G)$ is closed under inverses. Thus it is a subgroup of H .

Now suppose that φ is injective. To prove that G is then isomorphic to $\varphi(G)$, we must show that φ is an isomorphism, that is, that it is also surjective onto $\varphi(G)$. By definition, since $\varphi(G) = \{h \in H \mid h = \varphi(g) \text{ for some } g \in G\}$, φ is surjective onto $\varphi(G)$. Thus φ is an isomorphism, and so $G \cong \varphi(G)$. \square

14. (4/9/23)

Let G and H be groups and let $\varphi : G \rightarrow H$ be a homomorphism. Define the *kernel* of φ to be $\{g \in G \mid \varphi(g) = 1_H\}$ (so the kernel is the set of elements in G which map to the identity in H , i.e., is the fiber over the identity of H). Prove that the kernel of φ is a subgroup of G . Prove that φ is injective if and only if the kernel of φ is the identity subgroup of G .

Proof. To show that the kernel of φ is a subgroup of G , we need to show that it is closed under the binary operation of G and that it is closed under inverses.

Let g_1, g_2 be in the kernel of φ . Then $\varphi(g_1) = \varphi(g_2) = 1_H$. Since φ is a homomorphism, $1_H = \varphi(g_1)\varphi(g_2) = \varphi(g_1g_2)$. So the product g_1g_2 of arbitrary elements in the kernel of G is also in the kernel of G . Thus the kernel of φ is closed under the binary operation of G .

From 1.b), $\varphi(1) = 1_H$. Also, $1_H = \varphi(1) = \varphi(gg^{-1}) = \varphi(g)\varphi(g^{-1}) = 1_H \cdot \varphi(g^{-1})$, which implies that $\varphi(g^{-1}) = 1_H$, so g^{-1} is also in the kernel of φ . The kernel of φ is closed under inverses, and thus it is a subgroup of G .

Now we will show that φ is injective if and only if the kernel of φ is $\{1_G\}$.

First, let φ be an injective homomorphism from G to H . So for any $g_1, g_2 \in G$, $\varphi(g_1) = \varphi(g_2)$ implies that $g_1 = g_2$. Let g be in the kernel of φ , so $\varphi(g) = 1_H$. Also, $\varphi(1_G) = 1_H$, which implies that $g = 1_G$, so the only unique element in the kernel of φ is 1_G .

Next, suppose the kernel of φ is $\{1_G\}$. So $g \in G, g \neq 1_G$ implies that $\varphi(g) \neq 1_H$. Suppose that $g_1, g_2 \in G, g_1, g_2 \neq 1_G$ such that $\varphi(g_1) = \varphi(g_2) = h \in H$. From 1.b), $\varphi(g_2) = h$ implies that $\varphi(g_2)^{-1} = \varphi(g_2^{-1}) = h^{-1}$. So $\varphi(g_1)\varphi(g_2^{-1}) = hh^{-1} = 1_H$. Because φ is a homomorphism, $\varphi(g_1g_2^{-1}) = 1_H$. Because the only element in the kernel of φ is 1_G , we must have $g_1g_2^{-1} = 1_G$, which implies that $g_1 = g_2$. Therefore φ is an injective map from G to H .

This completes the proof that φ is injective if and only if the kernel of φ is $\{1_G\}$. \square

15. (4/9/23)

Define a map $\pi : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $\pi((x, y)) = x$. Prove that π is a homomorphism and find the kernel of π .

Proof. To show that π is a homomorphism, let $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$. Then $\pi((x_1, y_1)(x_2, y_2)) = \pi((x_1x_2, y_1y_2)) = x_1x_2$. Also, $\pi((x_1, y_1)) \cdot \pi((x_2, y_2)) = x_1x_2$. Thus π is a homomorphism.

By definition, the kernel of π is the set $\{(x, y) \in \mathbb{R}^2 \mid \pi(x, y) = 1\}$. Now $\pi((x, y)) = 1$ if and only if $x = 1$. Note that $x = 1 \Rightarrow \pi((1, y)) = 1$ and $\pi((x, y)) = 1 \Rightarrow x = 1$. So the kernel of π is $\{(1, y) \in \mathbb{R}^2 \mid y \in \mathbb{R}\}$. \square

16. (4/10/23)

Let A and B be groups and let G be their direct product, $A \times B$. Prove that the maps $\pi_1 : G \rightarrow A$ and $\pi_2 : G \rightarrow B$ defined by $\pi_1((a, b)) = a$ and $\pi_2((a, b)) = b$ are homomorphisms and find their kernels.

Proof. To show that π_1 and π_2 are homomorphisms, let $(a_1, b_1), (a_2, b_2) \in G$. Then $\pi_1((a_1, b_1)(a_2, b_2)) = \pi_1((a_1 a_2, b_1 b_2)) = a_1 a_2$ and $\pi_1((a_1, b_1)) \cdot \pi_1((a_2, b_2)) = a_1 a_2$, so π_1 is a homomorphism. Similarly, $\pi_2((a_1, b_1)(a_2, b_2)) = b_1 b_2 = \pi_2((a_1, b_1)) \cdot \pi_2((a_2, b_2))$, so it is also a homomorphism.

Now by identical proof to 15., the kernel of π_1 is $\{(1, b) \in G \mid b \in B\}$ and the kernel of π_2 is $\{(a, 1) \in G \mid a \in A\}$. \square