

# Dummit & Foote Ch. 3.4: Composition Series and the Hölder Program

Scott Donaldson

Oct. 2023

## 1. (11/2/23)

Prove that if  $G$  is an abelian simple group then  $G \cong Z_p$  for some prime  $p$  (do not assume  $G$  is a finite group).

*Proof.* Since  $G$  is simple, the only normal subgroups of  $G$  are 1 and  $G$  itself. However, since  $G$  is abelian, any subgroup of  $G$  must be normal, so it follows that  $G$  contains *no* subgroups other than 1 and itself.

If  $x_1, x_2 \in G$  are distinct generators for  $G$ , then  $\langle x_1 \rangle$  and  $\langle x_2 \rangle$  would be distinct subgroups of  $G$ ; therefore  $G$  is generated by a single element and is a cyclic group. Let us write  $G = \langle x \rangle$ . If  $G$  were infinite, then for any  $n > 1$ ,  $\langle x^n \rangle$  would be a distinct subgroup of  $G$ , so  $G$  must be finite.

Finally, if  $n$  divides  $|G|$ , then from Chapter 2, Theorem 7.(3),  $G$  contains a proper subgroup of order  $n$ . Therefore  $|G|$  has no divisors other than 1 and itself, so we have  $|G| = p$  for some prime  $p$ . We conclude that  $G \cong Z_p$  for some prime  $p$ .  $\square$

## 2. (11/3/23)

Exhibit all 3 composition series for  $Q_8$  and all 7 composition series for  $D_8$ . List the composition factors in each case.

The 3 composition series for  $Q_8$  are:

1.  $1 \leq \langle -1 \rangle \leq \langle i \rangle \leq Q_8$
2.  $1 \leq \langle -1 \rangle \leq \langle j \rangle \leq Q_8$
3.  $1 \leq \langle -1 \rangle \leq \langle k \rangle \leq Q_8$

In each series, each composition factor is isomorphic to  $Z_2$  (thus each  $N_i$  is normal in  $N_{i+1}$ ; since there is only one left coset it must equal the only right coset).

The 7 composition series for  $D_8$  are:

1.  $1 \leq \langle s \rangle \leq \langle s, r^2 \rangle \leq D_8$

2.  $1 \leq \langle sr^2 \rangle \leq \langle s, r^2 \rangle \leq D_8$
3.  $1 \leq \langle r^2 \rangle \leq \langle s, r^2 \rangle \leq D_8$
4.  $1 \leq \langle r^2 \rangle \leq \langle r \rangle \leq D_8$
5.  $1 \leq \langle r^2 \rangle \leq \langle sr, r^2 \rangle \leq D_8$
6.  $1 \leq \langle sr \rangle \leq \langle sr, r^2 \rangle \leq D_8$
7.  $1 \leq \langle sr^3 \rangle \leq \langle sr, r^2 \rangle \leq D_8$

Again each composition factor is isomorphic to  $Z_2$ .

### 3. (11/3/23)

Find a composition series for the quasidihedral group of order 16 (cf. Exercise 11, Section 2.5). Deduce that  $QD_{16}$  is solvable.

*Solution.* Recall that  $QD_{16} = \langle \sigma, \tau \mid \sigma^8 = \tau^2 = 1, \sigma\tau = \tau\sigma^3 \rangle$ .

A composition series for  $QD_{16}$  is:

$$1 \leq \langle \sigma^4 \rangle \leq \langle \sigma^2 \rangle \leq \langle \sigma \rangle \leq QD_{16},$$

where each composition factor is isomorphic to  $Z_2$ . Since  $Z_2$  is abelian, each composition factor is solvable, and so  $QD_{16}$  is solvable.  $\square$