

Dummit & Foote Ch. 3.1: Quotient Groups and Homomorphisms

Scott Donaldson

Aug. - Sep. 2023

Let G and H be groups.

1. (8/21/23)

Let $\varphi : G \rightarrow H$ be a homomorphism and let $E \leq H$. Prove that $\varphi^{-1}(E) \leq G$ (i.e., the preimage or pullback of a subgroup under a homomorphism is a subgroup). If $E \trianglelefteq H$ prove that $\varphi^{-1}(E) \trianglelefteq G$. Deduce that $\ker \varphi \trianglelefteq G$.

Proof. Let $x, y \in \varphi^{-1}(E) \subseteq G$. Suppose that $\varphi(x) = a, \varphi(y) = b, a, b \in E \leq H$. Since φ is a homomorphism, we have $\varphi(y^{-1}) = \varphi(y)^{-1} = b^{-1}$. Then:

$$\varphi(xy^{-1}) = \varphi(x)\varphi(y^{-1}) = \varphi(x)\varphi(y)^{-1} = ab^{-1} \in E,$$

which implies that $xy^{-1} \in \varphi^{-1}(E)$. It follows that, by the subgroup criterion, $\varphi^{-1}(E) \leq G$. \square

2. (8/23/23)

Let $\varphi : G \rightarrow H$ be a homomorphism of groups with kernel K and let $a, b \in \varphi(G)$. Let $X \in G/K$ be the fiber above a and Y be the fiber above b , i.e., $X = \varphi^{-1}(a), Y = \varphi^{-1}(b)$. Fix an element $x \in X$ (so $\varphi(x) = a$). Prove that if $XY = Z$ in the quotient group G/K and z is any member of Z , then there is some $y \in Y$ such that $xy = z$.

Proof. We know that, for any $x \in X, y \in Y$, $\varphi(x) = a$ and $\varphi(y) = b$. Since φ is a homomorphism, it follows that $\varphi(xy) = \varphi(x)\varphi(y) = ab$, and so the image of any element of $XY = Z$ under φ is $ab \in H$.

Next, consider the element $x^{-1}z \in G$, as well as its image under φ . Since φ is a homomorphism, we have $\varphi(x^{-1}) = \varphi(x)^{-1}$. So $\varphi(x^{-1}z) = \varphi(x^{-1})\varphi(z) = \varphi(x)^{-1}\varphi(z) = a^{-1}ab = b$. The set Y consists of all elements of G whose image under φ is b , and so we must have $x^{-1}z \in Y$.

Now if we fix some element $x \in X$, then for any $z \in Z$, we have $x^{-1}z \in Y$ such that its product with x is z : $xx^{-1}z = z$. \square

3. (8/23/23)

Let A be an abelian group and let B be a subgroup of A . Prove that A/B is abelian. Give an example of a non-abelian group G containing a proper normal subgroup N such that G/N is abelian.

Proof. Because A is abelian, all subgroups of A are normal, so A/B is well-defined for every $B \leq A$.

Let $C, D \in A/B$ with $C = cB$ and $D = dB$ for some $c, d \in A$. Then:

$$CD = (cB)(dB) = (cd)B = (dc)B = (dB)(cB) = DC,$$

which implies that A/B is abelian.

Now if we let G be the dihedral group D_8 , then G is non-abelian. Let N be the cyclic subgroup generated by $r : \{1, r, r^2, r^3\}$. The only coset of N is sN ; together these two sets cover G . Then $G/N = \{N, sN\}$. There is only one group of order 2 up to isomorphism, and it is abelian. Thus G/N is abelian. \square

4. (8/23/23)

Prove that in the quotient group G/N , $(gN)^\alpha = (g^\alpha)N$ for all $\alpha \in \mathbb{Z}$.

Proof. We start by induction: In the base case, $\alpha = 1$, we have $(gN)^1 = gN = (g^1)N$. Next, suppose that for some $\alpha > 1$, we have $(gN)^\alpha = (g^\alpha)N$. Then:

$$(gN)^{\alpha+1} = (gN)^\alpha gN = g^\alpha N \cdot gN = (g^{\alpha+1})N,$$

as desired. We have now proven that $(gN)^\alpha = (g^\alpha)N$ for $\alpha \geq 1$.

Next, consider $(gN)^\alpha (gN)^{-\alpha}$, where $\alpha \geq 1$. In the quotient group G/N , for any subset $X \in G/N$, we must have $X^\alpha X^{-\alpha} = N$ (the identity of G/N), so $(gN)^\alpha (gN)^{-\alpha} = N$. From above, $(gN)^\alpha = (g^\alpha)N$, so $(g^\alpha)N \cdot (gN)^{-\alpha} = N$. Also, from the operation on left cosets, we know that $N = (g^\alpha)N \cdot (g^{-\alpha})N$. Since both $(g^\alpha)N \cdot (gN)^{-\alpha} = N$ and $(g^\alpha)N \cdot (g^{-\alpha})N = N$, we must have $(gN)^{-\alpha} = (g^{-\alpha})N$. We have now proven for all nonzero integers.

Finally, we note that $(gN)^0 = N$ (the identity of G/N) and that $(g^0)N = eN = N$, so $(gN)^0 = (g^0)N$. This concludes the proof that $(gN)^\alpha = (g^\alpha)N$ for all $\alpha \in \mathbb{Z}$. \square

5. (8/23/23)

Use the preceding exercise to prove that the order of the element gN in G/N is n , where n is the smallest positive integer such that $g^n \in N$ (and gN has infinite order if no such positive integer exists). Give an example to show that the order of gN in G/N may be strictly smaller than the order of g in G .

Proof. Let $gN \in G/N$, and let n be the smallest positive integer such that $g^n \in N$. Suppose that $g^n = h \in N$.

From Exercise 4., $(gN)^n = (g^n)N = hN = N$ (because $h \in N$), so the order of gN must divide n .

Suppose (toward contradiction) that the order of gN is k , where $k < n$. Then $(gN)^k = (g^k)N = N$, which implies that g^k lies in N , contradicting our assumption that n is the smallest such positive integer. Therefore the order of gN is n .

If there is no positive integer n such that $g^n \in N$, then for all $k \in \mathbb{Z}^+$, we have $(gN)^k = (g^k)N \neq N$, so gN has infinite order.

As an example where $|gN| < |g|$, let $G = Z_9 = \langle x \rangle$ and let $N = \langle x^3 \rangle$. Because all cyclic groups are abelian, N is normal in G , and so G/N is well-defined. The quotient group G/N contains three elements: N, xN , and $(x^2)N$. The element $xN \in G/N$ has order 3: $(xN)^3 = (x^3)N = N$ (because $x^3 \in N$). However, the generating element $x \in G$ has order 9. \square

6. (8/24/23)

Define $\varphi : \mathbb{R}^\times \rightarrow \{\pm 1\}$ by letting $\varphi(x)$ be x divided by the absolute value of x . Describe the fibers of φ and prove that φ is a homomorphism.

Proof. We consider the two cases where $x < 0$ and $x > 0$ (0 is not an element of \mathbb{R}^\times). If $x > 0$, then $\varphi(x) = x/|x| = x/x = 1$. If $x < 0$, then $\varphi(x) = x/|x| = x/-x = -1$. Therefore the fiber above -1 is every negative real number and the fiber above 1 is every positive real number.

To show that φ is a homomorphism, we let $x, y \in \mathbb{R}^\times$ and again consider the different cases: Where x and y are both positive, where they are both negative, and where one is positive and the other negative.

If both x and y are positive, then $\varphi(x)\varphi(y) = 1 \cdot 1 = 1$ and $\varphi(xy) = \frac{xy}{|xy|} = \frac{xy}{xy} = 1$, so $\varphi(x)\varphi(y) = \varphi(xy)$.

If both x and y are negative, then $\varphi(x)\varphi(y) = -1 \cdot -1 = 1$ and $\varphi(xy) = \frac{xy}{|xy|} = \frac{xy}{xy} = 1$, so $\varphi(x)\varphi(y) = \varphi(xy)$.

Suppose x is positive and y is negative. Then $\varphi(x)\varphi(y) = 1 \cdot -1 = -1$ and $\varphi(xy) = \frac{xy}{|xy|} = \frac{xy}{-xy} = -1$, so $\varphi(x)\varphi(y) = \varphi(xy)$.

Thus, in every case of $x, y \in \mathbb{R}^\times$, we have $\varphi(x)\varphi(y) = \varphi(xy)$, and φ is thus a homomorphism. \square

7. (8/24/23)

Define $\pi : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $\pi((x, y)) = x + y$. Prove that π is a surjective homomorphism and describe the kernel and fibers of π geometrically.

Proof. First, to show that π is surjective, let $z \in \mathbb{R}$. Now $z = z + 0$, so $(z, 0)$ is an element of \mathbb{R}^2 such that $\pi((z, 0)) = z + 0 = z$.

Next, to show that π is a homomorphism, let $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$. We have $\pi((x_1, y_1) + (x_2, y_2)) = \pi((x_1 + x_2, y_1 + y_2)) = x_1 + x_2 + y_1 + y_2$, and $\pi((x_1, y_1)) + \pi((x_2, y_2)) = x_1 + y_1 + x_2 + y_2$. By the commutativity of addition in \mathbb{R} , these are equal to each other, and so π is a surjective homomorphism.

The kernel of π consists of all points $(x, y) \in \mathbb{R}^2$ such that $x + y = 0$, that is, the diagonal line running from the upper-left to the bottom-right of the Cartesian plane. Geometrically, the fibers of π are translations of this line, such that for any $z \in \mathbb{R}$, the fiber of π above z is the diagonal line intersecting both $(z, 0)$ and $(0, z)$. \square

8. (8/24/23)

Let $\varphi : \mathbb{R}^\times \rightarrow \mathbb{R}^\times$ be the map sending x to the absolute value of x . Prove that φ is a homomorphism and find the image of φ . Describe the kernel and the fibers of φ .

Proof. Let $x, y \in \mathbb{R}^\times$ (so $x \neq 0, y \neq 0$). If both x and y are positive or both are negative, then:

$$\varphi(xy) = |xy| = |x||y| = \varphi(x)\varphi(y),$$

and if x is positive and y is negative, then:

$$\varphi(xy) = |xy| = x(-y) = |x||y| = \varphi(x)\varphi(y),$$

so φ is a homomorphism.

The image of φ consists of every positive real number. The kernel of φ is the set $\{x \in \mathbb{R}^\times \mid |x| = 1\}$, that is, $\{\pm 1\}$. For a given element $z > 0$, the fiber of φ above z is the set $\{\pm z\}$. \square

9. (8/25/23)

Define $\varphi : \mathbb{C}^\times \rightarrow \mathbb{R}^\times$ by $\varphi(a + bi) = a^2 + b^2$. Prove that φ is a homomorphism and find the image of φ . Describe the kernel and the fibers of φ geometrically (as subsets of the plane).

Proof. To show that φ is a homomorphism, let $z_1 = a_1 + b_1i, z_2 = a_2 + b_2i \in \mathbb{C}^\times$. We calculate:

$$\begin{aligned} \varphi(z_1 z_2) &= \varphi((a_1 + b_1i)(a_2 + b_2i)) \\ &= \varphi((a_1 a_2 - b_1 b_2) + (a_1 b_2 + a_2 b_1)i) \\ &= (a_1 a_2 - b_1 b_2)^2 + (a_1 b_2 + a_2 b_1)^2 \\ &= a_1^2 a_2^2 - 2a_1 a_2 b_1 b_2 + b_1^2 b_2^2 + a_1^2 b_2^2 + 2a_1 a_2 b_1 b_2 + a_2^2 b_1^2 \\ &= a_1^2 a_2^2 + b_1^2 b_2^2 + a_1^2 b_2^2 + a_2^2 b_1^2, \text{ and} \\ \varphi(z_1) \varphi(z_2) &= \varphi(a_1 + b_1i) \varphi(a_2 + b_2i) = (a_1^2 + b_1^2)(a_2^2 + b_2^2) \\ &= a_1^2 a_2^2 + b_1^2 b_2^2 + a_1^2 b_2^2 + a_2^2 b_1^2, \end{aligned}$$

which proves that φ is a homomorphism.

The image of a complex number $a + bi$ under φ is $a^2 + b^2$, which is always non-negative because it is the sum of two non-negative numbers. Since both \mathbb{C}^\times and \mathbb{R}^\times exclude 0, the image of φ is therefore all positive real numbers.

The kernel of φ are those complex numbers whose image under φ is 1. Geometrically, φ is a map from a point in the complex plane to its length, or distance from zero. Therefore the kernel of φ is the unit circle in the complex plane. The fibers of a given positive real number x is the circle of radius x centered at the origin in the complex plane. \square

10. (8/28/23)

Let $\varphi : \mathbb{Z}/8\mathbb{Z} \rightarrow \mathbb{Z}/4\mathbb{Z}$ by $\varphi(\bar{a}) = \bar{a}$. Show that this is a well-defined, surjective homomorphism and describe its fibers and kernel explicitly (showing that φ is well-defined involves the fact that \bar{a} has a different meaning in the domain and range of φ).

Proof. The map φ is well-defined because it assigns to each member of $\mathbb{Z}/8\mathbb{Z}$ a single, unique element of $\mathbb{Z}/4\mathbb{Z}$. Let $a \in \{0, \dots, 7\}$ be equal to $\bar{a} \bmod 8$. Then we have $\varphi(\bar{a}) = \varphi(a)$. Further, φ assigns each $a \in \{0, \dots, 7\}$ to $a \bmod 4$; that is, it assigns 0 and 4 to 0, 1 and 5 to 1, 2 and 6 to 2, and 3 and 7 to 3. This also shows that φ is surjective, since each $\bar{a} \in \mathbb{Z}/4\mathbb{Z}$ (represented by $a = \bar{a} \bmod 4$) has a preimage in $\mathbb{Z}/8\mathbb{Z}$.

The kernel of φ is $\{0, 4\} \leq \mathbb{Z}/8\mathbb{Z}$, and the fiber of any $a \in \mathbb{Z}/4\mathbb{Z}$ is the tuple $\{a, a + 4\}$. \square

11. (8/28/23)

Let F be a field and let $G = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid a, b, c \in F, ac \neq 0 \right\} \leq GL_2(F)$.

- (a) Prove that the map $\varphi : \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mapsto a$ is a surjective homomorphism from G onto F^\times (recall that F^\times is the multiplicative group of nonzero elements in F). Describe the fibers and kernel of φ .

Proof. To show that φ is surjective, let $a \in F^\times$ (so $a \neq 0$). Then we have $\varphi\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}\right) = a$, so φ is onto.

Next, to show that it is a homomorphism, we note that:

$$\varphi\left(\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} d & e \\ 0 & f \end{pmatrix}\right) = \varphi\left(\begin{pmatrix} ad & ae + bf \\ 0 & cf \end{pmatrix}\right) = ad = \varphi\left(\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}\right)\varphi\left(\begin{pmatrix} d & e \\ 0 & f \end{pmatrix}\right),$$

so φ is also a homomorphism.

The kernel of φ is $\left\{\begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix} \mid b, c \in F, c \neq 0\right\}$, and the fiber of φ over a given element $a \in F^\times$ is $\left\{\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid b, c \in F, c \neq 0\right\}$. \square

- (b) Prove that the map $\psi : \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mapsto (a, c)$ is a surjective homomorphism from G onto $F^\times \times F^\times$. Describe the fibers and kernel of ψ .

Proof. To show that ψ is surjective, let $(a, c) \in F^\times \times F^\times$ (so $a, c \neq 0$). Then we have $\psi\left(\begin{pmatrix} a & 0 \\ 0 & c \end{pmatrix}\right) = (a, c)$, so ψ is onto.

Next, to show that it is a homomorphism, we note that:

$$\begin{aligned} \psi\left(\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} d & e \\ 0 & f \end{pmatrix}\right) &= \psi\left(\begin{pmatrix} ad & ae + bf \\ 0 & cf \end{pmatrix}\right) = (ad, cf) \\ &= (a, c)(d, f) = \psi\left(\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}\right)\psi\left(\begin{pmatrix} d & e \\ 0 & f \end{pmatrix}\right), \end{aligned}$$

so ψ is also a homomorphism.

The kernel of ψ is the preimage of $(1, 1)$, that is, $\left\{\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \mid b \in F\right\}$, and the fiber of ψ over a given element $(a, c) \in F^\times \times F^\times$ is $\left\{\begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid b \in F\right\}$. \square

- (c) Let $H = \left\{\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \mid b \in F\right\}$. Prove that H is isomorphic to the additive group F .

Proof. As usual, to show that H is isomorphic to the additive group F , we must show that there exists a bijective homomorphism $\varphi : H \rightarrow F$. Define φ by $\varphi\left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\right) = b$. We will show that it is an isomorphism.

First, φ is injective: Suppose that $\varphi\left(\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}\right) = \varphi\left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\right) = c$. Then we have $a = c$ and $b = c$, so the two matrices are the same, and φ is injective.

Next, φ is surjective: Let $b \in F$. Then we have $\varphi\left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\right) = b$.

Finally, φ is a homomorphism:

$$\varphi\left(\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\right) = \varphi\left(\begin{pmatrix} 1 & a+b \\ 0 & 1 \end{pmatrix}\right) = a+b = \varphi\left(\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}\right) + \varphi\left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\right).$$

\square

12. (8/30/23)

Let G be the additive group of real numbers, let H be the multiplicative group of complex numbers of absolute value 1 (the unit circle S^1 in the complex plane) and let $\varphi : G \rightarrow H$ be the homomorphism $\varphi : r \mapsto e^{2\pi ir}$. Draw the points on the real line which lie in the kernel of φ . Describe similarly the elements in the fibers of φ above the points -1 , i , and $e^{4\pi i/3}$ of H .

Proof. The kernel of φ is the set $\{r \in \mathbb{R} \mid e^{2\pi ir} = 1\}$. Recall that $e^{2\pi ir} = \cos 2\pi r + i \sin 2\pi r$, so the values of r for which $e^{2\pi ir} = 1$ are those where $\cos 2\pi r = 1$, that is, all of the integers.

We similarly obtain the fiber of φ above -1 by considering when $\cos 2\pi r = -1$, which occurs when $r = 1/2, 3/2, 5/2, \dots$, that is, $r \in \{n + \frac{1}{2} \mid n \in \mathbb{Z}\}$. For the fiber above i , we must have $\sin 2\pi r = 1$, which occurs when $r = 1/4, 5/4, 9/4, \dots$, that is, $r \in \{n + \frac{1}{4} \mid n \in \mathbb{Z}\}$. Finally, we have $4\pi/3 = \frac{2}{3} \cdot 2\pi$, so the fiber above $e^{4\pi i/3}$ is $\{n + \frac{2}{3} \mid n \in \mathbb{Z}\}$.

We can also write these as cosets of \mathbb{Z} , so the fibers are $\frac{1}{2} + \mathbb{Z}$, $\frac{1}{4} + \mathbb{Z}$, and $\frac{2}{3} + \mathbb{Z}$, respectively. \square

13. (8/31/23)

Repeat the preceding exercise with the map φ replaced by the map $\varphi : r \mapsto e^{4\pi ir}$.

Proof. In this case, the kernel of φ consists of values of r for which $e^{4\pi ir} = 1 \Rightarrow \cos 4\pi r = 1$. The period is now halved, so this occurs when $r \in \{1/2, 1, 3/2, \dots\}$; the kernel is $\{\frac{n}{2} \mid n \in \mathbb{Z}\}$.

The fiber of φ above -1 has $\cos 4\pi r = -1$, when $r = 1/4, 3/4, 5/4, \dots$, that is, $r \in \{\frac{1}{4} + \frac{n}{2} \mid n \in \mathbb{Z}\}$. Above i , we have $\sin 4\pi r = 1$, so $r \in \{\frac{1}{8}, \frac{5}{8}, \dots\}$, and the fiber is $\{\frac{1}{8} + \frac{n}{2} \mid n \in \mathbb{Z}\}$. Finally, above $4\pi/3$, the fiber is $\{\frac{1}{3} + \frac{n}{2} \mid n \in \mathbb{Z}\}$.

If we denote the kernel in this exercise as $\frac{1}{2}\mathbb{Z}$, then as cosets, the fibers are $\frac{1}{4} + \frac{1}{2}\mathbb{Z}$, $\frac{1}{8} + \frac{1}{2}\mathbb{Z}$, and $\frac{1}{3} + \frac{1}{2}\mathbb{Z}$, respectively. \square