

6.5

Work done by a  
variable force along  
a line:

Def<sup>n</sup>: The work  
done by a variable  
force  $F(x)$  in moving

Feb 4-11:39 AM

an object along the  
x-axis from  
 $x=a$  to  $x=b$  is

$$W = \int_a^b F(x) dx$$

Feb 4-11:42 AM

(Ex) When a  
particle is located  
at a distance  $x$  m  
from the origin,  
a force of  $x^2 + 2x$   
pounds acts on it.  
How much work is

Feb 4-11:43 AM

done in moving it  
from  $x=1$  to  $x=3$  m.

$$W = \int_1^3 F(x) dx$$

$$= \int_1^3 (x^2 + 2x) dx$$

Feb 4-11:44 AM

$$= \left. \frac{x^3}{3} + \frac{2x^2}{2} \right|_1^3$$

$$= \left( \frac{3^3}{3} + 3^2 \right) - \left( \frac{1^3}{3} + 1^2 \right)$$

$$= \frac{50}{3} \text{ Nm} = \boxed{\frac{50}{3} \text{ J}}$$

Feb 4-11:45 AM

Hooke's law for  
springs:

Hooke's law  
states that the  
force required to  
maintain a spring  
stretched or compressed

Feb 4-11:47 AM

$x$  units from its natural length is proportion to  $x$ .

$$F = Kx$$

Where  $K$  is a positive constant called spring constant.

Feb 4-11:48 AM

(Ex) Find the force required to stretch a 20 cm long spring, with a spring constant of 100 N/m, to a length of 21 cm.

Feb 4-11:51 AM

$$K = 100 \text{ N/m}$$

$$x = 21 - 20 = 1 \text{ cm} = 0.01 \text{ m}$$

$$F = Kx$$

$$F = (100)(0.01)$$

$$F = 1 \text{ N}$$

Feb 4-11:54 AM

(Ex) A spring has a natural length of 2 ft. A body of 10 lb hanging on the spring stretches it to the total length of 2.4 ft.

Feb 4-11:55 AM


(a) Find the spring constant.

(b) How far beyond its natural length would a body of 50 lb stretch the spring?

Feb 4-11:57 AM

(c) How much work is required to stretch the spring from its natural length to a length of 3 ft?

(a)  $F = Kx$



Feb 4-11:58 AM

$$\Rightarrow 10 = K(0.4)$$

$$\Rightarrow K = 25 \text{ lb/ft}$$

(b)  $F = Kx$   
 $50 = 25(x)$   
 $x = 2 \text{ ft}$

Feb 4-12:01 PM

(c)  $W = \int_0^1 F(x) dx$   
 $= \int_0^1 25x dx$   
 $= 25 \left. \frac{x^2}{2} \right|_0^1$

Feb 4-12:05 PM

$$= \frac{25}{2} \text{ lb ft}$$

Feb 4-12:05 PM