

Trilinear Higgs boson coupling variations for di-Higgs production with full NLO QCD predictions in Powheg

G. Heinrich¹, S. Jones², M. Kerner³, G. Luisoni¹ and L. Scyboz¹

¹ Max-Planck-Institut für Physik, Föhringer Ring 6, 80805 München, Germany

² Theoretical Physics Department, CERN, Geneva, Switzerland

³ Physik-Institut, Universität Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

E-mail: gudrun@mpp.mpg.de, s.jones@cern.ch, mkerner@physik.uzh.ch, luisonig@gmail.com, scyboz@mpp.mpg.de

Abstract. The couplings of the Higgs boson to other particles are increasingly well measured by the ATLAS and CMS experiments. The Higgs boson trilinear self-coupling however is still largely unconstrained, mainly due to the low cross-section for Higgs boson pair production. We present inclusive and differential results for the NLO QCD corrections to Higgs boson pair production with the full top-quark mass dependence, where the Higgs trilinear coupling is varied to non-SM values. The fixed-order calculation is supplemented by parton showering within the Powheg-B0X-V2 event generator, and both Pythia8 and Herwig7 parton-shower algorithms are implemented in a preliminary study of shower effects.

1. Introduction

Impressive experimental constraints have been set on the Higgs boson couplings to vector bosons and heavy fermions [1, 2, 3, 4]. The Higgs potential, in contrast, leaves more room for New Physics. In particular, the Higgs boson trilinear self-coupling λ can be experimentally constrained by exclusion limits on Higgs boson pair production $pp \rightarrow hh$ [5, 6], where the best limit on $\kappa_\lambda = \lambda/\lambda_{\text{SM}}$ is currently given by ATLAS with $-5.0 < \kappa_\lambda < 12.0$ at 95% confidence level. Higher-order corrections to Higgs pair production were first calculated in the heavy top-quark mass limit (HTL) $m_t \rightarrow \infty$, where the top-quark degrees of freedom are integrated out [7, 8, 9, 10]. The NLO QCD corrections with the full top-quark mass dependence were only computed more recently [11, 12, 13]. The latter are based on numerical evaluations of the two-loop contribution to $gg \rightarrow hh$. For non-SM values of the Higgs couplings, results were computed at NLO QCD in the full theory for a class of extensions of the SM in Ref. [14].

In the following, an implementation of the full NLO QCD corrections into the Powheg-B0X-V2 event generator [15, 16, 17] is presented. In this framework, the Higgs trilinear self-coupling can be varied, as well as the top-Higgs Yukawa coupling. Total cross sections are computed for $\sqrt{s} = 13, 14$ and 27 TeV at the (HE-)LHC. Differential results are shown for $\sqrt{s} = 14$ TeV. The fixed-order calculation is then matched to both Pythia8 [18] and Herwig7 [19, 20] parton showers. For a more detailed description, the reader is referred to Ref. [21].

2. Description of the calculation

The calculation is based on the setup presented in Ref. [22] for the case of the SM. The leading-order amplitude has been computed analytically. The real-emission contributions were

implemented using an interface [23] between the **Powheg-BOX** and **GoSAM** [24, 25], where the reduction of the one-loop amplitude has been performed with **Ninja** [26], using master integrals from **golem95C** [27, 28], **OneLOop** [29] and **VBFNLO** [30, 31]. The two-loop amplitude for the full virtual contribution was adapted from Refs. [11, 12], which used an extension of the **GoSAM** package to two loops [32]. There, the integral reduction was performed with **REDUZE2** [33], and the integrals were numerically evaluated with **SECDEC3** [34]. For a faster convergence, the integration was performed within a Quasi-Monte-Carlo implementation using a rank-1 shifted lattice rule [35, 36]. The integrals were computed with 16 dual **NVIDIA TESLA K20X** GPUs. The top-quark and Higgs masses have been set to $m_t = 173$ GeV and $m_h = 125$ GeV. Thus, the integrals depend only on the two Mandelstam invariants \hat{s} and \hat{t} .

A grid for the two-loop amplitude was constructed in both variables using 5291 pre-sampled phase-space points. We split the amplitude in two contributions: diagrams containing the trilinear Higgs coupling are called *triangle-like*, and those that do not are called *box-like* (see Fig. 1 for two diagrams at NLO QCD).



Figure 1. Triangle-like (left) and box-like (right) diagrams contribute to the full amplitude. The former contain the Higgs self-coupling, while the latter do not.

At any order in QCD, the squared matrix-element can thus be written as a second-order polynomial in λ :

$$M_\lambda \equiv |\mathcal{M}_\lambda|^2 = \mathcal{M}_B^* \mathcal{M}_B + \lambda (\mathcal{M}_B^* \mathcal{M}_T + \mathcal{M}_T^* \mathcal{M}_B) + \lambda^2 \mathcal{M}_T^* \mathcal{M}_T. \quad (1)$$

The two-loop amplitude for an arbitrary value of λ can be reconstructed from the squared matrix-element computed for three different values of λ . In our case, we chose $\kappa_\lambda = \lambda_{\text{BSM}}/\lambda_{\text{SM}} \in \{-1, 0, 1\}$. A new grid is generated at runtime for the user-defined value of λ , where the amplitude for each pre-sampled phase-space point is calculated as:

$$M_\lambda = M_0 \cdot (1 - \lambda^2) + \frac{M_1}{2} \cdot (\lambda + \lambda^2) + \frac{M_{-1}}{2} \cdot (-\lambda + \lambda^2). \quad (2)$$

The grid produced for the two-loop amplitude is fed to an interpolation framework, which interfaces the result at *any* phase-space point $M_\lambda(\hat{s}, \hat{t})$ to **Powheg**.

3. Total and differential cross-sections for variations of the trilinear coupling

The results given below are produced using the **PDF4LHC15_nlo_30_pdfas** sets [37, 38, 39, 40] interfaced to **Powheg** via **LHAPDF6** [41], with the corresponding value of α_s . The top-quark mass is renormalised in the on-shell scheme and is set to $m_t = 173$ GeV, as in the virtual amplitude. The mass of the Higgs boson is fixed to $m_h = 125$ GeV, and the top-quark and Higgs widths are set to zero. Jets are clustered using the anti- k_T algorithm [42] as implemented in **FastJet** [43, 44], with a jet distance parameter of $R = 0.4$ and a minimum transverse momentum requirement of $p_T = 20$ GeV. The central renormalisation and factorisation scales are set to $\mu_R = \mu_F = \mu_0 = m_{\text{hh}}/2$. Scale uncertainties are estimated by 3-point variations $\mu_R = \mu_F = c \mu_0$, with $c \in \{0.5, 1.0, 2.0\}$.

Total cross-sections for Higgs pair production at the (HE-)LHC are shown in Table 1, for centre-of-mass energies of $\sqrt{s} = 13, 14$ and 27 TeV and different values of the Higgs self-coupling $\kappa_\lambda = \lambda_{\text{BSM}}/\lambda_{\text{SM}}$. They are accompanied by their relative scale uncertainties, which are of the order $\mathcal{O}(10-20\%)$. Notably, the K -factors at 14 TeV show a sizeable dependence on the trilinear coupling κ_λ . In the HTL at NLO QCD, Ref. [45] suggested a variation of the K -factors with κ_λ of the order $\mathcal{O}(2-3\%)$. In the full theory, the K -factors are found to vary between 1.56 and 2.15 for variations of the trilinear coupling $-5 \leq \kappa_\lambda \leq 12$, see Fig. 2.

$\lambda_{\text{BSM}}/\lambda_{\text{SM}}$	$\sigma_{\text{NLO}}@13\text{TeV}$ [fb]	$\sigma_{\text{NLO}}@14\text{TeV}$ [fb]	$\sigma_{\text{NLO}}@27\text{TeV}$ [fb]	K-factor@14TeV
-1	$116.71^{+16.4\%}_{-14.3\%}$	$136.91^{+16.4\%}_{-13.9\%}$	$504.9^{+14.1\%}_{-11.8\%}$	1.86
0	$62.51^{+15.8\%}_{-13.7\%}$	$73.64^{+15.4\%}_{-13.4\%}$	$275.29^{+13.2\%}_{-11.3\%}$	1.79
1	$27.84^{+11.6\%}_{-12.9\%}$	$32.88^{+13.5\%}_{-12.5\%}$	$127.7^{+11.5\%}_{-10.4\%}$	1.66
2	$12.42^{+13.1\%}_{-12.0\%}$	$14.75^{+12.0\%}_{-11.8\%}$	$59.10^{+10.2\%}_{-9.7\%}$	1.56
2.4	$11.65^{+13.9\%}_{-12.7\%}$	$13.79^{+13.5\%}_{-12.5\%}$	$53.67^{+11.4\%}_{-10.3\%}$	1.65
3	$16.28^{+16.2\%}_{-15.3\%}$	$19.07^{+17.1\%}_{-14.1\%}$	$69.84^{+14.6\%}_{-12.1\%}$	1.90
5	$81.74^{+20.0\%}_{-15.6\%}$	$95.22^{+19.7\%}_{-11.5\%}$	$330.61^{+17.4\%}_{-13.6\%}$	2.14

Table 1. Total cross-sections for Higgs boson pair production at NLO QCD at (HE-)LHC for centre-of-mass energies of $\sqrt{s} = 13, 14$ and 27 TeV. The scale uncertainties are given in percent.




Figure 2. The dependence of the K -factor on the trilinear Higgs self-couplings κ_λ is given at $\sqrt{s} = 14$ TeV in the full theory.

In Fig. 3, distributions of the invariant mass m_{hh} of the Higgs boson pair system are displayed for different values of κ_λ . They exhibit a characteristic dip around $m_{hh} \sim 350$ GeV for values of the trilinear coupling around $\kappa_\lambda = 2.4$. This value of the trilinear self-coupling corresponds to a maximally destructive interference between triangle-like and box-like diagrams. For $\kappa_\lambda = 1$, the maximal destructive interference happens at the hh production threshold and therefore does not manifest itself as a dip, while for κ_λ values larger than ~ 3 the triangle-type contributions start to dominate.




Figure 3. Distributions of the Higgs boson pair invariant mass m_{hh} for various values of κ_λ at $\sqrt{s} = 14$ TeV. The uncertainty bands are from scale variations as described in the text.

Note that since the contributions can be separated in triangle- and box-like diagrams, the top-Higgs Yukawa coupling y_t can easily be varied within the same code. A non-SM value of y_t yields in Eq. (1):

$$|\mathcal{M}_\lambda|^2 = y_t^4 \left[\mathcal{M}_B^* \mathcal{M}_B + \frac{\kappa_\lambda}{y_t} (\mathcal{M}_B^* \mathcal{M}_T + \mathcal{M}_T^* \mathcal{M}_B) + \left(\frac{\kappa_\lambda}{y_t} \right)^2 \mathcal{M}_T^* \mathcal{M}_T \right]. \quad (3)$$

The cross-section can be computed by setting κ_λ in the code to the desired value of the ratio κ_λ/y_t , and rescaling the result by an overall factor y_t^4 . For example, $\sigma(y_t = 1.2, \kappa_\lambda = 1) = (1.2)^4 \sigma(y_t = 1, \kappa_\lambda = 1/1.2)$. Fig. 4 shows the distribution of m_{hh} for values of the top-Higgs Yukawa coupling that are still not experimentally excluded [4].




Figure 4. The distribution of the Higgs boson pair invariant mass m_{hh} for values of the top-Higgs Yukawa coupling $y_t \in \{0.8, 1, 1.2\}$.

4. Parton-shower matched results

We now consider NLO distributions matched to a parton shower. The Les Houches Events (LHE) [46] files produced by Powheg are used as input to the Pythia8.235 and Herwig7.1.4

parton showers. In the case of **Herwig7**, both the default angular-ordered \tilde{q} and the dipole showers are compared. The radiation-regulating **hdamp** parameter in **Powheg** is set to **hdamp** = 250 GeV. Multiple-parton interactions and hadronisation are switched off. The default tunes are used for both parton showers.

Fig. 5 displays the transverse momentum of the Higgs boson pair p_T^{hh} and the separation between the two Higgs bosons $\Delta R^{\text{hh}} = \sqrt{(\eta_1 - \eta_2)^2 + (\phi_1 - \phi_2)^2}$. Considering first the distribution of p_T^{hh} , both **Herwig7** parton showers (PH7- \tilde{q} and PH7-dipole) generate similar results and reproduce the fixed-order NLO prediction in the far- p_T^{hh} range. In contrast, **Pythia8** agrees with **Herwig7** only for small transverse momenta, while it produces much harder radiation in the tail of the distribution. The same comments apply to the ΔR^{hh} observable in the region $0 < \Delta R^{\text{hh}} < \pi$ where shower contributions are important. Large parton-shower matching uncertainties in Higgs boson pair production have already been discussed in Ref. [47].




Figure 5. The transverse momentum p_T^{hh} of the Higgs boson pair and the separation between the two Higgs bosons ΔR^{hh} are shown for the fixed-order NLO calculation and three parton showers, in the $\kappa_\lambda = 1$ case.

5. Conclusion

We have presented a new program package for Higgs boson pair production at NLO QCD with full top-quark mass dependence. In this package, the trilinear Higgs self-coupling can be varied explicitly. Within the same code, simultaneous variations of the top-Higgs Yukawa coupling can also be produced. The public code for the **Powheg-BOX-V2** event generator can be found at the website <http://powhegbox.mib.infn.it> in the **User-Processes-V2/ggHH** subdirectory. In addition, approximations related to the heavy top limit (HTL) can be enabled for comparison purposes. We have found that the full m_t -dependent NLO QCD corrections lead to K -factors which exhibit a sizeable dependence on the value of the trilinear Higgs self-coupling, which is not present in the HTL. We have compared fixed-order predictions at NLO QCD to parton-shower matched results. Both the **Pythia8** and **Herwig7** (\tilde{q} and dipole) parton showers can be matched directly to LHE files produced by **Powheg**. Full particle-level events can be produced with our framework, including Higgs boson decays and hadronisation.

Acknowledgments

This research was supported in part by the COST Action CA16201 ‘Particleface’ of the European Union, and by the Swiss National Science Foundation (SNF) under grant number 200020-175595.

References

- [1] Aad G *et al.* (ATLAS, CMS) 2016 *JHEP* **08** 045 (*Preprint* 1606.02266)

- [2] Aaboud M *et al.* (ATLAS) 2018 *JHEP* **03** 095 (*Preprint* 1712.02304)
- [3] Sirunyan A M *et al.* (CMS) 2018 *Submitted to: Eur. Phys. J.* (*Preprint* 1809.10733)
- [4] Sirunyan A M *et al.* (CMS) 2018 *Submitted to: Phys. Lett.* (*Preprint* 1812.06504)
- [5] Sirunyan A M *et al.* (CMS) 2019 *Phys. Rev. Lett.* **122** 121803 (*Preprint* 1811.09689)
- [6] Aad G *et al.* (ATLAS) 2019 (*Preprint* 1906.02025)
- [7] Dawson S, Dittmaier S and Spira M 1998 *Phys. Rev.* **D58** 115012 (*Preprint* hep-ph/9805244)
- [8] de Florian D and Mazzitelli J 2013 *Phys. Rev. Lett.* **111** 201801 (*Preprint* 1309.6594)
- [9] Grigo J, Melnikov K and Steinhauser M 2014 *Nucl. Phys.* **B888** 17–29 (*Preprint* 1408.2422)
- [10] de Florian D, Grazzini M, Hanga C, Kallweit S, Lindert J M, Maierhöfer P, Mazzitelli J and Rathlev D 2016 *JHEP* **09** 151 (*Preprint* 1606.09519)
- [11] Borowka S, Greiner N, Heinrich G, Jones S, Kerner M, Schlenk J, Schubert U and Zirke T 2016 *Phys. Rev. Lett.* **117** 012001 [Erratum: *Phys. Rev. Lett.* 117, no. 7, 079901 (2016)] (*Preprint* 1604.06447)
- [12] Borowka S, Greiner N, Heinrich G, Jones S, Kerner M, Schlenk J and Zirke T 2016 *JHEP* **10** 107 (*Preprint* 1608.04798)
- [13] Baglio J, Campanario F, Glaus S, Mühlleitner M, Spira M and Streicher J 2018 (*Preprint* 1811.05692)
- [14] Buchalla G, Capozzi M, Celis A, Heinrich G and Scyboz L 2018 *JHEP* **09** 057 (*Preprint* 1806.05162)
- [15] Nason P 2004 *JHEP* **11** 040 (*Preprint* hep-ph/0409146)
- [16] Frixione S, Nason P and Oleari C 2007 *JHEP* **11** 070 (*Preprint* 0709.2092)
- [17] Alioli S, Nason P, Oleari C and Re E 2010 *JHEP* **06** 043 (*Preprint* 1002.2581)
- [18] Sjostrand T, Ask S, Christiansen J R, Corke R, Desai N, Ilten P, Mrenna S, Prestel S, Rasmussen C O and Skands P Z 2015 *Comput. Phys. Commun.* **191** 159–177 (*Preprint* 1410.3012)
- [19] Bellm J *et al.* 2016 *Eur. Phys. J.* **C76** 196 (*Preprint* 1512.01178)
- [20] Bellm J *et al.* 2017 (*Preprint* 1705.06919)
- [21] Heinrich G, Jones S P, Kerner M, Luisoni G and Scyboz L 2019 (*Preprint* 1903.08137)
- [22] Heinrich G, Jones S P, Kerner M, Luisoni G and Vryonidou E 2017 *JHEP* **08** 088 (*Preprint* 1703.09252)
- [23] Luisoni G, Nason P, Oleari C and Tramontano F 2013 *JHEP* **10** 083 (*Preprint* 1306.2542)
- [24] Cullen G, Greiner N, Heinrich G, Luisoni G, Mastrolia P, Ossola G, Reiter T and Tramontano F 2012 *Eur. Phys. J.* **C72** 1889 (*Preprint* 1111.2034)
- [25] Cullen G *et al.* 2014 *Eur. Phys. J.* **C74** 3001 (*Preprint* 1404.7096)
- [26] Peraro T 2014 *Comput. Phys. Commun.* **185** 2771–2797 (*Preprint* 1403.1229)
- [27] Binoth T, Guillet J P, Heinrich G, Pilon E and Reiter T 2009 *Comput. Phys. Commun.* **180** 2317–2330 (*Preprint* 0810.0992)
- [28] Cullen G, Guillet J P, Heinrich G, Kleinschmidt T, Pilon E *et al.* 2011 *Comput. Phys. Commun.* **182** 2276–2284 (*Preprint* 1101.5595)
- [29] van Hameren A 2011 *Comput. Phys. Commun.* **182** 2427–2438 (*Preprint* 1007.4716)
- [30] Arnold K *et al.* 2009 *Comput. Phys. Commun.* **180** 1661–1670 (*Preprint* 0811.4559)
- [31] Baglio J *et al.* 2014 (*Preprint* 1404.3940)
- [32] Jones S P 2016 *PoS* **LL2016** 069 (*Preprint* 1608.03846)
- [33] von Manteuffel A and Studerus C 2012 (*Preprint* 1201.4330)
- [34] Borowka S, Heinrich G, Jones S P, Kerner M, Schlenk J and Zirke T 2015 *Comput. Phys. Commun.* **196** 470–491 (*Preprint* 1502.06595)
- [35] Borowka S, Heinrich G, Jahn S, Jones S P, Kerner M and Schlenk J 2018 *Comp. Phys. Comm.* (*Preprint* 1811.11720)
- [36] Jones S P 2019 *These Proceedings*
- [37] Butterworth J *et al.* 2015 (*Preprint* 1510.03865)
- [38] Dulat S, Hou T J, Gao J, Guzzi M, Huston J, Nadolsky P, Pumplin J, Schmidt C, Stump D and Yuan C P 2016 *Phys. Rev.* **D93** 033006 (*Preprint* 1506.07443)
- [39] Harland-Lang L A, Martin A D, Motylinski P and Thorne R S 2015 *Eur. Phys. J.* **C75** 204 (*Preprint* 1412.3989)
- [40] Ball R D *et al.* (NNPDF) 2015 *JHEP* **04** 040 (*Preprint* 1410.8849)
- [41] Buckley A, Ferrando J, Lloyd S, Nordström K, Page B, Rüfenacht M, Schönherr M and Watt G 2015 *Eur. Phys. J.* **C75** 132 (*Preprint* 1412.7420)
- [42] Cacciari M, Salam G P and Soyez G 2008 *JHEP* **04** 063 (*Preprint* 0802.1189)
- [43] Cacciari M and Salam G P 2006 *Phys. Lett.* **B641** 57–61 (*Preprint* hep-ph/0512210)
- [44] Cacciari M, Salam G P and Soyez G 2012 *Eur. Phys. J.* **C72** 1896 (*Preprint* 1111.6097)
- [45] Gröber R, Mühlleitner M, Spira M and Streicher J 2015 *JHEP* **09** 092 (*Preprint* 1504.06577)
- [46] Alwall J *et al.* 2007 *Comput. Phys. Commun.* **176** 300–304 (*Preprint* hep-ph/0609017)
- [47] Jones S and Kuttimalai S 2018 *JHEP* **02** 176 (*Preprint* 1711.03319)