

AN ALGORITHM FOR THE NUMERICAL SOLUTION OF DIFFERENTIAL EQUATIONS OF FRACTIONAL ORDER

ZAID M. ODIBAT* AND SHAHER MOMANI

ABSTRACT. We present and discuss an algorithm for the numerical solution of initial value problems of the form $D_*^\alpha y(t) = f(t, y(t))$, $y(0) = y_0$, where $D_*^\alpha y$ is the derivative of y of order α in the sense of Caputo and $0 < \alpha \leq 1$. The algorithm is based on the fractional Euler's method which can be seen as a generalization of the classical Euler's method. Numerical examples are given and the results show that the present algorithm is very effective and convenient.

AMS Mathematics Subject Classification : 65L05, 26A33

Key words and phrases : Generalized Taylor's formula; fractional Euler's method; fractional differential equation; Caputo fractional derivative.

1. Introduction

In this paper we introduce an algorithm for the numerical solution of initial value problems of the form

$$D_*^\alpha y(t) = f(t, y(t)), \quad y(0) = y_0, \quad 0 < \alpha \leq 1, \quad (1)$$

where D_*^α denotes the Caputo fractional differential operator [1]. The development of this algorithm is motivated by a few classical and many recent applications of fractional differential equations. Among the classical problems we mention areas like the modelling of the behavior of viscoelastic materials in mechanics [2]. More recently fractional calculus has been applied to continuum and statistical mechanics for viscoelasticity problems, Brownian motion and fractional diffusion-wave equations and many physical phenomena [1-12].

Most nonlinear fractional differential equations do not have analytic solutions, so approximations and numerical techniques must be used [13-19]. The decomposition method [20-28] and the variational iteration method [29-45] are relatively new approaches to provide an analytical approximate solution to linear

Received March 18, 2007. Revised October 19, 2007. *Corresponding author.
 © 2008 Korean SIGCAM and KSCAM .

and nonlinear problems, and they are particularly valuable as tools for scientists and applied mathematicians, because they provide immediate and visible symbolic terms of analytic solutions, as well as numerical approximate solutions to both linear and nonlinear differential equations. The application of the two methods is successfully extended to obtain an analytical approximate solutions to linear and nonlinear differential equations of fractional order [24-28,32,45-47]. A comparison between the variational iteration method and Adomian decomposition method for solving fractional differential equations is given in [46,47]. The fact that the variational iteration method solves nonlinear equations without using Adomian polynomials can be considered as an advantage of this method over Adomian decomposition method.

A few numerical methods for fractional differential equations have been presented in the literature [13-19]. However many of these methods are used for very specific types of differential equations, often just linear equations or even smaller classes. Recently, Diethelm et al. [19] introduced a method for the numerical solution of the nonlinear fractional differential equation (1) which can be seen as a generalization of the classical Adams-Bashforth-Moulton scheme for first order differential equations.

The structure of this paper is as follows. We begin by introducing some necessary definitions and mathematical preliminaries of the fractional calculus theory which are required for establishing our results. In sections 3 and 4, we introduce the modified trapezoidal rule and a new generalization of Taylor's formula that involves Caputo derivatives, respectively. In section 5, we derive the fractional Euler's method that is a generalization of the classical Euler's method for the numerical solution of ordinary differential equations. The algorithm itself is presented in details in section 6. In section 7, we present three examples to show the efficiency and the simplicity of the algorithm. Finally, in section 8, we give the conclusions.

2. Basic definitions

For the concept of fractional derivative we will adopt Caputo's definition which is a modification of the Riemann-Liouville definition and has the advantage of dealing properly with initial value problems in which the initial conditions are given in terms of the field variables and their integer order which is the case in most physical processes.

Definition 1. A real function $f(x)$, $x > 0$, is said to be in the space C_μ , $\mu \in \mathbb{R}$ if there exists a real number $p(> \mu)$, such that $f(x) = x^p f_1(x)$, where $f_1(x) \in C[0, \infty)$, and it is said to be in the space C_μ^m iff $f^{(m)} \in C_\mu$, $m \in \mathbb{N}$.

Definition 2. The Riemann-Liouville fractional integral operator of order $\alpha \geq 0$, of a function $f \in C_\mu$, $\mu \geq -1$, is defined as

$$J^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \quad \alpha > 0, \quad x > 0,$$

$$J^0 f(x) = f(x).$$

Properties of the operator J^α can be found in [3,4,8], we mention only the following:

For $f \in C_\mu, \mu \geq -1, \alpha, \beta \geq 0$ and $\gamma > -1$:

$$(1) J^\alpha J^\beta f(x) = J^{\alpha+\beta} f(x), \quad (2) J^\alpha J^\beta f(x) = J^\beta J^\alpha f(x),$$

$$(3) J^\alpha x^\gamma = \frac{\Gamma(\gamma+1)}{\Gamma(\alpha+\gamma+1)} x^{\alpha+\gamma}.$$

The Riemann-Liouville derivative has certain disadvantages when trying to model real-world phenomena with fractional differential equations. Therefore, we shall introduce a modified fractional differential operator D_*^α proposed by M. Caputo in his work on the theory of viscoelasticity [1]. Caputo's definition, which is a modification of the Riemann-Liouville definition, has the advantage of dealing properly with initial value problems in which the initial conditions are given in terms of the field variables and their integer order which is the case in most physical processes.

Definition 3. The fractional derivative of $f(x)$ in the Caputo sense is defined as

$$D_*^\alpha f(x) = J^{m-\alpha} D^m f(x) = \frac{1}{\Gamma(m-\alpha)} \int_0^x (x-t)^{m-\alpha-1} f^{(m)}(t) dt,$$

for $m-1 < \alpha \leq m, m \in \mathbb{N}, x > 0, f \in C_{-1}^m$.

Also, we need here two of its basic properties:

Lemma 1. If $m-1 < \alpha \leq m, m \in \mathbb{N}$ and $f \in C_\mu^m, \mu \geq -1$, then

$$D_*^\alpha J^\alpha f(x) = f(x), \quad (2)$$

$$J^\alpha D_*^\alpha f(x) = f(x) - \sum_{k=0}^{m-1} f^{(k)}(0^+) \frac{x^k}{k!}, \quad x > 0. \quad (3)$$

3. Modified trapezoidal rule

In this section, we present a review of the modified trapezoidal rule, which is introduced in [48]. This rule is used to approximate the fractional integral $J^\alpha f(t)$ by a weighted sum of function values at specified points. Suppose that the interval $[0, a]$ is subdivided into k subintervals $[t_j, t_{j+1}]$ of equal width $h = a/k$ by using the nodes $t_j = jh$, for $j = 0, 1, \dots, k$. The modified trapezoidal rule

$$T(f, h, \alpha) = ((k-1)^{\alpha+1} - (k-\alpha-1)k^\alpha) \frac{h^\alpha f(0)}{\Gamma(\alpha+2)} + \frac{h^\alpha f(a)}{\Gamma(\alpha+2)} \\ + \sum_{j=1}^{k-1} ((k-j+1)^{\alpha+1} - 2(k-j)^{\alpha+1} + (k-j-1)^{\alpha+1}) \frac{h^\alpha f(t_j)}{\Gamma(\alpha+2)}, \quad (4)$$

is an approximation to the fractional integral

$$(J^\alpha f(t))(a) = T(f, h, \alpha) - E_T(f, h, \alpha), \quad a > 0, \quad \alpha > 0. \quad (5)$$

Furthermore, if $f(t) \in \mathbf{C}^2[0, a]$, there is a constant C_α depending only on α so that the error term $E_T(f, h, \alpha)$ has the form

$$|E_T(f, h, \alpha)| \leq C_\alpha \|f''\|_\infty a^\alpha h^2 = O(h^2). \quad (6)$$

It is clear that if $\alpha = 1$, then the modified trapezoidal rule (4) reduces to the classical trapezoidal rule. This rule is simple for computational performance for all values of α and h . For more details about the rule and its applications, we refer the reader to [48].

4. Generalized Taylor's formula

In this section we introduce a new generalization of Taylor's formula that involves Caputo fractional derivatives. This generalization is presented in [49]. We begin by introducing the generalized mean value theorem.

Theorem 1. (Generalized mean value theorem) *Suppose that $f(x) \in \mathbf{C}[0, a]$ and $D_*^\alpha f(x) \in \mathbf{C}(0, a]$, for $0 < \alpha \leq 1$. Then we have*

$$f(x) = f(0+) + \frac{1}{\Gamma(\alpha)} (D_*^\alpha f)(\xi) \cdot x^\alpha, \quad (7)$$

with $0 \leq \xi \leq x$, $\forall x \in (0, a]$.

Proof. From the definitions of the Riemann-Liouville fractional integral operator and the Caputo fractional derivative operator, we have

$$(J^\alpha D_*^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} (D_*^\alpha f)(t) dt. \quad (8)$$

Using the integral mean value theorem, we get

$$(J^\alpha D_*^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} (D_*^\alpha f)(\xi) \int_0^x (x-t)^{\alpha-1} dt = \frac{1}{\Gamma(\alpha)} (D_*^\alpha f)(\xi) \cdot x^\alpha, \quad (9)$$

for $0 \leq \xi \leq x$. On the other hand, from equation (3), we have

$$(J^\alpha D_*^\alpha f)(x) = f(x) - f(0+). \quad (10)$$

So, from (9) and (10), the generalized mean value theorem (7) is obtained. \square

In case of $\alpha = 1$, the generalized mean value theorem reduces to the classical mean value theorem. Before we present the generalized Taylor's formula in the Caputo sense, we need the following relation.

Theorem 2. *Suppose that $D_*^{n\alpha} f(x)$, $D_*^{(n+1)\alpha} f(x) \in \mathbf{C}(0, a]$, for $0 < \alpha \leq 1$. Then we have*

$$(J^{n\alpha} D_*^{n\alpha} f)(x) - (J^{(n+1)\alpha} D_*^{(n+1)\alpha} f)(x) = \frac{x^{n\alpha}}{\Gamma(n\alpha + 1)} (D_*^{n\alpha} f)(0+), \quad (11)$$

where

$$D_*^{n\alpha} = D_*^\alpha D_*^\alpha \dots D_*^\alpha \quad (n - \text{times}).$$

Proof. The proof can be obtained using the properties of the Riemann-Liouville fractional integral operator and the Caputo fractional derivative operator and the relation:

$$\begin{aligned} (J^{n\alpha} D_*^{n\alpha} f)(x) - (J^{(n+1)\alpha} D_*^{(n+1)\alpha} f)(x) &= J^{n\alpha} \left((D_*^{n\alpha} f)(x) - (J^\alpha D_*^\alpha)(D_*^{n\alpha} f)(x) \right) \\ &= J^{n\alpha} (D_*^{n\alpha} f)(0+). \quad \square \end{aligned}$$

Theorem 3. (Generalized Taylor's formula) Suppose that $D_*^{k\alpha} f(x) \in \mathcal{C}(0, a]$ for $k = 0, 1, \dots, n+1$, where $0 < \alpha \leq 1$. Then we have

$$f(x) = \sum_{i=0}^n \frac{x^{i\alpha}}{\Gamma(i\alpha + 1)} (D_*^{i\alpha} f)(0+) + \frac{(D_*^{(n+1)\alpha} f)(\xi)}{\Gamma((n+1)\alpha + 1)} x^{(n+1)\alpha}, \quad (12)$$

with $0 \leq \xi \leq x$, $\forall x \in (0, a]$.

Proof. From (11), we have

$$\sum_{i=0}^n (J^{i\alpha} D_*^{i\alpha} f)(x) - (J^{(n+1)\alpha} D_*^{(n+1)\alpha} f)(x) = \sum_{i=0}^n \frac{x^{i\alpha}}{\Gamma(i\alpha + 1)} (D_*^{i\alpha} f)(0+), \quad (13)$$

that is,

$$f(x) - (J^{(n+1)\alpha} D_*^{(n+1)\alpha} f)(x) = \sum_{i=0}^n \frac{x^{i\alpha}}{\Gamma(i\alpha + 1)} (D_*^{i\alpha} f)(0+). \quad (14)$$

Applying the integral mean value theorem to (14) yields

$$\begin{aligned} (J^{(n+1)\alpha} D_*^{(n+1)\alpha} f)(x) &= \frac{(D_*^{(n+1)\alpha} f)(\xi)}{\Gamma((n+1)\alpha + 1)} \int_0^x (x-t)^{(n+1)\alpha} dt \\ &= \frac{(D_*^{(n+1)\alpha} f)(\xi)}{\Gamma((n+1)\alpha + 1)} x^{(n+1)\alpha}. \end{aligned} \quad (15)$$

Now, if we Substitute (15) into (14), then the generalized Taylor's formula (12) is obtained. \square

In case of $\alpha = 1$, the generalized Taylor's formula (12) reduces to the classical Taylor's formula.

5. Fractional Euler's method

In this section we shall derive the fractional Euler's method that we have developed for the numerical solution of initial value problems with Caputo derivatives. The method is a generalization of the classical Euler's method. Consider the initial value problem

$$D_*^\alpha y(t) = f(t, y(t)), \quad y(0) = y_0, \quad 0 < \alpha \leq 1, \quad t > 0. \quad (16)$$

Let $[0, a]$ be the interval over which we want to find the solution of the problem (16). In actuality, we will not find a function $y(t)$ that satisfies the initial value problem (16). Instead, a set of points $\{(t_j, y(t_j))\}$ is generated, and the points are used for our approximation.

For convenience we subdivide the interval $[0, a]$ into k subintervals $[t_j, t_{j+1}]$ of equal width $h = a/k$ by using the nodes $t_j = jh$, for $j = 0, 1, \dots, k$. Assume that $y(t)$, $D_*^\alpha y(t)$ and $D_*^{2\alpha} y(t)$ are continuous on $[0, a]$ and use the generalized Taylor's formula (12) to expand $y(t)$ about $t = t_0 = 0$. For each value t there is a value c_1 so that

$$y(t) = y(t_0) + (D_*^\alpha y(t))(t_0) \frac{t^\alpha}{\Gamma(\alpha + 1)} + (D_*^{2\alpha} y(t))(c_1) \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)}. \quad (17)$$

When $(D_*^\alpha y(t))(t_0) = f(t_0, y(t_0))$ and $h = t_1$ are substituted into equation (17), the result is an expression for $y(t_1)$:

$$y(t_1) = y(t_0) + f(t_0, y(t_0)) \frac{h^\alpha}{\Gamma(\alpha + 1)} + (D_*^{2\alpha} y(t))(c_1) \frac{h^{2\alpha}}{\Gamma(2\alpha + 1)}. \quad (18)$$

If the step size h is chosen small enough, then we may neglect the second-order term (involving $h^{2\alpha}$) and get

$$y(t_1) = y(t_0) + \frac{h^\alpha}{\Gamma(\alpha + 1)} f(t_0, y(t_0)). \quad (19)$$

The process is repeated and generates a sequence of points that approximates the solution $y(t)$. The general formula for fractional Euler's method is

$$\begin{aligned} t_{j+1} &= t_j + h, \\ y(t_{j+1}) &= y(t_j) + \frac{h^\alpha}{\Gamma(\alpha + 1)} f(t_j, y(t_j)), \end{aligned} \quad (20)$$

for $j = 0, 1, \dots, k-1$. It is clear that if $\alpha = 1$, then the fractional Euler's method (20) reduces to the classical Euler's method.

6. The algorithm

In this section we shall derive the fundamental algorithm for the numerical solution of the initial value problem (16). The new algorithm is based on the modified trapezoidal rule and the fractional Euler's method. Our approach depends on the analytical property that the initial value problem (16) is equivalent to the integral equation

$$y(t) = J^\alpha f(t, y(t)) + y(0). \quad (21)$$

Let $[0, a]$ be the interval over which we want to find the approximate the solution. Suppose that the interval $[0, a]$ is subdivided into k subintervals $[t_j, t_{j+1}]$ of equal

width $h = a/k$ by using the nodes $t_j = jh$, for $j = 0, 1, \dots, k$. To obtain the solution point $(t_1, y(t_1))$, we substitute $t = t_1$ into (21) and we get

$$y(t_1) = (J^\alpha f(t, y(t)))(t_1) + y(0). \quad (22)$$

Now if the modified trapezoidal rule (4) is used to approximate $(J^\alpha f(t, y(t)))(t_1)$ with step size $h = t_1 - t_0$, then the result is

$$y(t_1) = \alpha \frac{h^\alpha f(t_0, y(t_0))}{\Gamma(\alpha + 2)} + \frac{h^\alpha f(t_1, y(t_1))}{\Gamma(\alpha + 2)} + y(0). \quad (23)$$

Notice that the formula on the right-hand side of (23) involves the term $y(t_1)$. So, we use an estimate for $y(t_1)$. Fractional Euler's method will suffice for this purpose. Substituting (19) into (23), yields

$$y(t_1) = \alpha \frac{h^\alpha f(t_0, y(t_0))}{\Gamma(\alpha + 2)} + \frac{h^\alpha f(t_1, y(t_0) + \frac{h^\alpha}{\Gamma(\alpha+1)} f(t_0, y(t_0)))}{\Gamma(\alpha + 2)} + y(0). \quad (24)$$

The process is repeated to generate a sequence of points that approximate the solution $y(t)$. At each step, the fractional Euler's method is used as a prediction, and then the modified trapezoidal rule is used to make a correction to obtain the finite value. The general formula for our algorithm is :

$$\begin{aligned} y(t_j) = & \frac{h^\alpha}{\Gamma(\alpha + 2)} \left((j-1)^{\alpha+1} - (j-\alpha-1)j^\alpha \right) f(t_0, y(t_0)) + y(0) \\ & + \frac{h^\alpha}{\Gamma(\alpha + 2)} \sum_{i=1}^{j-1} \left((j-i+1)^{\alpha+1} - 2(j-i)^{\alpha+1} + (j-i-1)^{\alpha+1} \right) f(t_i, y(t_i)) \\ & + \frac{h^\alpha}{\Gamma(\alpha + 2)} f \left(t_j, y(t_{j-1}) + \frac{h^\alpha}{\Gamma(\alpha + 1)} f(t_{j-1}, y(t_{j-1})) \right). \end{aligned} \quad (25)$$

The new algorithm is simple for computational performance for all values of α and h . It is clear that the behavior of the method is independent of the parameter α and, as we will see in the next section, the accuracy of the approximation depends on the step size h .

7. Numerical examples

To give a clear overview of the methodology as a numerical tool, we apply the proposed algorithm on three different examples of differential equations of fractional order.

Example 1. Our first example deals with the homogeneous linear equation

$$D_*^\alpha y(t) = -y(t), \quad y(0) = 1, \quad t > 0, \quad (26)$$

where $0 < \alpha \leq 1$.

The exact solution of equation (26) is given by

$$y(t) = E_\alpha(-t^\alpha), \quad (27)$$

where

$$E_\alpha(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + 1)}, \quad (28)$$

is the Mittag-Leffler function of order α . In view of the algorithm (25), we obtain the following iteration formula:

$$\begin{aligned} y(t_j) = & 1 - \frac{h^\alpha}{\Gamma(\alpha + 2)} \left((j-1)^{\alpha+1} - (j-\alpha-1)j^\alpha \right) \\ & - \frac{h^\alpha}{\Gamma(\alpha + 2)} \sum_{i=1}^{j-1} \left((j-i+1)^{\alpha+1} - 2(j-i)^{\alpha+1} + (j-i-1)^{\alpha+1} \right) y(t_i) \\ & + \frac{h^\alpha}{\Gamma(\alpha + 2)} \left(\frac{h^\alpha}{\Gamma(\alpha + 1)} - 1 \right) y(t_{j-1}). \end{aligned} \quad (29)$$

Table 1. Numerical values for Example 1 when $\alpha = 0.5, 0.75$ and 1 with $h = 0.001$.

	$\alpha = 0.5$		$\alpha = 0.75$		$\alpha = 1.0$	
t	$u_{Approx.}$	u_{Exact}	$u_{Approx.}$	u_{Exact}	$u_{Approx.}$	u_{Exact}
0.0	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000
0.1	0.723955	0.723578	0.828254	0.828251	0.904837	0.904837
0.2	0.644094	0.643788	0.732588	0.732585	0.818730	0.818731
0.3	0.592277	0.592018	0.660341	0.660337	0.740817	0.740818
0.4	0.553831	0.553606	0.602124	0.602121	0.670319	0.670320
0.5	0.523355	0.523157	0.553605	0.553603	0.606530	0.606531
0.6	0.498203	0.498025	0.512287	0.512285	0.548811	0.548812
0.7	0.476864	0.476703	0.476556	0.476555	0.496585	0.496585
0.8	0.458394	0.458246	0.445293	0.445292	0.449328	0.449329
0.9	0.442157	0.442021	0.417683	0.417682	0.406569	0.406570
1.0	0.427709	0.427584	0.393109	0.393108	0.367079	0.367079

First we investigate the effect of allowing the value of α to vary in the interval $(0, 1]$ with fixed step size $h = 0.001$. The results are given in Table 1. It is clear that the approximate solutions are in high agreement with the exact solutions and the solution continuously depends on the time-fractional derivative. The absolute errors when $\alpha = 0.5$ and different values of the step size h are given in Table 2. The results show that the accuracy can be improved if we use much smaller values of h .

Example 2. Our second example covers the inhomogeneous linear equation

$$D_*^\alpha y(t) = \frac{2}{\Gamma(3-\alpha)} t^{2-\alpha} - \frac{1}{\Gamma(2-\alpha)} t^{1-\alpha} - y(t) + t^2 - t, \quad y(0) = 0, \quad t > 0, \quad (30)$$

where $0 < \alpha \leq 1$.

Table 2. Absolute errors for Example 1 when $\alpha = 0.5$ and with varying step sizes h .

t	$h = 0.1$	$h = 0.01$	$h = 0.004$	$h = 0.0001$
0.0	0.0000e-00	0.0000e-00	0.0000e-00	0.0000e-00
0.1	2.7642e-02	2.5504e-03	1.2841e-03	4.1574e-05
0.2	6.3532e-02	2.4222e-03	1.1042e-03	3.2673e-05
0.3	2.1765e-02	2.1647e-03	9.5509e-04	2.7271e-05
0.4	6.6783e-03	1.9368e-03	8.4009e-04	2.3510e-05
0.5	1.0867e-02	1.7463e-03	7.9939e-04	2.0697e-05
0.6	1.0016e-02	1.5871e-03	6.7599e-04	1.8494e-05
0.7	9.8945e-03	1.4526e-03	6.1530e-04	1.6713e-05
0.8	9.4682e-03	1.3377e-03	5.6423e-04	1.5231e-05
0.9	9.0545e-03	1.2381e-03	5.2061e-04	1.3995e-05
1.0	8.6295e-03	1.1521e-03	4.8291e-04	1.2932e-05

The exact solution of equation (30) is given by

$$y(t) = t^2 - t, \quad (31)$$

In view of the algorithm (25), we obtain the following iteration formula:

$$\begin{aligned}
y(t_j) = & \frac{h^\alpha}{\Gamma(\alpha+2)} \sum_{i=1}^{j-1} \left((j-i+1)^{\alpha+1} - 2(j-i)^{\alpha+1} + (j-i-1)^{\alpha+1} \right) \\
& \times \left(\frac{2}{\Gamma(3-\alpha)} t_i^{2-\alpha} - \frac{1}{\Gamma(2-\alpha)} t_i^{1-\alpha} - y(t_i) + t_i^2 - t_i \right) \\
& + \frac{h^\alpha}{\Gamma(\alpha+2)} \left(\frac{2}{\Gamma(3-\alpha)} t_j^{2-\alpha} - \frac{1}{\Gamma(2-\alpha)} t_j^{1-\alpha} - y(t_{j-1}) \right. \\
& + t_j^2 - t_j - \frac{h^\alpha}{\Gamma(\alpha+1)} \left(\frac{2}{\Gamma(3-\alpha)} t_{j-1}^{2-\alpha} - \frac{1}{\Gamma(2-\alpha)} t_{j-1}^{1-\alpha} - y(t_{j-1}) \right. \\
& \left. \left. + t_{j-1}^2 - t_{j-1} \right) \right). \quad (32)
\end{aligned}$$

The linear equation (30) is solved by Diethelm et al. [19] using the fractional Adams-Bashforth-Moulton method. Tables 3 shows the exact solution and the approximate solution for equation (30) using the algorithm (25) for different values of α . The results compare well with those obtained by the Adams-Bashforth-Moulton method [19].

Example 3. *The third example deals with the nonlinear equation*

$$D_*^\alpha y(t) = y(t)^2 - \frac{2}{(t+1)^2}, \quad y(0) = -2, \quad t > 0, \quad (33)$$

where $0 < \alpha \leq 1$.

Table 3. Numerical values for Example 2 when $\alpha = 0.5, 0.75$ and 1 with $h = 0.001$.

	$\alpha = 0.5$		$\alpha = 0.75$		$\alpha = 1.0$	
t	$u_{Appro.}$	u_{Exact}	$u_{Appro.}$	u_{Exact}	$u_{Appro.}$	u_{Exact}
0.0	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000
0.1	-0.089804	-0.090000	-0.089921	-0.090000	-0.089555	-0.090000
0.2	-0.159788	-0.160000	-0.159942	-0.160000	-0.159590	-0.160000
0.3	-0.209807	-0.210000	-0.209954	-0.210000	-0.209629	-0.210000
0.4	-0.239842	-0.240000	-0.239963	-0.240000	-0.239664	-0.240000
0.5	-0.249893	-0.250000	-0.249971	-0.250000	-0.249696	-0.250000
0.6	-0.239953	-0.240000	-0.239977	-0.240000	-0.239725	-0.240000
0.7	-0.210019	-0.210000	-0.209982	-0.210000	-0.209751	-0.210000
0.8	-0.160091	-0.160000	-0.159987	-0.160000	-0.159775	-0.160000
0.9	-0.090168	-0.090000	-0.089991	-0.090000	-0.089796	-0.090000
1.0	-0.000249	0.000000	-0.000005	0.000000	-0.000184	0.000000

The exact solution of equation (33) in case of $\alpha = 1$ is given by

$$y(t) = -\frac{2}{(t+1)}. \quad (34)$$

In view of the algorithm (25), we obtain the following iteration formula:

$$\begin{aligned}
y(t_j) = & 2 \frac{h^\alpha}{\Gamma(\alpha+2)} \left((j-1)^{\alpha+1} - (j-\alpha-1)j^\alpha \right) - 2 \\
& + \frac{h^\alpha}{\Gamma(\alpha+2)} \sum_{i=1}^{j-1} \left((j-i+1)^{\alpha+1} - 2(j-i)^{\alpha+1} + (j-i-1)^{\alpha+1} \right) \left(y^2(t_i) - \frac{2}{(t_i+1)^2} \right) \\
& + \frac{h^\alpha}{\Gamma(\alpha+2)} \left(\left(y(t_{j-1}) + \frac{h^\alpha}{\Gamma(\alpha+1)} \left(y^2(t_{j-1}) - \frac{2}{(t_{j-1}+1)^2} \right) \right)^2 - \frac{2}{(t_j+1)^2} \right)
\end{aligned} \quad (35)$$

Table 4 shows the approximate solutions for equation (33) for different values of α using the algorithm (25). The value of $\alpha = 1$ is the only case for which we know the exact solution $y(t) = -\frac{2}{(t+1)}$ and our approximate solution in this case is in high agreement with the exact solution. Of course the accuracy can be improved if we use much smaller values of h .

8. Conclusions

The fundamental goal of this work has been to construct a numerical scheme for the numerical solution of linear and nonlinear differential equations of fractional order. The goal has been achieved by using the proposed algorithm (25). This algorithm is based on the modified trapezoidal rule and the fractional Euler's method.

There are few important points to make here. First, the algorithm were used in a direct way without using linearization, perturbation or restrictive assumptions. Second, the algorithm is a reliable and very useful one for the numerical

evaluation of the Mittag-Leffler function and the other functions arising in fractional calculus. Finally, the recent appearance of fractional differential equations as models in some fields of applied mathematics makes it necessary to investigate methods of solution for such equations (analytical and numerical) and we hope that this work is a step in this direction.

Table 4. Numerical values for Example 3 with varying order α and $h = 0.001$.

t	$\alpha = 0.2$	$\alpha = 0.4$	$\alpha = 0.6$	$\alpha = 0.8$	$\alpha = 1.0$
0.0	-2.00000	-2.00000	-2.00000	-2.00000	-2.00000
0.1	-1.57191	-1.58885	-1.56451	-1.73541	-1.81818
0.2	-1.48646	-1.48991	-1.53362	-1.59189	-1.66666
0.3	-1.42059	-1.41460	-1.44369	-1.48262	-1.53846
0.4	-1.36655	-1.35254	-1.36994	-1.39284	-1.42857
0.5	-1.32108	-1.29978	-1.30711	-1.31637	-1.33333
0.6	-1.28222	-1.25413	-1.25246	-1.24987	-1.25000
0.7	-1.24861	-1.21415	-1.20427	-1.19122	-1.17647
0.8	-1.21926	-1.17879	-1.16136	-1.13896	-1.11111
0.9	-1.19341	-1.14725	-1.12283	-1.09202	-1.05263
1.0	-1.17049	-1.11893	-1.08802	-1.04957	-0.99999

REFERENCES

1. M. Caputo, *Linear models of dissipation whose Q is almost frequency independent. Part II*, J. Roy. Astr. Soc. **13** (1967), 529-539.
2. P. T. Torvik and R. L. Bagley, *On the appearance of the fractional derivative in the behavior of real materials*, J. Appl. Mech. **51** (1996), 294-298.
3. K. B. Oldham, *J. Spanier, The Fractional Calculus*, Academic Press, New York, 1974.
4. K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley and Sons, Inc., New York, 1993.
5. H. Beyer and S. Kempfle, *Definition of physically consistent damping laws with fractional derivatives*, Z. Angew Math. Mech. **75** (1995), 623-635.
6. F. Mainardi, *Fractional relaxation-oscillation and fractional diffusion-wave phenomena*, Chaos, Solitons & Fractals **7** (1996), 1461-1477.
7. R. Gorenflo and F. Mainardi, *Fractional calculus: Integral and differential equations of fractional order*. In the book *Fractals and fractional calculus*, (eds.: Carpinteri and Mainardi), New York, 1997.
8. Y. Luchko and R. Gorenflo, *The initial value problem for some fractional differential equations with the Caputo derivative*, Fachbereich Mathematik und Informatik, Freie Universität Berlin, (1998).
9. I. Podlubny, *Fractional differential equations*, Academic Press, San Diego, CA, 1999.
10. K. Diethelm and N. Ford, *Analysis of fractional differential equations*, J. Math. Anal. Appl. **265** (2002), 229-248.
11. F. Huang and F. Liu, *The time-fractional diffusion equation and fractional advection-dispersion equation*, ANZIAM J. **46** (2005), 1-14.
12. F. Huang and F. Liu, *The fundamental solution of the space-time fractional advection-dispersion equation*, J. Appl. Math. and Computing **18(2)** (2005), 339-350.

13. L. Blank, *Numerical treatment of differential equations of fractional order*, Numerical analysis report 287, Manchester center for computational mathematics, Manchester, 1996.
14. R. Gorenflo, *Fractional calculus: Some numerical methods*. In the book *Fractals and fractional calculus*, (eds.: Carpinteri and Mainardi), New York, 1997.
15. I. Podlubny, *Numerical solution of ordinary fractional differential equations by the fractional difference method*. In the book *Advances in Difference Equations*, (eds.: S.Elaydi, I.Gyori, G.Ladas), Gordon and Breach Sci. Publ., Amsterdam, 1997
16. K. Diethelm, *An algorithm for the numerical solution for differential equations of fractional order*, Elec. Transact. Numer. Anal. **5** (1997), 1-6.
17. K. Diethelm and G. Walz, *Numerical solution for fractional differential equations by extrapolation*, Numerical algorithms **16** (1997), 231-253.
18. K. Diethelm, N. Ford and A. Freed, *A Predictor-Corrector approach for the numerical solution of fractional differential equations*, Nonlinear Dynamics **29** (2002), 3-22.
19. K. Diethelm, N. Ford and A. Freed, *Detailed error analysis for a fractional Adams Method*, Numer. Algorithms **36** (2004), 31-52.
20. G. Adomian, *A review of the decomposition method in applied mathematics*, J. Math. Anal. Appl. **135** (1988), 501-544.
21. G. Adomian, *Solving Frontier Problems of Physics: The Decomposition method*, Kluwer Academic Publishers, Boston, 1994.
22. A. Wazwaz, *A new algorithm for calculating Adomian polynomials fo nonlinear operators*, Appl. Math. Comput. **111** (2000), 53-69.
23. A. Wazwaz and S. El-Sayed, *A new modificatrion of the Adomian decomposition method for linear and nonlinear operators*, Appl. Math. Comput. **122** (2001), 393-405.
24. N. Shawagfeh, *Analytical approximate solutions for nonlinear fractional differential equations*, Appl. Math. Comput. **131** (2002), 517-529.
25. S. Momani, *An explicit and numerical solutions of the fractional KdV equation*, Math. Comput. Simul. **70(2)** (2005), 110-118.
26. S. Momani, *Non-perturbative analytical solutions of the space- and time-fractional Burgers equations*, Chaos, Solitons & Fractals **28(4)** (2006), 930-937.
27. S. Momani and Z. Odibat, *Analytical solution of a time-fractional Navier-Stokes equation by Adomian decomposition method*, Appl. Math. Comput. **177(2)** (2006), 488-494.
28. Z. Odibat and S. Momani, *Approximate solutions for boundary value problems of time-fractional wave equation*, Appl. Math. Comput. **181(1)** (2006), 767-774.
29. J. H. He, *Variational iteration method for delay differential equations*, Commun. Nonlin. Sci. Numer. Simulat. **2(4)** (1997), 235-36.
30. J. H. He, *Semi-inverse method of establishing generalized principles for fluid mechanics with emphasis on turbomachinery aerodynamics*, Int. J. Turbo Jet-Engines **14(1)** (1997), 23-28.
31. J. H. He, *Approximate solution of non linear differential equations with convolution product nonlinearities*, Computer Meth. Appl. Mech. Eng. **167** (1998), 69-73.
32. J. H. He, *Approximate analytical solution for seepage flow with fractional derivatives in porous media*, Computer Meth. Appl. Mech. Eng. **167** (1998), 57-68.
33. J. H. He, *Variational iteration method- a kind of non-linear analytical technique: some examples*, Int. J. Nonlin. Mech. **34** (1999), 699-708.
34. J. H. He, *Variational iteration method for autonomous ordinary differential systems*, Appl. Math. Comput. **114** (2000), 115-123.
35. J. H. He, *Variational theory for linear magneto-electro-elasticity*, Int. J. Nonlin. Sci. Numer. Simulat. **2(4)** (2001), 309-316.
36. J. H. He, *Variational principle for some nonlinear partial differential equations with variable coefficients*, Chaos, Solitons & Fractals **19(4)** (2004), 847-851.
37. J. H. He, *Asymptotic methods for strongly nonlinear equations*, Int. J. Modern Phys. B **20(10)** (2006), 1141-1199.

38. J. H. He, *Non-perturbative methods for strongly nonlinear problems*, Berlin: dissertation, de-Verlag im Internet GmbH, 2006.
39. J. H. He and X. H. Wu, *Construction of solitary solution and compacton-like solution by variational iteration method*, Chaos, Solitons & Fractals **29(1)** (2006), 108-113.
40. A. A. Soliman, *A numerical simulation and explicit solutions of KdV-Burgers' and Lax's seventh-order KdV equations*, Chaos, Solitons & Fractals **29(2)** (2006), 294-302.
41. E. M. Abulwafa, M. A. Abodu and A. A. Mohmoud, *The solution of nonlinear coagulation problem with mass loss*, Chaos, Solitons & Fractals **29(2)** (2006), 313-330.
42. N. Bildik and A. Konuralp, *The use of variational iteration method, differential transform method and Adomian decomposition method for solving different types of nonlinear partial differential equations*, Int. J. Nonlin. Sci. Numer. Simulat. **7(1)** (2006), 65-70.
43. T. H. Hao, *Search for variational principles in electrodynamics by Lagrange method*, Int. J. Nonlin. Sci. Numer. Simulat. **6(2)** (2005), 209-210.
44. S. Momani and S. Abuasad, *Application of He's variational iteration method to Helmholtz equation*, Chaos, Solitons & Fractals **27(5)** (2006), 1119-1123.
45. Z. Odibat and S. Momani, *Application of Variational Iteration Method to Nonlinear Differential Equations of Fractional Order*, Int. J. Nonlin. Sci. Numer. Simulat. **7(1)** (2006), 15-27.
46. S. Momani and Z. Odibat, *Numerical comparison of methods for solving linear differential equations of fractional order*, Chaos, Solitons & Fractals **31(5)** (2007), 1248-1255.
47. S. Momani and Z. Odibat, *Analytical approach to linear fractional partial differential equations arising in fluid mechanics*, Physics Letters A **355(4-5)** (2006), 271-279.
48. Z. Odibat, *Approximations of fractional integral and Caputo fractional derivatives*, Appl. Math. Comput. **178(2)** (2006), 527-533.
49. Z. Odibat and N. Shawagfeh, *Generalized Taylor's formula*, Appl. Math. Comput. **186(1)** (2007), 286-293.

Zaid M. Odibat received his Ph.D from the University of Jordan (Jordan) in 2002. Since 2002 he has been at Al-Balqa applied University. His research interests focus on ordinary and partial differential equations, fractional calculus and numerical analysis.

Prince Abdullah Bin Ghazi Faculty of Science and IT, Al-Balqa Applied University, Salt, Jordan

e-mail: odibat@bau.edu.jo

Shaher Momani received his Ph.D from the University of Wales (UK) in 1991 under the direction of Ken Walters (FRS). Since 1991 he has been at Mutah University in Jordan. He left Mutah University to spend one year at Yarmouk University (2000-2001) and three years at United Arab Emirates University (2001-2004). His research interests focus on the numerical solution of fractional differential equations in fluid mechanics, non-Newtonian fluid mechanics and numerical analysis.

Department of Mathematics, Mutah University, P. O. Box 7, Al-Karak, Jordan

e-mail: shaherm@yahoo.com