

QSOonic: fast quasar continuum fitting

Naim Göksel Karaçaylı ^{1,2,3}

¹ Center for Cosmology and AstroParticle Physics, The Ohio State University, 191 West Woodruff Avenue, Columbus, OH 43210, USA ² Department of Astronomy, The Ohio State University, 4055 McPherson Laboratory, 140 W 18th Avenue, Columbus, OH 43210, USA ³ Department of Physics, The Ohio State University, 191 West Woodruff Avenue, Columbus, OH 43210, USA

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Ivelina Momcheva](#) 

Reviewers:

- [@gcalderone](#)
- [@corentinravoux](#)

Submitted: 18 January 2024

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

Summary

Distant quasars show absorption features at wavelengths shorter than the Ly α emission line in their spectrum due to intervening neutral hydrogen gas. The Ly α forest technique uses these absorption lines to map out the matter distribution in vast volumes when the universe was 1.5–3 billion years old ($2 \lesssim z \lesssim 5$). This map of the cosmos successfully constrained cosmological models using the Baryon Acoustic Oscillations feature in the 3D correlation function ([du Mas des Bourboux et al., 2020](#)), various dark matter models ([Iršič et al., 2017](#)) and the sum of the neutrino masses ([Palanque-Delabrouille et al., 2015](#)) using the 1D Ly α forest power spectrum ([Karaçaylı et al., 2022, 2024](#)).

The density of the intervening gas is inferred from the transmitted flux fraction which is the absorption depth relative to the unabsorbed quasar continuum. The field has devised a simple continuum model to homogenize the analysis across quasars so that the errors in the estimated continuum are contained to well-known modes. This model assumes each quasar continuum C_q is a scaling of a mean quasar continuum shape \bar{C} by a polynomial of $\ln \lambda$. For example, an order 1 polynomial introduces amplitude a_q and slope b_q terms for each quasar such that $C_q = (a_q + b_q \ln \lambda) \bar{C}$. The quasar continuum fitting algorithm is then an iterative solution that minimizes χ^2 for each quasar and updates the mean quasar continuum using the stack of all quasar continua after every iteration ([du Mas des Bourboux et al., 2020](#); [Karaçaylı et al., 2024](#)). This method requires efficient parallelization due to computational time and memory needed.

Statement of need

QSOonic is an MPI-parallelized, highly optimized Python package specifically built for the Dark Energy Spectroscopic Instrument (DESI), which will collect approximately almost a million Ly α quasar spectra in the near future ([DESI Collaboration et al., 2016](#)). QSOonic is designed to overcome the numerical challenge posed by this large quantity of spectra by MPI parallelization that can distribute the memory and workload across multiple compute nodes, and provide an efficient API to read and manipulate DESI quasar spectra to allow implementation of other algorithms within the Ly α forest framework. It provides detailed intermediate data products including best-fit parameters for each quasar and a delete-one Jackknife estimate for the covariance matrix used in pipeline noise calibration correction calculation. These intermediate products are crucial to ascertain the precision of the fitted continua and the pipeline noise corrections.

QSOonic is built on the same algorithm as [picca](#), but through MPI parallelization and efficient numerical algorithms, it is faster by an order of magnitude. Unlike [picca](#), QSOonic has robust convergence criteria, provides a generic API for DESI quasar spectra, and saves more intermediate data products.

43 QSONic will be used in future scientific publications of the 1D Ly α forest power spectrum and
44 the 3D correlation function measurements from DESI data.

45 References

- 46 DESI Collaboration, Aghamousa, A., Aguilar, J., Ahlen, S., Alam, S., Allen, L. E., Allende
47 Prieto, C., Annis, J., Bailey, S., Balland, C., Ballester, O., Baltay, C., Beaufore, L.,
48 Bebek, C., Beers, T. C., Bell, E. F., Bernal, J. L., Besuner, R., Beutler, F., ... Zu, Y.
49 (2016). The DESI Experiment Part I: Science, Targeting, and Survey Design. *arXiv e-Prints*,
50 arXiv:1611.00036. <https://arxiv.org/abs/1611.00036>
- 51 du Mas des Bourboux, H., Rich, J., Font-Ribera, A., de Sainte Agathe, V., Farr, J., Etourneau,
52 T., Le Goff, J.-M., Cuceu, A., Balland, C., Bautista, J. E., Blomqvist, M., Brinkmann,
53 J., Brownstein, J. R., Chabanier, S., Chaussidon, E., Dawson, K., González-Morales,
54 A. X., Guy, J., Lyke, B. W., ... Youles, S. (2020). The Completed SDSS-IV Extended
55 Baryon Oscillation Spectroscopic Survey: Baryon Acoustic Oscillations with Ly α Forests.
56 *Astrophysical Journal*, 901(2), 153. <https://doi.org/10.3847/1538-4357/abb085>
- 57 Iršič, V., Viel, M., Haehnelt, M. G., Bolton, J. S., Cristiani, S., Becker, G. D., D'Odorico, V.,
58 Cupani, G., Kim, T.-S., Berg, T. A. M., López, S., Ellison, S., Christensen, L., Denney, K.
59 D., & Worseck, G. (2017). New constraints on the free-streaming of warm dark matter
60 from intermediate and small scale Lyman- α forest data. *Phys. Rev. D.*, 96(2), 023522.
61 <https://doi.org/10.1103/PhysRevD.96.023522>
- 62 Karaçaylı, N. G., Martini, P., Guy, J., Ravoux, C., Karim, M. L. A., Armengaud, E., Walther,
63 M., Aguilar, J., Ahlen, S., Bailey, S., Bautista, J., Beltran, S. F., Brooks, D., Cabayol-
64 Garcia, L., Chabanier, S., Chaussidon, E., Chaves-Montero, J., Dawson, K., Cruz, R.
65 de la, ... Zhou, Z. (2024). Optimal 1D Ly α Forest Power Spectrum Estimation – III.
66 DESI early data. *Monthly Notices of the Royal Astronomical Society*, stae171. <https://doi.org/10.1093/mnras/stae171>
- 67 Karaçaylı, N. G., Padmanabhan, N., Font-Ribera, A., Iršič, V., Walther, M., Brooks, D.,
68 Gaztañaga, E., Kehoe, R., Levi, M., Ntelis, P., Palanque-Delabrouille, N., & Tarlé, G.
69 (2022). Optimal 1D Ly α forest power spectrum estimation - II. KODIAQ, SQUAD,
70 and XQ-100. *Monthly Notices of the Royal Astronomical Society*, 509(2), 2842–2855.
71 <https://doi.org/10.1093/mnras/stab3201>
- 72 Palanque-Delabrouille, N., Yèche, C., Lesgourgues, J., Rossi, G., Borde, A., Viel, M., Aubourg,
73 E., Kirkby, D., LeGoff, J.-M., Rich, J., Roe, N., Ross, N. P., Schneider, D. P., &
74 Weinberg, D. (2015). Constraint on neutrino masses from SDSS-III/BOSS Ly α forest
75 and other cosmological probes. *J. Cosmology Astropart. Phys.*, 2015(2), 045–045.
76 <https://doi.org/10.1088/1475-7516/2015/02/045>
- 77