

# CRNPy: An Open-Source Python Library for Cosmic Ray Neutron Probe Data Processing

Joaquin Peraza Rud<sup>1</sup>, Tyson Ochsner<sup>2</sup>, and Andres Patrignani<sup>1</sup>

<sup>1</sup> Department of Agronomy, Kansas State University, Manhattan, KS, USA. <sup>2</sup> Department of Plant and Soil Sciences, Oklahoma State University, Stillwater, OK, USA.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Jayaram Hariharan](#)

## Reviewers:

- [@jlarsen-usgs](#)
- [@danpower101](#)

Submitted: 09 August 2023

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

CRNPy is a Python library that facilitates the processing, analysis, and conversion of raw cosmic-ray neutron counts obtained with stationary and roving cosmic-ray neutron probes (CRNP) into volumetric soil water content. The CRNPy library includes routines for atmospheric, biomass, and road corrections, along with one-dimensional and two-dimensional filtering. The library extends its utility by offering horizontal and vertical footprint determination, uncertainty estimation, depth extrapolation operators, and routines to assist users with field calibration. The design of the CRNPy library prioritizes reproducibility, ease of use, and compatibility across instruments, facilitating its adoption by manufacturers, end users, and researchers working to integrate non-invasive soil moisture sensing in agricultural and hydrological applications.


## Statement of Need

Cosmic ray neutron probes (CRNP) are non-invasive soil moisture sensors that fill the niche between point-level and satellite sensors. However, the conversion of raw CRNP data into soil moisture requires multiple corrections and filtering steps that are described across various peer-reviewed articles. To circumvent this limitation and enhance reproducibility, the CRNPy library offers a simple, instrument-agnostic, and integrated solution with minimal dependencies. Compared to the existing `crspy` (Power et al., 2021) library, CRNPy avoids stringent data naming conventions and external data requirements. A flexible data naming convention enables a seamless integration with output files from different instrument manufacturers and the lack of external data requirements makes the library more compact (only ~65 KB) and straight forward to install. Compared to the `corny` (Schrön, Accessed: 2024) toolbox's GUI-based workflow, CRNPy's modular design based on Python functions promotes integration and reproducibility within data analysis pipelines and interactive development environments like Jupyter Lab notebooks. In addition, its straightforward installation using the Python Package Index, the minimal dependencies—most included with the Anaconda open-source ecosystem—and the comprehensive datasets with included examples in the form of Jupyter notebooks, provide an accessible start for CRNP data processing. The CRNPy library emphasizes easy maintenance and community-driven improvements since users can expand its capabilities by adding regular Python functions to the core module. The compact size and simple structure of the CRNPy library can also enable future integration into cloud-based services, IoT sensors, and system-on-chip technologies, broadening its use and customization potential.


## Library features

The CRNPy library integrates standard routines for processing CRNP data, with features including:


- 40 ■ Utilization of core scientific Python libraries like Numpy (Harris et al., 2020), Pandas (McKinney & others, 2011), SciPy (Virtanen et al., 2020), and Matplotlib (Hunter & Dale, 2007), that are readily available within the Anaconda environment. All CRNPy functions are compatible with Numpy arrays or Pandas series for robust data science functionality.
- 45 ■ Utility functions for obtaining site-specific lattice water, geomagnetic cutoff rigidity (Smart & Shea, 2001), and neutron monitor references (Klein et al., 2009), which are required for pre-processing raw neutron counts.
- 48 ■ Flexible input data handling from delimited text files without stringent naming conventions for columns, which keeps scripts simpler, increases reproducibility, and minimizes human error. This aspect also enables a more versatile, modular, and customizable workflow (Figure 1 and Figure 2) that adapts to instrument outputs from different manufacturers.
- 51 ■ Detection of possible outliers using range detection based on user-provided lower and upper boundaries, interquartile range, z-scores, and a scaled mean absolute difference (Iglewicz & Hoaglin, 1993).
- 55 ■ Corrections for atmospheric pressure as described by Zreda et al. (2012), air humidity (Rosolem et al., 2013), and incoming neutron flux following the guidelines from Zreda et al. (2012); Hawdon et al. (2014); McJannet & Desilets (2023). The article by Andreasen et al. (2017) provides an overall description of these correction methods included in CRNPy (Figure 3a and Figure 3b).
- 60 ■ Corrections to account for additional hydrogen pools in above- and below-ground plant biomass (Baatz et al., 2015; Wahbi et al., 2018).
- 62 ■ Corrections to account for the impact of road soil moisture conditions during roving surveys (Schrön et al., 2018).
- 64 ■ Conversion of corrected counts into volumetric soil water content following the approach suggested by Desilets et al. (2010).
- 66 ■ Determination of neutron count uncertainty following the method detailed in Jakobi et al. (2020) (see Figure 3c).
- 68 ■ Estimation of sensing depth by determining the volume that accounts for 86% of the origin of the counted neutrons (Franz et al., 2012; Schrön et al., 2017).
- 70 ■ An exponential filter operator (Albergel et al., 2008) to extend near-surface soil moisture conditions to the rootzone (Franz et al., 2020; Rossini & Patrignani, 2021), see Figure 3d.
- 72 ■ Utility functions for spatial filtering and basic interpolation routines required to process CRNP roving surveys (see Figure 4)
- 74 ■ Additional functions for temporal interpolation and filtering required to process stationary CRNP observations (see Figure 3).




**Figure 1:** Example workflow for stationary CRNP, dashed lines represent optional steps.



**Figure 2:** Example workflow for roving CRNP, dashed lines represent optional steps.



**Figure 3:** Outputs of a stationary device in each of the steps of the workflow, A) Impact of each factor in the correction process. B) Difference between raw and corrected counts. C) Resulting volumetric water content time series with the propagated neutron count uncertainty. D) Depth that accounts for the 86% of the observed counts.



**Figure 4:** Contour plot of the roving transect spatial output. Each marker represents the estimated center of the observation.

## Acknowledgements

This work was supported by the USDA National Institute of Food and Agriculture through the Agriculture and Food Research Initiative Competitive Grant no. 2019-68012-29888 and through Multistate project W4188. The authors declare no competing interests. This is contribution no. 24-179-J of the Kansas Agricultural Experiment Station.

## References

- Albergel, C., Rüdiger, C., Pellarin, T., Calvet, J.-C., Fritz, N., Froissard, F., Suquia, D., Petitpa, A., Pigué, B., & Martin, E. (2008). From near-surface to root-zone soil moisture using an exponential filter: An assessment of the method based on in-situ observations and model simulations. *Hydrology and Earth System Sciences*, 12(6), 1323–1337. <https://doi.org/10.5194/hess-12-1323-2008>
- Andreasen, M., Jensen, K. H., Desilets, D., Franz, T. E., Zreda, M., Bogen, H. R., & Looms, M. C. (2017). Status and perspectives on the cosmic-ray neutron method for soil moisture estimation and other environmental science applications. *Vadose Zone Journal*, 16(8), 1–11. <https://doi.org/10.2136/vzj2017.04.0086>
- Baatz, R., Bogen, H., Hendricks Franssen, H.-J., Huisman, J., Montzka, C., & Vereecken, H. (2015). An empirical vegetation correction for soil water content quantification using

- 93 cosmic ray probes. *Water Resources Research*, 51(4), 2030–2046. <https://doi.org/10.1002/2014WR016443>
- 94
- 95 Desilets, D., Zreda, M., & Ferré, T. P. (2010). Nature's neutron probe: Land surface  
96 hydrology at an elusive scale with cosmic rays. *Water Resources Research*, 46(11).  
97 <https://doi.org/10.1029/2009WR008726>
- 98 Franz, T. E., Wahbi, A., Zhang, J., Vreugdenhil, M., Heng, L., Dercon, G., Strauss, P., Brocca,  
99 L., & Wagner, W. (2020). Practical data products from cosmic-ray neutron sensing for  
100 hydrological applications. *Frontiers in Water*, 2, 9. <https://doi.org/10.3389/frwa.2020.00009>
- 101
- 102 Franz, T. E., Zreda, M., Ferre, T., Rosolem, R., Zweck, C., Stillman, S., Zeng, X., &  
103 Shuttleworth, W. (2012). Measurement depth of the cosmic ray soil moisture probe  
104 affected by hydrogen from various sources. *Water Resources Research*, 48(8). <https://doi.org/10.1029/2012WR011871>
- 105
- 106 Harris, C. R., Millman, K. J., Van Der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau,  
107 D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., & others. (2020). Array programming  
108 with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- 109 Hawdon, A., McJannet, D., & Wallace, J. (2014). Calibration and correction procedures  
110 for cosmic-ray neutron soil moisture probes located across australia. *Water Resources*  
111 *Research*, 50(6), 5029–5043. <https://doi.org/10.1002/2013WR015138>
- 112 Hunter, J., & Dale, D. (2007). The matplotlib user's guide. *Matplotlib 0.90. 0 User's Guide*.
- 113 Iglewicz, B., & Hoaglin, D. C. (1993). *Volume 16: How to detect and handle outliers*. Quality  
114 Press.
- 115 Jakobi, J., Huisman, J. A., Schrön, M., Fiedler, J., Brogi, C., Vereecken, H., & Bogaen, H. R.  
116 (2020). Error estimation for soil moisture measurements with cosmic ray neutron sensing  
117 and implications for rover surveys. *Frontiers in Water*, 2, 10. <https://doi.org/10.3389/frwa.2020.00010>
- 118
- 119 Klein, K.-L., Steigies, C., & NMDB Team. (2009). WWW.NMDB.EU: The real-time Neutron  
120 Monitor database. *EGU General Assembly Conference Abstracts*, 5633.
- 121 McJannet, D., & Desilets, D. (2023). Incoming neutron flux corrections for cosmic-ray soil  
122 and snow sensors using the global neutron monitor network. *Water Resources Research*,  
123 59(4), e2022WR033889. <https://doi.org/10.1029/2022WR033889>
- 124 McKinney, W., & others. (2011). Pandas: A foundational python library for data analysis and  
125 statistics. *Python for High Performance and Scientific Computing*, 14(9), 1–9.
- 126 Power, D., Rico-Ramirez, M. A., Desilets, S., Desilets, D., & Rosolem, R. (2021). Cosmic-ray  
127 neutron sensor PYthon tool (crspy 1.2. 1): An open-source tool for the processing of  
128 cosmic-ray neutron and soil moisture data. *Geoscientific Model Development*, 14(12),  
129 7287–7307. <https://doi.org/10.5194/gmd-14-7287-2021>
- 130 Rosolem, R., Shuttleworth, W. J., Zreda, M., Franz, T. E., Zeng, X., & Kurc, S. A. (2013).  
131 The effect of atmospheric water vapor on neutron count in the cosmic-ray soil moisture  
132 observing system. *Journal of Hydrometeorology*, 14(5), 1659–1671. <https://doi.org/10.1175/JHM-D-12-0120.1>
- 133
- 134 Rossini, P., & Patrignani, A. (2021). Predicting rootzone soil moisture from surface observations  
135 in cropland using an exponential filter. *Soil Science Society of America Journal*, 85(6),  
136 1894–1902. <https://doi.org/10.1002/saj2.20319>
- 137 Schrön, M. (Accessed: 2024). *CORNish PASDy - COsmic-ray neutron flavored PASDy*.  
138 [https://git.ufz.de/CRNS/cornish\\_pasdy](https://git.ufz.de/CRNS/cornish_pasdy).

- 139 Schrön, M., Köhli, M., Scheffele, L., Iwema, J., Bogen, H. R., Lv, L., Martini, E., Baroni, G.,  
140 Rosolem, R., Weimar, J., Mai, J., Cuntz, M., Rebmann, C., Oswald, S. E., Dietrich, P.,  
141 Schmidt, U., & Zacharias, S. (2017). Improving calibration and validation of cosmic-ray  
142 neutron sensors in the light of spatial sensitivity. *Hydrology and Earth System Sciences*,  
143 21(10), 5009–5030. <https://doi.org/10.5194/hess-21-5009-2017>
- 144 Schrön, M., Rosolem, R., Köhli, M., Piuissi, L., Schröter, I., Iwema, J., Kögler, S., Oswald,  
145 S. E., Wollschläger, U., Samaniego, L., Dietrich, P., & Zacharias, S. (2018). Cosmic-ray  
146 neutron rover surveys of field soil moisture and the influence of roads. *Water Resources*  
147 *Research*, 54(9), 6441–6459. <https://doi.org/10.1029/2017WR021719>
- 148 Smart, D., & Shea, M. (2001). *Geomagnetic cutoff rigidity computer program: Theory,*  
149 *software description and example.*
- 150 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,  
151 Burovski, E., Peterson, P., Weckesser, W., Bright, J., & others. (2020). SciPy 1.0:  
152 Fundamental algorithms for scientific computing in python. *Nature Methods*, 17(3),  
153 261–272. <https://doi.org/10.1038/s41592-019-0686-2>
- 154 Wahbi, A., Heng, L., Dercon, G., Wahbi, A., & Avery, W. (2018). In situ destructive sampling.  
155 *Cosmic Ray Neutron Sensing: Estimation of Agricultural Crop Biomass Water Equivalent*,  
156 5–9. [https://doi.org/10.1007/978-3-319-69539-6\\_2](https://doi.org/10.1007/978-3-319-69539-6_2)
- 157 Zreda, M., Shuttleworth, W., Zeng, X., Zweck, C., Desilets, D., Franz, T., & Rosolem, R.  
158 (2012). COSMOS: The cosmic-ray soil moisture observing system. *Hydrology and Earth*  
159 *System Sciences*, 16(11), 4079–4099. <https://doi.org/10.5194/hess-16-4079-2012>