Tables Suzanne Dufault July 10, 2019

Table 1: Characteristics of suicides, overdoses, and person-time distribution, UAW-GM cohort 1941-2015

Characteristic	Person-years (N = $1,505,574$)	Suicide ($N = 276$)	Overdoses $(N = 67)$
Age			
mean (SD)	$49.92\ (15.11)$	$38.26 \ (10.16)$	$36.31 \ (8.43)$
Sex, $\%$			
Males	87.95	95.65	95.52
Females	12.05	4.35	4.48
Race, $\%$			
Whites	82.10	88.41	74.63
Blacks	17.90	11.59	25.37
Plant, %			
Plant 1	23.01	17.39	19.40
Plant 2	42.31	49.64	65.67
Plant 3	34.55	32.25	14.93
Calendar year of hire			
min	1938	1940	1942
mean (SD)	1962 (11.47)	1962 (11.35)	1970 (10.95)
max	1987	1979	1978
Years since hire			
mean (SD)	$42.94\ (10.87)$	$23.22\ (13.63)$	23.96 (12.34)

Suicide was determined by ICD-9 codes E950-E959 and ICD-10 codes X60-X84, Y87, and U03. Overdose was determined by ICD-9 codes E850-E858, E980.0-E980.5 (undetermined intent) and ICD-10 codes X40-X44 and Y10-Y14.