


Steven R. Dunbar
Department of Mathematics
203 Avery Hall
University of Nebraska-Lincoln
Lincoln, NE 68588-0130
<http://www.math.unl.edu>
Voice: 402-472-3731
Fax: 402-472-8466

Topics in Probability Theory and Stochastic Processes Steven R. Dunbar

Lévy's Construction of Brownian Motion



Study Tip



Rating

Mathematicians Only: prolonged scenes of intense rigor.



Section Starter Question

If Z_1 and Z_2 are independent $N(0, \sigma^2)$ random variables, then what can be said about $Z_1 + Z_2$ and $Z_1 - Z_2$?



Key Concepts

1. Brownian Motion can be defined in a natural way by first constructing a set of independent normal random variables on the dyadic rationals and then combining them inductively to have the right incremental distributions for Brownian Motion.
 2. The resulting construction of a continuous function B_t on $[0,1]$ has the properties:
 - (a) $B_0 = 0$,
 - (b) for all $t_1 < t_2 < \dots < t_{k-1} < t_k$ in $[0, 1]$, $B_{t_k} - B_{t_{k-1}}, \dots, B_{t_2} - B_{t_1}$ are independent with joint distribution $(N(0, t_k - t_{k-1}), \dots, N(0, t_2 - t_1))$,
 - (c) and almost surely $t \mapsto B_t$ is continuous on $[0, 1]$.
-



Vocabulary

1. A one-dimensional **standard Brownian Motion** is a stochastic process $B : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ (alternatively a collection of random variables (indexed by $x \in [0, \infty)$) with the following properties:
 - (a) $B(0) = 0$ almost surely.
 - (b) For all $t_1 < t_2 < \dots < t_{k-1} < t_k$ in $[0, 1]$, $B_{t_k} - B_{t_{k-1}}, \dots, B_{t_2} - B_{t_1}$ are independent with joint distribution $(N(0, t_k - t_{k-1}), \dots, N(0, t_2 - t_1))$,
 - (c) Almost surely, the path $x \rightarrow B(x)$ (called a sample path) is a continuous function on $[0, \infty)$.
 2. $D_n = \{k/2^n : 0 \leq k \leq 2^n\}$ and $D = \bigcup_n D_n$ are the **dyadic rationals**, a countable dense set in $[0, 1]$.
-

Notation

1. $Z \sim N(0, 1)$ – A normally distributed random variable with mean 0 and variance 1.
2. $\{A_i\}_{i=0}^\infty$ – A sequence of events in some probability space.
3. $\limsup_{i \rightarrow \infty} A_i = \bigcap_{i=0}^\infty \bigcup_{j=i}^\infty A_j = \{A_i, \text{i.o.}\}$ – An event which occurs “infinitely often”.
4. $B : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ or B_t – The standard one-dimensional Brownian Motion as a stochastic process.
5. $D_n = \{k/2^n : 0 \leq k \leq 2^n\}$ and $D = \bigcup_n D_n$ are the **dyadic rationals**, a countable dense set in $[0, 1]$.
6. $\{Z_d\}$ – A countable collection of $N(0, 1)$ independent random variables indexed by D .
7. B_d – Random variables derived from Z_d , indexed by D with the independent increments property on D .

8. $d^+ = d + 2^{-n}$ and $d^- = d - 2^{-n}$ – The neighbors of d , which are actually members of D_{n-1} .
9. $r < s < t$ – Arbitrary points in D .
10. $F_0(t)$ for $n = 0, 1, 2, \dots$ – Piecewise linear functions defined in terms of the random variables Z_d .
11. $t_1 < t_2 < t_3$ – Points in $[0, 1]$.
12. $t_i = \lim_{n \rightarrow \infty} t_{i,n}$ for $i = 1, 2, 3$ – Sequences of dyadics from D with limit t_i .
13. $B_t^{(n)}$ – Independent standard Brownian Motion stochastic processes on the unit interval.



Mathematical Ideas

Preliminary Lemmas

Lemma 1. *If $Z \sim N(0, 1)$ then $\mathbb{P}[Z \geq x] \leq \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{x} \cdot e^{-x^2/2}$.*

Proof.

$$\begin{aligned}
 \mathbb{P}[Z \geq x] &= \int_x^\infty \frac{1}{\sqrt{2\pi}} e^{-u^2/2} du \\
 &\leq \int_x^\infty \frac{1}{\sqrt{2\pi}} \frac{u}{x} e^{-u^2/2} du \\
 &= \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{x} \int_x^\infty u e^{-u^2/2} du \\
 &= \frac{1}{\sqrt{2\pi}} \cdot \frac{1}{x} \cdot e^{-x^2/2}
 \end{aligned}$$

□

Remark. For $x \geq 1/(\sqrt{2\pi}) \approx 0.398$, the simpler but cruder estimate is useful: $\mathbb{P}[Z \geq x] \leq e^{-x^2/2}$.

Lemma 2 (Borel-Cantelli). *Let $\{A_i\}_{i=0}^\infty$ be a sequence of events in some probability space. Let $\limsup_{i \rightarrow \infty} A_i = \bigcap_{i=0}^\infty \bigcup_{j=i}^\infty A_i = \{A_i, i.o.\}$, where *i.o.* stands for infinitely often.*

1. *If $\sum_{i=0}^\infty \mathbb{P}[A_i] < \infty$, then $\mathbb{P}[\{A_i, i.o.\}] = 0$.*
2. *If the events $\{A_i\}_{i=0}^\infty$ are pairwise independent, and $\sum_{i=0}^\infty \mathbb{P}[A_i] = \infty$, then $\mathbb{P}[\{A_i, i.o.\}] = 1$.*

Proof. Omitted here. □

Lévy's Construction of Brownian Motion

Definition. A one-dimensional **standard Brownian Motion** is a stochastic process $B : \Omega \times [0, \infty) \rightarrow \mathbb{R}$ (alternatively a collection of random variables indexed by $x \in [0, \infty)$) with the following properties:

1. $B_0 = 0$,
2. for all $t_1 < t_2 < \dots < t_{k-1} < t_k$ in $[0, 1]$, $B_{t_k} - B_{t_{k-1}}, \dots, B_{t_2} - B_{t_1}$ are independent with joint distribution $(N(0, t_k - t_{k-1}), \dots, N(0, t_2 - t_1))$,
3. and almost surely $t \mapsto B_t$ is continuous on $[0, 1]$.

Theorem 3 (Wiener, 1923). *One-dimensional standard Brownian Motion B_t exists.*

Remark. Although Norbert Wiener mathematically described Brownian Motion first in 1923, the following construction is due to Paul Lévy in 1948.

Proof. Step 1: Construct a set of random values on the dyadics in $[0, 1]$ satisfying the independent normal increments property.

- (a) Let $D_n = \{k/2^n : 0 \leq k \leq 2^n\}$ and $D = \bigcup_n D_n$ be the **dyadic rationals**, a countable dense set in $[0, 1]$.
- (b) Let $\{Z_d\}$ be a countable collection of $N(0, 1)$ independent random variables indexed by D .

- (c) Inductively define a some new values B_d indexed by D with the independent increments property on D . Begin by defining $B_0 = 0$ and $B_1 = Z_1$.
- (d) Claim: It is possible to make a collection B_d , with $d \in D$, such that for $r < s < t$ in D , $B_t - B_s \sim N(0, t - s)$ is independent of $B_s - B_r \sim N(0, s - r)$ and furthermore, that the B_d with $d \in D_n$ are globally independent of Z_d for $d \in D \setminus D_n$.
- (e) Proof of Claim by Induction: The base case for $n = 0$ with D_0 holds since $r = 0$, $t = 1$ and there is no s . Then $B_1 - B_0 = Z_1 - 0 = Z_1 \sim N(0, 1)$. $B_1 = Z_1$ is independent of all other B_d by constructions and $B_0 = 0$ is independent of all other B_d .
The induction step assumes the claim is true for $n - 1$. Define for $d \in D_n \setminus D_{n-1}$

$$B_d = \frac{B_{d^-} + B_{d^+}}{2} + \frac{Z_d}{2^{(n+1)/2}}$$

where $d^+ = d + 2^{-n}$ and $d^- = d - 2^{-n}$ are the neighbors of d , which are actually members of D_{n-1} . Since $B_{d^-} - B_{d^+} \sim N(0, 1/2^{n-1})$ by the induction hypothesis, then $\frac{B_{d^-} - B_{d^+}}{2} \sim N(0, 1/2^{n+1})$. Furthermore, since $Z_d \sim N(0, 1)$, then $\frac{Z_d}{2^{(n+1)/2}} \sim N(0, 1/2^{n+1})$. Together, these mean $B_d - B_{d^-} = \frac{B_{d^+} - B_{d^-}}{2} + \frac{Z_d}{2^{(n+1)/2}} \sim N(0, 1/2^n)$ and $B_{d^+} - B_d = \frac{B_{d^+} - B_{d^-}}{2} - \frac{Z_d}{2^{(n+1)/2}} \sim N(0, 1/2^n)$. This is enough to establish the first part of the induction claim. Note that B_d depends only on values from D_n so it is independent of all other values in $D \setminus D_n$. This establishes the second part of the induction claim.

Step 2: Using the B_d values, define some functions by linear interpolation between some values in D_n .

- (a) Let $F_0(x) = xZ_1$. Let

$$F_n(x) = \begin{cases} 2^{-(n+1)/2} Z_d & x \in D_n \setminus D_{n-1} \\ 0 & x \in D_{n-1} \\ \text{linear} & \text{between adjacent points in } D_n \end{cases}$$

These functions are continuous on $[0, 1]$. A graph of $F_i(t)$ for $i = 0, \dots, 6$ for some random normal values selected with a pseudo random number generator are in Figure 1.

- (b) Claim: $B_d = \sum_{i=0}^n F_i(d) = \sum_{i=0}^{\infty} F_i(d)$.
- (c) Proof of Claim by Induction: The base case is $n = 0$ with $F_0(0) = 0 = B_0$ and $F_0(1) = Z_1 = B_1$. The induction step supposes the claim holds for $n - 1$. Let $d \in D_n \setminus D_{n-1}$ which is where new values of $\sum_{i=0}^n F_i(d)$ need to be checked. For $0 \leq i \leq n - 1$, F_i is linear on $[d^-, d^+]$ so

$$\sum_{i=0}^{n-1} F_i(d) = \sum_{i=0}^n \frac{F_i(d^-) + F_i(d^+)}{2} = \frac{B_{d^-} + B_{d^+}}{2}.$$

Additionally, $F_n(d) = 2^{-(n+1)/2} Z_d$, so together

$$\sum_{i=0}^n F_i(d) = \frac{B_{d^-} + B_{d^+}}{2} + 2^{-(n+1)/2} Z_d = B_d$$

establishing the claim.

Step 3: Apply the Borel-Cantelli Lemma to show that $|Z_d| \sim c\sqrt{n}$ cannot happen infinitely often.

- (a) Using the estimate for normal random variables in the Lemma, $\mathbb{P}[|Z_d| \geq c\sqrt{n}] \leq \frac{2}{c\sqrt{n}} \frac{1}{\sqrt{2\pi}} e^{-c^2 n/2} < e^{-c^2 n/2}$ for all $n \geq 1$ (as long as $c > \sqrt{\frac{2}{\pi}} \approx 0.798$).
- (b) Then

$$\sum_{n=0}^{\infty} \sum_{d \in D_n} \mathbb{P}[|Z_d| \geq c\sqrt{n}] \leq \sum_{n=0}^{\infty} 2^n e^{-c^2 n/2} = \sum_{n=0}^{\infty} e^{(\log 2 - c^2/2)n}.$$

- (c) So if $c > \sqrt{2 \log 2} \approx 1.177$, the series converges.
- (d) By the Borel-Cantelli Lemma, then there is a random but finite N so that for $n > N$, $d \in D_n$, $|Z_d| \leq c\sqrt{n}$.
- (e) This means $\max_{t \in [0,1]} |F_n(t)| \leq c\sqrt{n} 2^{-n/2}$.
- (f) Then by the Weierstrass M-test, $\sum_{n=0}^{\infty} F_n(t)$ is uniformly and absolutely convergent on $[0, 1]$ to a continuous function, say $B_t = \sum_{n=0}^{\infty} F_n(t)$. A suggestive graph of $\sum_{n=0}^{\infty} F_n(t)$ approximating B_t in in Figure 2.

Step 4: Check that all increments have the correct finite dimensional joint distributions.

- (a) Let $t_1 < t_2 < t_3$ be 3 points in $[0, 1]$. Each is the limit of sequences of dyadics from D , $t_i = \lim_{n \rightarrow \infty} t_{i,n}$ for $i = 1, 2, 3$.
- (b) Then

$$B_{t_3} - B_{t_2} = \lim_{n \rightarrow \infty} (B_{t_{3,n}} - B_{t_{2,n}})$$

is the limit of Gaussian sequences with mean 0 and variance $\lim_{n \rightarrow \infty} (t_{3,n} - t_{2,n}) = t_3 - t_2$. Likewise for $B_{t_2} - B_{t_1}$.

- (c) The sequential increments are independent since $t_{1,n} < t_{2,n} < t_{3,n}$ (for sufficiently large n), and therefore the limit increments $B_{t_3} - B_{t_2}$ and $B_{t_2} - B_{t_1}$ are independent.

Step 5: Finally construct Brownian Motion on $[0, \infty)$.

- (a) By the preceding construction, for $n \geq 1$ let $B_t^{(n)}$ be independent standard Brownian Motion processes on the unit interval.
- (b) Let $B_t = B_{t - [t]}^{([t])} + \sum_{0 \leq i \leq [t]} B_1^{(i)}$

□

Remark. The slopes of the linear parts of $F_n(x)$ are

$$\frac{\pm 2^{-(n+1)/2} Z_d}{1/2^n} = \pm 2^{(n-1)/2} Z_d = O(2^{n/2})$$

which suggests that the limit function $\sum_{i=0}^{\infty} F_i(d)$ will be non-differentiable. See Figure 2. This is not enough to prove non-differentiability, but is suggestive.






Figure 1: Graphs of $F_n(t)$ for $n = 0, \dots, 6$.

Figure 2: Graph of $\sum_{n=0}^6 F_n(t)$ suggesting the form of Brownian Motion and its non-differentiability.

Section Ending Answer

$Z_1 + Z_2$ and $Z_1 - Z_2$ and independent random variables each having the distribution $N(0, 2\sigma^2)$.

Sources

This section is adapted from: *An Invitation to Sample Paths of Brownian Motion*, Chapter 3, by Yuval Peres, An Invitation to Sample Paths of Brownian Motion.



Algorithms, Scripts, Simulations

Algorithm

Scripts

Data: Depth of approximation

Result: Plots of approximation of Brownian Motion

- 1 *Initializations*
- 2 Load interpolation and plotting packages
- 3 Set the depth of approximation
- 4 Set Z_1 and define function F_0
- 5 Create list of dyadic rationals to depth
- 6 *Function Creation* Define the values of F_i on D_i to depth
- 7 Use linear interpolation to create functions F_i
- 8 *Create Plots*
- 9 Plot the functions F_i and save the plot
- 10 Create the Brownian Motion function as a sum of F_i
- 11 Plot the Brownian Motion approximation and save the plot

Scripts

Julia

```
1 using Plots, DataInterpolations
2
3 N = 6
4
5 Z1 = randn()
6 F0 = t -> t * Z1
7
8 D = [ [j//2^i for j=0:2^i] for i = 1:N]
9 Fdata = [ [ iseven(j) ? 0 : randn()/2^((i+1)/2)
10           for j=0:2^i] for i = 1:N]
11 Finterps = [ LinearInterpolation(Fdata[i], D[i
12           ]) for i in 1:N ]
13
14 ts = range(0, 1, length=2^(N+1)+1)
15 F0s = F0.(ts)
16
17 Fs = [Finterps[i].(ts) for i in 1:6]
18 plot(ts, [F0s, Fs[1,:], Fs[2,:], Fs[3,:], Fs
19         [4,:], Fs[5,:], Fs[6,:]], label=false)
20 savefig("individualFs.pdf")
21
22 sumF = F0s + Fs[1] + Fs[2] + Fs[3] + Fs[4] + Fs
23         [5] + Fs[6]
24 plot(ts, sumF, label=false)
25 savefig("LevyBrownianMotion.pdf")
```



Problems to Work for Understanding



Reading Suggestion:

References



Outside Readings and Links:

1. *An Invitation to Sample Paths of Brownian Motion*, by Yuval Peres, An Invitation to Sample Paths of Brownian Motion
2. *Brownian Motion* by Peter Mörters and Yuval Peres, Brownian Motion

Solutions

I check all the information on each page for correctness and typographical errors. Nevertheless, some errors may occur and I would be grateful if you would alert me to such errors. I make every reasonable effort to present current and accurate information for public use, however I do not guarantee the accuracy or timeliness of information on this website. Your use of the information from this website is strictly voluntary and at your risk.

I have checked the links to external sites for usefulness. Links to external websites are provided as a convenience. I do not endorse, control, monitor, or guarantee the information contained in any external website. I don't guarantee that the links are active at all times. Use the links here with the same caution as you would all information on the Internet. This website reflects the thoughts, interests and opinions of its author. They do not explicitly represent official positions or policies of my employer.

Information on this website is subject to change without notice.

Steve Dunbar's Home Page, <http://sdunbarne.github.io>
Email to Steve Dunbar, `sdunbarne` at `gmail dot com`
Last modified: Processed from L^AT_EX source on December 20, 2024